



Sorghum x Sudangrass Root Morphological Changes Affected by Additional N Application while Ratooning

저자 (Authors)	Gamgon Kim, Wonsang Park, Nayoung Choi, Chae-In Na
출처 (Source)	한국작물학회 학술발표대회 논문집 , 2019.4, 88-88(1 pages)
발행처 (Publisher)	한국작물학회 Korean Society Of Crop Science
URL	http://www.dbpia.co.kr/journal/articleDetail?nodeId=NODE08753376
APA Style	Gamgon Kim, Wonsang Park, Nayoung Choi, Chae-In Na (2019). Sorghum x Sudangrass Root Morphological Changes Affected by Additional N Application while Ratooning. 한국작물학회 학술발표대회 논문집 , 88-88
이용정보 (Accessed)	이화여자대학교 203.255.***.68 2020/05/18 04:01 (KST)

저작권 안내

DBpia에서 제공되는 모든 저작물의 저작권은 원저작자에게 있으며, 누리미디어는 각 저작물의 내용을 보증하거나 책임을 지지 않습니다. 그리고 DBpia에서 제공되는 저작물은 DBpia와 구독계약을 체결한 기관소속 이용자 혹은 해당 저작물의 개별 구매자가 비영리적으로만 이용할 수 있습니다. 그러므로 이에 위반하여 DBpia에서 제공되는 저작물을 복제, 전송 등의 방법으로 무단 이용하는 경우 관련 법령에 따라 민, 형사상의 책임을 질 수 있습니다.

Copyright Information

Copyright of all literary works provided by DBpia belongs to the copyright holder(s) and Nurimedia does not guarantee contents of the literary work or assume responsibility for the same. In addition, the literary works provided by DBpia may only be used by the users affiliated to the institutions which executed a subscription agreement with DBpia or the individual purchasers of the literary work(s) for non-commercial purposes. Therefore, any person who illegally uses the literary works provided by DBpia by means of reproduction or transmission shall assume civil and criminal responsibility according to applicable laws and regulations.

PA-54

Sorghum x Sudangrass Root Morphological Changes Affected by Additional N Application while Ratooning

Gamgon Kim¹, Wonsang Park¹, Nayoung Choi¹, Chae-In Na^{1*}

¹Department of Agronomy, Gyeongsang National University, 52828

[Introduction]

Plant root functions as nutrient and water uptake, physical supports. Although there is a direct relationship between the performance of roots and overall crop yield and quality, it is known to be ‘the hidden half of plant’ due to the difficulty of root research. However, there have been significant improvements in root morphological measure recently. Hence, sorghum × sudangrass hybrid (forage sorghum) is a candidate for summer forage and cellulosic biofuel. It has great regrowth potential and excellent biomass yield. The focus of the current research was the root morphological changes affected by summer harvest and following additional N application in the ratooning plant.

[Materials and Method]

The field research was conducted at Gyeongsang National University research farm located in Jinju City, Gyeongnam. The soil was sandy, so it is ideal for massive field root sampling efforts. The plant received 100-81-81 kg/ha fertilizer before planting. Summer harvest was conducted August 10th 2018. The experimental design was a randomized complete block design with a split-plot arrangement. Forage sorghum cultivars; HoneychewBMR (early flowering) and Greenstar (late flowering) was main-plot, and, after summer harvest, additional N level (0, 50, 100, 150 kgN/ha; N0, N50, N100, and N150) was sub-plot. 7 to 10 crown roots were collected twice at the harvests (Oct. 17th and Nov. 21st; Date1 and Date2, respectively). Five representative crown roots from each plot were washed thoroughly then placed specialized scanner for WinRHIZO PRO system (Regent instruments, Canada) to obtain root length, surface area, volume, avg. diameter. Data were statistically analyzed using SAS 9.4.

[Result and discussion]

For early flowering cultivar HoneychewBMR, N100, and N150 showed greater root length, surface area, and volume comparing to N0 and N50 in Date1. In the case of root volume, N100 had avg. of 5.5cm³ which was 2.3-folds greater than N0. However, this different was disappeared in Date2.

For late flowering cultivar Greenstar, N0 had the lowest surface area, and volume compared to the other N treatments in Date1. In the case of root volume, N50 had avg. of 4.9cm³ which was 4.8-folds greater than N0. Unlike early flowering cultivar, N0 had lowest avg. diameter, volume comparing to the other N treatments in Date 2. This is an indication that each cultivar requires a different level of N application for root regrowth.

The Current research showed different belowground characteristics (surface area, volume, length, and avg. diameter) changes to increasing N level in two cultivars. Thus, root morphology can attribute to evaluating plant growth in field based research.

[Acknowledgements]

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (No. NRF-2017R1C1B5016480).

*Corresponding author: Tel. +82-55-772-1878, E-mail, nachaein@gnu.ac.kr