

V5A4 – HABIRO COHOMOLOGY

SUMMER SEMESTER 2025

WERN JUIN GABRIEL ONG

PRELIMINARIES

These notes roughly correspond to the course **V5A4 – Habiro Cohomology** taught by Prof. Peter Scholze at the Universität Bonn in the Summer 2025 semester. These notes are \LaTeX -ed after the fact with significant alteration and are subject to misinterpretation and mistranscription. Use with caution. Any errors are undoubtedly my own and any virtues that could be ascribed to these notes ought be attributed to the instructor and not the typist. Recordings of the lecture are available at the following link:

archive.mpim-bonn.mpg.de/id/eprint/5155/

CONTENTS

Preliminaries	1
1. Lecture 1 – 11th April 2025	2
2. Lecture 2 – 2nd May 2025	8
3. Lecture 3 – 9th May 2025	13
References	17

1. LECTURE 1 – 11TH APRIL 2025

Recall that the construction of the Habiro ring of a number field [GS+24, Sch24] was motivated by an expectation of the instructor, circa 2017, that there exists some form of “Habiro cohomology.” Within this larger aspirational framework, the Habiro ring of a number field serves as the zero-dimensional case where the variety is a discrete collection of points. More precisely, in the case of the Habiro ring of a number field, there are certain q -series related to perturbative Chern-Simons theory giving rise to an explicit approach to Habiro rings of number fields. In particular, these q -series from perturbative Chern-Simons theory as computed by Garoufalidis and Zagier arise as elements of the abstract Habiro ring of a number field.

The goal of this course, then, is to explicate this aspirational framework of Habiro cohomology that synthesizes the concrete approach of Garoufalidis-Zagier with the instructor’s abstract approach. In particular, we will define a new explicit cohomology theory for algebraic varieties that has specializations to classical cohomology theories: de Rham cohomology as well as p -adic étale cohomology, crystalline cohomology, and prismatic cohomology for all primes p . Moreover, this cohomology theory will extend to the rigid-analytic setting of Berkovich spaces.

Let recall a modern definition of Weil-type cohomology theories for algebraic varieties: functors

$$\mathrm{Sch}_k^{\mathrm{sft}} \longrightarrow \mathrm{Pr}_A^{\mathrm{L}}$$

where $\mathrm{Sch}_k^{\mathrm{sft}}$ is the category of separated finite type schemes over k and $\mathrm{Pr}_A^{\mathrm{L}}$ the category of presentable A -linear categories with a six-functor formalism and satisfying the Künneth formula. In particular this excludes some cohomology theories such as motivic cohomology.

The state of the art of Weil-type cohomology theories for algebraic varieties can be summarized in the following diagram.

The instructor remarks that this is his favorite diagram.




FIGURE 1. Cohomology theories for algebraic varieties. Or: the instructor’s favorite diagram.

- Betti cohomology $X \mapsto \mathcal{D}(X(\mathbb{C}), \mathbb{Z}) \otimes (-)$ produces a cohomology theory for complex schemes. But coefficients can be taken in any field by base change.
- de Rham cohomology $X \mapsto \mathbf{DMod}(X)$ associating to a scheme its category of D -modules produces a cohomology theory for k -schemes (modulo technicalities). This produces a k -vector space for a k -scheme, hence has coefficients equal to the characteristic of the scheme.
- Étale cohomology as defined by Grothendieck $X \mapsto \mathcal{D}_{\text{ét}}(X_{\text{ét}}, \mathbb{Z}/\ell^n \mathbb{Z})$ produces for a k -scheme X , a cohomology theory with $\mathbb{Z}/\ell^n \mathbb{Z}$ -coefficients with ℓ of characteristic distinct from that of k . That étale cohomology is able to produce cohomology in coefficients modulo powers of ℓ is represented by the thickening of the horizontal. Note that étale cohomology satisfies the Künneth formula, but not its categorical variant.
- Crystalline cohomology after Grothendieck, Berthelot, Caro, et. al. that associates to a k -scheme where k is of positive characteristic a cohomology theory $X \mapsto \mathbf{DMod}(X)$ that associates to X its category of arithmetic D -modules and which satisfies the categorical Künneth formula. This produces a module over the Witt vectors $W(k)$ of k for a k -scheme, and is represented by vertical thickenings at the characteristic.
- Prismatic cohomology was defined by Bhatt-Scholze [BS22] as a universal cohomology theory at the (p, p) -point by computing the structure sheaf cohomology of the prismatic site $X \mapsto R\Gamma_{\Delta}(X)$ where X is a scheme over \mathcal{O}_K where K is a mixed characteristic local field which has coefficients valued in prisms.¹

FIGURE 2. Prismatic cohomology at the (p, p) -point.

Moreover, the diagram reflects several important comparison phenomena between the abovementioned cohomology theories:

¹It would be more precise to state this using “derived category of sheaves” associated to prismatic cohomology, namely the category of F -gauges à la Bhatt-Lurie [Bha22], but we do not comment on this further.

- The intersection of the lines corresponding to Betti and de Rham cohomology at the (∞, ∞) -point is substantiated by the comparison isomorphism between singular cohomology with \mathbb{C} -coefficients and de Rham cohomology via the Riemann-Hilbert correspondence.
- The intersection of the lines corresponding to étale and Betti cohomology at the (∞, p) -points are substantiated by the Artin's comparison isomorphism between étale and Betti cohomology.
- The intersection of the thickenings of crystalline cohomology meeting de Rham cohomology along the diagonal at the (p, p) -point is substantiated by the isomorphism between crystalline cohomology reduced modulo p and de Rham cohomology.
- Prismatic cohomology as depicted in Figure 1 admits specializations to de Rham, crystalline, and étale cohomology. Prismatic cohomology is additionally compatible with the structures of the various cohomology theories around the (p, p) -point, specializing to the action of the Frobenius in crystalline cohomology, the Hodge-Tate filtration in the case of de Rham cohomology, and the action of the absolute Galois group $\text{Gal}(\overline{\mathbb{Q}_p}/\mathbb{Q}_p)$ in the case of étale cohomology.
- The “prismatization” at the (∞, ∞) -point is the content of classical complex Hodge theory, which considers Hodge filtrations on de Rham cohomology and associated objects.

Observe, then, that de Rham cohomology is the unifying cohomology theory on the diagonal, while prismatic cohomology only exists at a fixed prime. One then wonders if there is a way to unify the cohomology theories along the diagonal. This is provided by Habiro cohomology, at least in the positive characteristic case.

The instructor remarks that he is unsure how to unify Habiro cohomology with classical Hodge theory.



FIGURE 3. The role of Habiro cohomology highlighted in blue generalizing prismatic cohomology at all primes. Compare Figure 1.

That is to say that Habiro cohomology, covering a neighborhood of the de Rham diagonal, specializes to prismatic cohomology at each prime, and spreads out further than prismatic cohomology along the horizontal étale branches in an appropriate sense.

The starting point of Habiro cohomology is the example of the q -de Rham prism, the definition of which we now recall.

Example 1.1. The q -de Rham prism is the prism $(\mathbb{Z}_p[[q-1]], [p]_q)$ where $[p]_q = \frac{1-p^n}{1-q}$ is the q -deformation of p with a Frobenius action by $q \mapsto q^p$. The quotient $\mathbb{Z}_p[[q-1]]/([p]_q)$ is precisely the quotient by the p -th cyclotomic polynomial and hence isomorphic to the cyclotomic extension $\mathbb{Z}_p[\zeta_p]$.

Computing the prismatic cohomology of $\mathbb{A}_{\mathbb{Z}_p[\zeta_p]}^1$ relative to the q -de Rham prism, one finds that this is computed by an obvious q -deformation of the de Rham complex. The cohomological comparisons of the preceding discussion suggest that there is a deformation of the de Rham complex given by

$$\nabla_q : \mathbb{Z}_p[\zeta_p][x][[q-1]] \longrightarrow \mathbb{Z}_p[\zeta_p][x][[q-1]]$$

by $x^n \mapsto [n]_q x^{n-1}$. It is not *a priori* clear why q -deformations appear in this setting. Moreover, the construction of prismatic cohomology over the q -de Rham prism is expected to be functorial in automorphisms of $\mathbb{A}_{\mathbb{Z}_p[\zeta_p]}^1$ but it is unclear if (and how) this construction is invariant under change of coordinates. Additionally, the q -deformation suggests that by removing p everywhere, one can find a construction independent that works for all primes p . In particular, the instructor conjectures in [Sch17] the following:

Conjecture 1.2 (Scholze; [Sch17, Conj. 1.1]). If R is a smooth \mathbb{Z} -algebra equipped with an étale map $\mathrm{Spec}(R) \rightarrow \mathbb{A}_{\mathbb{Z}}^d$, there is a cohomology theory for smooth proper varieties over R valued in finitely generated $R[[q-1]]$ -modules with a q -connection.

The q -connection captures precisely the difficulties with coordinate transformations articulated above, and the specialization at $q = 1$ recovers the de Rham cohomology of X with a Gauss-Manin connection. This suggests that algebraic varieties have a canonical q -deformation with connection compatible with the Gauss-Manin connection on classical de Rham cohomology, and was proven after p -adic completion in [BS22] and in general by Ferdinand Wagner in [Wag24] using the machinery of adelic gluing.

Theorem 1.3 (Wagner; [Wag24, Thm. 1.7]). Let R be a smooth framed \mathbb{Z} -algebra. There is an isomorphism between the $(q-1)$ -completed q -de Rham–Witt complex and the cohomology of the quotient of the q -Hodge complex by $(q^m - 1)$.

Let us consider an example of this phenomenon.

Example 1.4. Consider the Legendre family of elliptic curves X with affine model $y^2 = x(x-1)(x-\lambda)$ over $R = \mathbb{Z}[\frac{1}{2}, \lambda, \frac{1}{\lambda(1-\lambda)}]$. We have $H_{\mathrm{dR}}^1(X)$ free of rank 2, containing the Hodge filtration $\mathrm{Fil}_{\mathrm{Hdg}}^1 = H^0(X, \Omega_{X/R}^1)$ with canonical differential

$\omega = \frac{dx}{y}$. Denoting ∇ the connection on $H_{\text{dR}}^1(X)$, we have $\omega, \nabla(\omega)$ a basis of $H_{\text{dR}}^1(X)$ and

$$\nabla^2(\omega) = \frac{1}{4\lambda(1-\lambda)} + \frac{2\lambda-1}{\lambda(1-\lambda)}\nabla(\omega).$$

A horizontal section is $f(\lambda) \cdot \lambda(1-\lambda) - f'(\lambda)\lambda(1-\lambda)\nabla(\omega)$ for a certain hypergeometric function $f(\lambda) = \sum_{n \geq 0} \prod_{i=0}^{n-1} \left(\frac{i+\frac{1}{2}}{i+1} \right)^2 \lambda^n$.

There is a q -analogue of hypergeometric functions.

Example 1.5. The q -hypergeometric function

$$\sum_{n \geq 0} \prod_{i=0}^{n-1} \left(\frac{[i + \frac{1}{2}]_q}{[i+1]_q} \right)^2 \lambda^n$$

satisfies a second order q -difference equation that deforms the Picard-Fuchs equation whose solutions describe periods of elliptic curves [nLab-a].

The example suggests that there is a possible connection between q -hypergeometric functions – the q -analogue of hypergeometric functions – and q -deformations of de Rham cohomology.

In the case of de Rham cohomology as in Example 1.4, there is not only a connection ∇ , but also a choice of canonical vector $\omega = \frac{dx}{y}$ obtained by the filtration. Then considering the differential equation the class satisfies produces the desired differential equation – the module and connection alone are insufficient to produce the differential equation. The main barrier to considering the q -analogue, then, was the lack of choice of such a class.

Recent computations of Shirai [Shi20] and work of Garoufalidis-Wheeler remedy this by producing explicit classes in q -de Rham cohomology, allowing the procedure above to be repeated.

This course will consider what happens to these q -deformations when q approaches a root of unity ζ_m , knowing that it recovers the classical construction at $q = 1$. Working over the Habiro ring

$$\mathcal{H} = \lim_{m,n \geq 1} \mathbb{Z}[q]/(1-q^n)^m = \lim_n \mathbb{Z}[q]/(q; q)_n$$

allows us to consider specializations at different roots of unity.

One issue that arises in trying to naïvely generalize Habiro cohomology to schemes of higher dimension is that the specialization of prismatic cohomology over the q -de Rham prism at $q = 1$ recovers de Rham cohomology, but at other roots of unity recovers only Hodge cohomology – this does not put all roots of unity on equal footing. But if the q -de Rham cohomology could be modified to be Hodge cohomology in an appropriate manner. This was shown by Meyer-Wagner in [MW24].

Theorem 1.6 (Meyer-Wagner; [MW24, Thm. 1.7]). Let R be a p -torsion free p -complete ring which is a quasiregular quotient over \mathbb{Z}_p and such that the Frobenius on R/p is semiperfect. If R admits a lift to a p -complete \mathbb{E}_1 ring spectrum \mathbb{S}_R such that $R \simeq \mathbb{S}_R \widehat{\otimes}_{\mathbb{S}_p} \mathbb{Z}_p$ then the q -Hodge filtration on the p -complete derived

q -de Rham complex is a q -deformation of the Hodge filtration on the (ordinary) p -complete derived de Rham complex.

The proof of Meyer-Wagner once again leverages highly technical machinery, in particular the relationship between prismatic cohomology and topological cyclic homology. However, there is a more computational way of achieving the same goal.

Theorem 1.7 (Scholze). There is an explicit ring stack over an analytic version of the Habiro ring yielding a full six-functor formalism.

Remark 1.8. This in particular yields a sheaf theory.

These are related to the constructions of the ring stacks for prismatic cohomology following Drinfeld [Dri20] and Bhatt-Lurie [BL22].

Here “ring stack” and “analytic” are to be taken in the sense of condensed mathematics [CS23].

While multiplication is easy to define in this ring, addition is not: in particular, the instructor remarks that he spent a whole day computing what $1 + 1$ is in this ring.

2. LECTURE 2 – 2ND MAY 2025

The goal of this course is to develop a theory of Habiro cohomology, a functor that associates to a smooth \mathbb{Z} -scheme X its Habiro cohomology – a module over the Habiro ring, or more generally its “category of constructible sheaves” which in this case we tentatively denote $\mathcal{D}_{\text{Hab}}(X)$ of “variations of Habiro structure.”

We begin with an exploration of what these structures are in terms of coordinates, and we will later show that the constructions we discuss are in fact independent of these coordinates. Let us make the notion of coordinates precise.

Definition 2.1 (Framed Algebra). A framed algebra is a pair (R, \square) where R is a smooth \mathbb{Z} -algebra and an étale map $\square : \text{Spec}(R) \rightarrow \mathbb{A}_{\mathbb{Z}}^d$ or $\square : \text{Spec}(R) \rightarrow \mathbb{G}_m^d$.

Remark 2.2. It is often simpler to consider the case where the coordinates are invertible, that is, the case of \mathbb{G}_m^d .

As a first pass, let us contemplate these constructions in the case where X is affine and equal to either $\mathbb{A}_{\mathbb{Z}}^d$ or \mathbb{G}_m^d and only later consider the generalization to the case where X is étale over one of these spaces. Moreover, under these assumptions, we need not make any completions and one can work over $\mathbb{Z}[q^{\pm}]$.

Recall Habiro cohomology subsumes de Rham cohomology in an appropriate sense, and takes the q -derivative – the Gaussian q -analogue of the derivative – as an input. These q -derivatives were first investigated by Jackson [Jac10].

Definition 2.3 (q -Derivative). Let R be $\mathbb{Z}[q^{\pm}][T_1, \dots, T_d]$ or $\mathbb{Z}[q^{\pm}][T_1^{\pm}, \dots, T_d^{\pm}]$. The q -derivative $\nabla_i^q : R \rightarrow R$ for $1 \leq i \leq d$ is defined by

$$\nabla_i^q(f(T_1, \dots, T_d)) = \frac{f(T_1, \dots, qT_i, \dots, T_d) - f(T_1, \dots, T_i, \dots, T_d)}{qT_i - T_i}.$$

Remark 2.4. More explicitly, this operation is given on monomials by

$$\nabla_i^q(T_1^{n_1} \dots T_d^{n_d}) = [n_i]_q \cdot T_1^{n_1} \dots T_i^{n_i-1} \dots T_d^{n_d}$$

where $[n]_q = \frac{1-q^n}{1-q}$ is the Gaussian q -analogue of n .

Remark 2.5. ∇_i^q is closely related $\gamma_i : R \rightarrow R$ the automorphism by

$$T_j \mapsto \begin{cases} T_j & j \neq i \\ qT_i & j = i \end{cases}$$

allowing us to write $\nabla_i^q(f) = \frac{\gamma_i(f) - f}{(q-1)T_i}$.

The q -derivative does not satisfy the Leibniz rule on the nose, but does so up to a twist by the automorphism γ_i of Remark 2.5.

Lemma 2.6. Let R be $\mathbb{Z}[q^{\pm}][T_1, \dots, T_d]$ or $\mathbb{Z}[q^{\pm}][T_1^{\pm}, \dots, T_d^{\pm}]$. Then for $f, g \in R$ we have equalities

$$\nabla_i^q(fg) = \gamma_i(f) \cdot \nabla_i^q(g) + g \cdot \nabla_i^q(f) = f \cdot \nabla_i^q(g) + \gamma_i(g) \cdot \nabla_i^q(f).$$

Proof. We first show the second equality. We use Remark 2.5 to observe that the latter two terms are given by

$$\gamma_i(f) \cdot \frac{\gamma_i(g) - g}{(q-1)T_i} + g \cdot \frac{\gamma_i(f) - f}{(q-1)T_i} = \frac{\gamma_i(f)\gamma_i(g) - \gamma_i(f)g + \gamma_i(f)g - fg}{(q-1)T_i} = \frac{\gamma_i(f)\gamma_i(g) - fg}{(q-1)T_i}$$

and

$$f \cdot \frac{\gamma_i(g) - g}{(q-1)T_i} + \gamma_i(g) \frac{\gamma_i(f) - f}{(q-1)T_i} = \frac{\gamma_i(g)f - fg + \gamma_i(f)\gamma_i(g) - \gamma_i(g)f}{(q-1)T_i} = \frac{\gamma_i(f)\gamma_i(g) - fg}{(q-1)T_i}$$

respectively, which are evidently equal.

We now show the first equality. Note that γ_i is an automorphism $R \rightarrow R$, and in particular a homomorphism so $\gamma_i(fg) = \gamma_i(f)\gamma_i(g)$ in which case we have

$$\frac{\gamma_i(fg) - fg}{(q-1)T_i} = \frac{\gamma_i(f)\gamma_i(g) - fg}{(q-1)T_i}$$

whence the claim. ■

We can now define the q -de Rham complex following Aomoto [Aom90].

Definition 2.7 (q -de Rham Complex of $\mathbb{A}_{\mathbb{Z}}^d$ and \mathbb{G}_m^d). Let $R = \mathbb{Z}[q^{\pm}][T]$ be $\mathbb{Z}[q^{\pm}][T_1, \dots, T_d]$ or $\mathbb{Z}[q^{\pm}][T_1^{\pm}, \dots, T_d^{\pm}]$. The q -de Rham complex of $\text{Spec}(R)$ is the complex

$$(2.1) \quad \begin{aligned} 0 &\longrightarrow \mathbb{Z}[q^{\pm}][T] \longrightarrow \mathbb{Z}[q^{\pm}][T]^{\oplus d} \longrightarrow \bigoplus_{i < j} \mathbb{Z}[q^{\pm}][T] \longrightarrow \dots \\ &\dots \longrightarrow \mathbb{Z}[q^{\pm}][T] \longrightarrow 0 \end{aligned}$$

with differentials given by the differentials for the Koszul complex of commuting operators $\nabla_1^q, \dots, \nabla_n^q$.

Remark 2.8. Recall that these are precisely the differentials for the classical de Rham complex. See [Stacks, Tag 0FKF] for an explicit description via equations.

Remark 2.9. Since the first differential $\mathbb{Z}[q^{\pm}][T] \rightarrow \mathbb{Z}[q^{\pm}][T]^{\oplus d}$ by $(\nabla_1^q, \dots, \nabla_d^q)$ does not satisfy the ordinary Leibniz rule, the complex (2.1) is not a differential graded algebra. Later, we will see that working in the derived (∞) -category, one can endow this with the structure of a commutative ring.

The complex (2.1) computes q -de Rham cohomology, or Aomoto-Jackson cohomology of $\text{Spec}(R)$. But to compute Habiro cohomology, we use a closely related variant based on a modified q -derivative.

Definition 2.10 (Modified q -Derivative). Let R be $\mathbb{Z}[q^{\pm}][T_1, \dots, T_d]$ or $\mathbb{Z}[q^{\pm}][T_1^{\pm}, \dots, T_d^{\pm}]$. The modified q -derivative is given by

$$\tilde{\nabla}_i^q(f(T_1, \dots, T_d)) = \frac{f(T_1, \dots, qT_i, \dots, T_d) - f(T_1, \dots, T_i, \dots, T_d)}{T_i}.$$

Remark 2.11. In other words, $\tilde{\nabla}_i^q(f) = (q-1)\nabla_i^q(f) = \frac{\gamma_i(f) - f}{T_i}$.

Recomputing everything using this modified derivative gives the q -Hodge complex.

Definition 2.12 (q -Hodge Complex of $\mathbb{A}_{\mathbb{Z}}^d$ and \mathbb{G}_m^d). Let $R = \mathbb{Z}[q^{\pm}][T]$ be $\mathbb{Z}[q^{\pm}][T_1, \dots, T_d]$ or $\mathbb{Z}[q^{\pm}][T_1^{\pm}, \dots, T_d^{\pm}]$. The q -Hodge complex of $\text{Spec}(R)$ is the complex

$$(2.2) \quad \begin{aligned} 0 \longrightarrow \mathbb{Z}[q^{\pm}][T] &\longrightarrow \mathbb{Z}[q^{\pm}][T]^{\oplus d} \longrightarrow \bigoplus_{i < j} \mathbb{Z}[q^{\pm}][T] \longrightarrow \dots \\ &\dots \longrightarrow \mathbb{Z}[q^{\pm}][T] \longrightarrow 0 \end{aligned}$$

with differentials given by the differentials for the Koszul complex of commuting operators $\tilde{\nabla}_1^1, \dots, \tilde{\nabla}_d^q$.

Remark 2.13. The nomenclature of Definitions 2.7 and 2.12 are justified by the fact that they recover the ordinary de Rham and Hodge complexes at $q = 1$.

Remark 2.14. An automorphism of $\mathbb{A}_{\mathbb{Z}}^d$ or \mathbb{G}_m^d would give rise to an automorphism of the complexes (2.1) and (2.2), at least as an object in the derived category, but it is extremely difficult to understand these automorphisms from this explicit perspective.

The instructor remarks that he does not believe in non-flat connections. We will henceforth omit the adjective “flat.”

In parallel to the correspondence between algebraic D -modules and modules with flat connection, one would expect the existence of a category of modules with an appropriate connection to play the role of $\mathcal{D}_{\text{Hab}}(X)$ alluded to earlier. To make this precise, we consider modules with q -connection. To simplify matters, we make these considerations on the Abelian and not ∞ -categorical level.

Definition 2.15 (q -Connections on Modules). Let $R = \mathbb{Z}[q^{\pm}][T]$ be $\mathbb{Z}[q^{\pm}][T_1, \dots, T_d]$ or $\mathbb{Z}[q^{\pm}][T_1^{\pm}, \dots, T_d^{\pm}]$. A module with (flat) q -connection is a $\mathbb{Z}[q^{\pm}][T]$ -module with commuting $\mathbb{Z}[q^{\pm}]$ -linear operations $\nabla_{i,M}^q : M \rightarrow M$ which satisfy the q -Leibniz rule

$$\nabla_{i,M}^q(fm) = \gamma_i(f) \cdot \nabla_{i,M}^q(m) + \nabla_i^q(f) \cdot m$$

for $f \in \mathbb{Z}[q^{\pm}][T]$ and $m \in M$.

Remark 2.16. To unwind any possible confusion between the similar-looking $\nabla_i^q : \mathbb{Z}[q^{\pm}][T] \rightarrow \mathbb{Z}[q^{\pm}][T], \nabla_{i,M}^q : M \rightarrow M$, we have

$$\underbrace{\underbrace{\gamma_i(f)}_{\in \mathbb{Z}[q^{\pm}][T]} \cdot \underbrace{\nabla_{i,M}^q(m)}_{\in M}}_{\in M} + \underbrace{\underbrace{\nabla_i^q(f)}_{\in \mathbb{Z}[q^{\pm}][T]} \cdot \underbrace{m}_{\in M}}_{\in M}$$

so everything type-checks.

Example 2.17. If $X = \mathbb{A}_{\mathbb{Z}}^1$ then recall that modules with connection are equivalent to modules over the Weyl algebra $\mathbb{Z}[q^{\pm}]\{T, \partial_q\}/(qT\partial_q - \partial_q T + 1)$ since we have the operators $T\partial_q, \partial_q T$ take T^n to $q[n]_q T^n, [n+1]_q T^n$, respectively, but $q[n]_q - [n+1]_q = q \cdot \frac{1-q^n}{1-q} - \frac{1-q^{n+1}}{1-q} = -1$. Passing to the associated-graded of the degree filtration, one gets commuting variables with the correct q -twists.

Similarly, we can construct modules with a modified q -connection.

Definition 2.18 (Modified q -Connections on Modules). Let $R = \mathbb{Z}[q^\pm][T]$ be $\mathbb{Z}[q^\pm][T_1, \dots, T_d]$ or $\mathbb{Z}[q^\pm][T_1^\pm, \dots, T_d^\pm]$. A module with modified q -connection is a $\mathbb{Z}[q^\pm][T]$ -module with commuting $\mathbb{Z}[q^\pm]$ -linear operations $\tilde{\nabla}_{i,M}^q : M \rightarrow M$ which satisfy the q -Leibniz rule

$$\tilde{\nabla}_{i,M}^q(fm) = \gamma_i(f) \cdot \tilde{\nabla}_{i,M}^q(m) + \tilde{\nabla}_i^q(f) \cdot m$$

for $f \in \mathbb{Z}[q^\pm][T]$ and $m \in M$.

Remark 2.19. For a more in-depth discussion of modules with q -connection, see Morrow-Tsuiji [MT21] and André [And01].

Remark 2.20. Let T_i be invertible. Unwinding the definition of the modified q -derivative, we have

$$\tilde{\nabla}_{i,M}^q(fm) = \gamma_i(f) \cdot \tilde{\nabla}_{i,M}^q(m) + (q-1)\nabla_i^q(f) \cdot m$$

where in particular we observe that the second summand has denominator T_i . Define a new operator

$$\tilde{\tilde{\nabla}}_{i,M}^q = T_i \cdot \tilde{\nabla}_{i,M}^q$$

which satisfies

$$\begin{aligned} \tilde{\tilde{\nabla}}_{i,M}^q(fm) &= \gamma_i(f) \cdot \tilde{\tilde{\nabla}}_{i,M}^q(m) + (\gamma_i(f) - f)m \\ &= \gamma_i(f) \left(\tilde{\tilde{\nabla}}_{i,M}^q(m) + m \right) - fm. \end{aligned}$$

In particular,

$$\left(\tilde{\tilde{\nabla}}_{i,M}^q + \text{id}_M \right) (fm) = \gamma_i(f) \left(\tilde{\tilde{\nabla}}_{i,M}^q + \text{id}_M \right) (m)$$

so denoting $\gamma_{i,M} = \left(\tilde{\tilde{\nabla}}_{i,M}^q + \text{id}_M \right)$, we have $\gamma_{i,M}(fm) = \gamma_i(f)\gamma_{i,M}(m)$ simplifying the relation.

The preceding discussion of Remark 2.20 implies the following.

Corollary 2.21. Let $R = \mathbb{Z}[q^\pm][T_1^\pm, \dots, T_d^\pm]$. There is an equivalence of categories between R -modules with modified q -connection and R -modules with commuting $\gamma_i : R \rightarrow R$ -semilinear endomorphisms $\gamma_{i,M} : M \rightarrow M$.

Note that for $R = \mathbb{Z}[q^\pm][T]$, $(-) \otimes_R (-)$ does not define a symmetric monoidal structure on the category of modules with q -connection: for $(M, \nabla_{i,M}^q), (N, \nabla_{i,N}^q)$ two modules with q -connection,

$$(M \otimes_R N, \nabla_{i,M}^q \otimes_R \text{id}_N + \text{id}_M \otimes_R \nabla_{i,N}^q)$$

is not a module with q -connection. One needs instead to take the twist

$$(M \otimes_R N, \nabla_{i,M}^q \otimes_R \text{id}_N + \gamma_{i,M} \otimes_R \nabla_{i,N}^q),$$

defining $\gamma_{i,M} : M \rightarrow M$ in an analogous way to Remark 2.20. While *a priori* appearing assymetric in M, N , there is in fact a canonical isomorphism between them.

Proposition 2.22. Let $R = \mathbb{Z}[q^\pm][T_1^\pm, \dots, T_d^\pm]$. The category of R -modules with q -connection is symmetric monoidal.

Proof Outline. Using the equivalence of Corollary 2.21, the latter category is symmetric monoidal, hence the former can be promoted to a symmetric monoidal category. ■

Proposition 2.23. Let $R = \mathbb{Z}[q^\pm][T_1^\pm, \dots, T_d^\pm]$. There is a fully faithful embedding from $(q-1)$ -torsion free R -modules with q -connection and R -modules with modified q -connection by $(M, \nabla_{i,M}^q) \mapsto (M, \tilde{\nabla}_{i,M}^q)$ with essential image those that are $(q-1)$ -torsion free and such that $\tilde{\nabla}_{i,M}^q \equiv 0 \pmod{(q-1)}$.

The discussion thus far has been done entirely in terms of coordinates. This prompts:

Question 2.24. To what extent are the cohomologies and categories discussed thus far independent of coordinates?

Let us consider the following example.

Example 2.25. Let $X = \mathbb{G}_m^d$. The modules with modified q -connection are quasicoherent sheaves on $(\mathbb{G}_m/q^\mathbb{Z})^d$ – the γ_i 's act by multiplication by q on the coordinates so the data of the endomorphisms $\gamma_{i,M}$ on the modules prescribe descent data to the quotient stack (ie. as an fpqc quotient).

Let us relate the discussion of complexes Definitions 2.7 and 2.12, their cohomologies, and these categories of modules with (modified) q -connections.

Proposition 2.26. (i) The q de Rham complex computes $R\mathrm{Hom}(\mathbb{1}, \mathbb{1})$ in the derived category of modules with q -connection on $\mathrm{Spec}(R)$.
(ii) The q -Hodge complex computes $R\mathrm{Hom}(\mathbb{1}, \mathbb{1})$ in the derived category of modules with modified q -connection on $\mathrm{Spec}(R)$.

Proof Outline of (i). Using the equivalence between modules with q -connection and modules over the Weyl algebra, we compute a resolution of the symmetric monoidal unit $\mathbb{Z}[q^\pm][T]$ in the category of modules over the Weyl algebra – which precisely recovers the de Rham complex, whence the claim. ■

Example 2.27. Consider the case of $\mathbb{A}_{\mathbb{Z}}^1$ taking $R = \mathbb{Z}[q^\pm][T]$. We compute $R\mathrm{Hom}(\mathbb{Z}[q^\pm][T], \mathbb{Z}[q^\pm][T])$ as $R\mathrm{Hom}(-, \mathbb{Z}[q^\pm][T])$ of a free resolution of $\mathbb{Z}[q^\pm][T]$ in the category of modules over the Weyl algebra $\mathbb{Z}[q^\pm]\{T, \partial_q\}/(qT\partial_q - \partial_q T + 1)$ (vis. Example 2.17). This produces

$$0 \rightarrow \mathbb{Z}[q^\pm][T] \xrightarrow{\nabla_1^q} \mathbb{Z}[q^\pm][T] \rightarrow 0$$

which is the q -de Rham complex (after passing back to modules with q -connection along the equivalence).

Moreover, in the setting of higher algebra, these promote canonically to commutative algebra objects.

Corollary 2.28. The q -de Rham complex and q -Hodge complex have canonical structures as \mathbb{E}_∞ -rings.

The instructor remarks that in the theory of analytic geometry the quotient would be the Tate elliptic curve for $d = 1$. See [CS23].

3. LECTURE 3 – 9TH MAY 2025

In Section 2, we constructed the q -de Rham and q -Hodge complexes for $\mathbb{A}_{\mathbb{Z}}^d, (\mathbb{G}_m)^d$ using the q -derivatives ∇_i^q and modified q -derivatives $\widehat{\nabla}_i^q$, respectively.

We now consider the construction of the q -de Rham and q -Hodge complexes more generally in the case where the T_i are invertible and using the logarithmic q -derivative $\nabla_i^{q, \log} = T_i \nabla_i^q$. In particular, we define the logarithmic $\nabla_i^{q, \log}$ and modified logarithmic q -derivative $\widehat{\nabla}_i^{q, \log}$ by

$$f \mapsto \frac{\gamma_i(f) - f}{q - 1}, f \mapsto \gamma_i(f) - f,$$

respectively.

Remark 3.1. The commutation relation for the ordinary logarithmic q -derivative are given by $\gamma_i T_i = q T_i \gamma_i$ since multiplying by T_i and applying the map $T_i \mapsto q T_i$ is the same as applying the map $T_i \mapsto q T_i$ and multiplying by $q T_i$.

For simplicity, take $X = \text{Spec}(R)$ to be affine for some smooth \mathbb{Z} -algebra R and $\square : X \rightarrow (\mathbb{G}_m)^d$ its (étale) framing induced by an étale ring map $\mathbb{Z}[T_1^{\pm}, \dots, T_d^{\pm}] = \mathbb{Z}[\underline{T}^{\pm}] \rightarrow R$. Naïvely we want to define a category of $R[q^{\pm}]$ -modules with commuting semilinear endomorphisms $\gamma_{i, M} : M \rightarrow M$ for each such module M . However, such semilinearity first requires a choice of $\gamma_{i, R} : R[q^{\pm}] \rightarrow R[q^{\pm}]$ extending the map $\gamma_i : \mathbb{Z}[q^{\pm}][\underline{T}^{\pm}] \rightarrow \mathbb{Z}[q^{\pm}][\underline{T}^{\pm}]$ but such a map need not exist. Geometrically speaking, the automorphisms γ_i on $(\mathbb{G}_m)^d$ need not lift along the map $\square : X \rightarrow (\mathbb{G}_m)^d$. This is where $(q - 1)$ -adic completion enters the picture: after $(q - 1)$ -adic completion, there are unique such $\gamma_{i, R} : R[[q - 1]] \rightarrow R[[q - 1]]$ which reduce to the identity modulo $(q - 1)$ induced by the infinitesimal lifting property of étale maps.

To wit, there is an equivalence of categories

$$\left\{ \begin{array}{c} \text{étale } \mathbb{Z}[q^{\pm}][\underline{T}^{\pm}]/(q-1)^n \\ \text{algebras} \end{array} \right\} \simeq \left\{ \begin{array}{c} \text{étale } \mathbb{Z}[\underline{T}^{\pm}] \\ \text{algebras} \end{array} \right\}.$$

On étale $\mathbb{Z}[q^{\pm}][\underline{T}^{\pm}]/(q - 1)^n$ -algebras, there are endomorphisms γ_i^* induced by $\gamma_i : \mathbb{Z}[q^{\pm}][\underline{T}^{\pm}] \rightarrow \mathbb{Z}[q^{\pm}][\underline{T}^{\pm}]$ which reduce to the identity on $\mathbb{Z}[\underline{T}^{\pm}]$ -algebras. In particular, the identity map on a $\mathbb{Z}[q^{\pm}][\underline{T}^{\pm}]/(q - 1)^n$ -algebra and γ_i^* are both lifts of the identity along the equivalence, showing that γ_i^* is the identity by functoriality. Thus, after $(q - 1)$ -adic completion, the automorphisms γ_i on $(\mathbb{G}_m)^d$ are infinitesimally close to 1, and hence lift uniquely along the framing map \square , and allowing us to define q -derivatives and the categories of modules with (modified) q -connection.

Theorem 3.2 (Bhatt-Scholze, [BS22, §16]; Wagner, [Wag24, Thm. 1.5]). Let (R, \square) be a smooth framed \mathbb{Z} -algebra. The complex $q\Omega_{(R, \square)/\mathbb{Z}[[q - 1]]}$ given by

$$R[[q - 1]] \xrightarrow{(\nabla_i^q)_{i=1}^d} \bigoplus_{i=1}^d R[[q - 1]] \longrightarrow \dots$$

as an object of $\mathcal{D}(\mathbb{Z}[[q - 1]])$ is canonically independent of the choice of coordinates.

Example 3.3. Consider the case of a smooth framed \mathbb{Q} -algebra (R, \square) where $\square : \text{Spec}(R) \rightarrow \mathbb{G}_m$. We can use Taylor's theorem to write

$$f(qT) = f(T) + \log(q)(\nabla^{\log} f)(T) + \frac{1}{2} \log(q)^2 (\nabla^{\log})^2 f(T) + \dots$$

where $\log(q) = \sum_{n \geq 0} (-1)^{n-1} \frac{(q-1)^n}{n} \in \mathbb{Q}[[q-1]]$ so

$$\nabla^{q, \log} = \frac{\log(q)}{q-1} \nabla^{\log} + \frac{1}{2} \frac{\log(q)^2}{q-1} (\nabla^{\log})^2 + \dots$$

and taking $\tilde{\nabla}^{q, \log} = \log(q) \nabla^{q, \log}$, we get $\tilde{\nabla}^{q, \log} = \exp(\tilde{\nabla}^{\log}) - 1$.

Example 3.3 yields the following more general result.

Proposition 3.4. Let (R, \square) be a smooth framed \mathbb{Q} -algebra. There is an symmetric monoidal equivalence of categories

$$\left\{ \begin{array}{c} (q-1)\text{-adically complete } R[[q-1]]\text{-modules} \\ \text{with } q\text{-connection} \end{array} \right\} \simeq \left\{ \begin{array}{c} (q-1)\text{-adically complete } R[[q-1]]\text{-modules} \\ \text{with connection} \end{array} \right\}.$$

This proves coordinate independence, since the latter category is coordinate independent.

Corollary 3.5. Let (R, \square) be a framed \mathbb{Q} -algebra and denote the category of $(q-1)$ -adically complete $R[[q-1]]$ -modules with q -connection by $q\text{Mod}_{R[[q-1]]}$. The q -de Rham complex $q\Omega_{(R, \square)/\mathbb{Q}}$ computes $R\text{Hom}_{q\text{Mod}_{R[[q-1]]}}(\mathbb{1}, \mathbb{1})$ and is canonically independent of coordinates.

The case of modified q -connections is more subtle.

Definition 3.6 (Logarithmic q -Connections). Let R be a \mathbb{Q} -algebra. A h -connection over $R[h]$ is an $R[h]$ -module M with a map $\tilde{\nabla}_M : M \rightarrow M \otimes_R \Omega_{R/\mathbb{Q}}^1$ satisfying $(\tilde{\nabla}_M)^2 = 0$ and

$$\tilde{\nabla}_M(fm) = h \cdot \nabla(f) \cdot m + f \cdot \tilde{\nabla}_M(m).$$

Such constructions appear in Hodge and twistor theory under the name of Higgs fields.

We would like to see an analogue of Proposition 3.4.

Proposition 3.7. Let (R, \square) be a smooth framed \mathbb{Q} -algebra. There is an symmetric monoidal equivalence of categories

$$\left\{ \begin{array}{c} (q-1)\text{-adically complete } R[[q-1]]\text{-modules} \\ \text{with modified } q\text{-connection s.t. } \tilde{\nabla}_{i,M}^{q, \log} \text{'s are top. nil.} \end{array} \right\} \simeq \left\{ \begin{array}{c} h\text{-adically complete } R[[q-1]]\text{-modules} \\ \text{with } h\text{-connection s.t. } \tilde{\nabla}_M \text{ is top. nil.} \end{array} \right\}$$

$$\left(M, (\tilde{\nabla}_{i,M}^{q, \log})_{i=1}^d \right) \longleftarrow \left(M, (\tilde{\nabla}_{i,M}^{\log})_{i=1}^d \right)$$

where $\tilde{\nabla}_{i,M}^{q, \log} = \exp(\tilde{\nabla}_{i,M}^{\log}) - 1$.

Remark 3.8. The topological nilpotence of the endomorphisms ensure coverage of the exponential.

Once again, observing that the right hand side is coordinate independent, we get coordinate independence for modules with modified q -connections.

Corollary 3.9. Let (R, \square) be a framed \mathbb{Q} -algebra and denote the category of $(q-1)$ -adically complete $R[[q-1]]$ -modules with modified q -connection where the operators $\widetilde{\nabla}_{i,M}^{q,\log}$ are topologically nilpotent by $q\widetilde{\text{Mod}}_{R[[q-1]]}$. The q -Hodge complex $q\text{Hdg}_{(R,\square)/\mathbb{Q}}$ computes $R\text{Hom}_{q\widetilde{\text{Mod}}_{R[[q-1]]}}(\mathbb{1}, \mathbb{1})$ and is canonically independent of coordinates.

Deferring the discussion of coordinate independence integrally – which can be done by similarly isolating subcategories of modules with convergence conditions on their q -connections – we seek to understand the preceding constructions not just in the $(q-1)$ -adically complete case to the Habiro case, namely at all roots of unity.

In the preceding discussion, $(q-1)$ -adic completion allowed us to leverage étaleness of the map to produce a unique lift of the endomorphism on $\mathbb{Z}[q^\pm][T^\pm]$ since γ_i was infinitesimally close to the identity after $(q-1)$ -adic completion. But noticing that ζ_p is p -adically close to 1, we can attempt a similar approach.

Example 3.10. Let (R, \square) be a framed \mathbb{Z} -algebra with $\square : \text{Spec}(R) \rightarrow \mathbb{G}_m$. This gives a map

$$\mathbb{Z}[T^\pm]_p^\wedge[[q-1]] \longrightarrow R_p^\wedge[[q-1]]$$

which on specialization to $q = \zeta_p$ yields

$$\mathbb{Z}_p[\zeta_p]\langle T^\pm \rangle \longrightarrow R_p^\wedge[\zeta_p]$$

where using that ζ_p is close to 1 p -adically, $\gamma : \mathbb{Z}_p[\zeta_p]\langle T^\pm \rangle \rightarrow \mathbb{Z}_p[\zeta_p]\langle T^\pm \rangle$ by $T \mapsto qT$ lifts uniquely to an endomorphism $\gamma_R : R_p^\wedge[\zeta_p] \rightarrow R_p^\wedge[\zeta_p]$. However, $R[\zeta_p] \hookrightarrow R_p^\wedge[\zeta_p]$ may not have image stable under γ_R , for example, in the case of $\mathbb{G}_m \setminus \{1\}$.

So as seen in the example above, we will require an alternative description. For this, we produce an endomorphism of $R_p^\wedge[\zeta_p]$ that does globalize using the Frobenius map $\varphi : \mathbb{Z}[T^\pm] \rightarrow \mathbb{Z}[T^\pm]$ by $T \mapsto T^p$ lifts uniquely to R_p^\wedge and reduces to the Frobenius map on $R/(p)$. This produces an isomorphism

$$\mathbb{Z}[T^{\pm 1/p}] \otimes_{\mathbb{Z}[T^\pm]} R_p^\wedge \longrightarrow R_p^\wedge$$

by $T^{1/p} \mapsto T$ and thus

$$\mathbb{Z}[\zeta_p, T^{\pm 1/p}] \otimes_{\mathbb{Z}[T^\pm]} R_p^\wedge \longrightarrow R_p^\wedge[\zeta_p]$$

by the Frobenius once more. The map $\gamma : R_p^\wedge[\zeta_p] \rightarrow R_p^\wedge[\zeta_p]$ is induced by the map $\text{id}_{R_p^\wedge} \otimes (T^{1/p} \mapsto \zeta_p T^{1/p})$ but $\mathbb{Z}[\zeta_p, T^{\pm 1/p}] \otimes_{\mathbb{Z}[T^\pm]} R_p^\wedge$ contains $R \otimes_{\mathbb{Z}[T^\pm]} \mathbb{Z}[\zeta_p, T^{\pm 1/p}]$ as a subring, and since γ is the identity on R , γ is stable as an endomorphism.

Definition 3.11. Let $m \geq 1$. $\mathbb{Z}[\zeta_m, T^\pm]$ -algebra $R^{(m)} = R \otimes_{\mathbb{Z}[T^\pm]} \mathbb{Z}[\zeta_m, T^{\pm 1/m}]$ with algebra structure given by $T_i \mapsto T_i^{1/m}$ with action by $\gamma_i^{(m)} = \text{id}_R \otimes (T_i \mapsto \zeta_m T_i)$ lifting $T_i \mapsto \zeta_m T_i$ on $\mathbb{Z}[\zeta_m, T^\pm]$.

Example 3.12. Let $X = \mathbb{G}_m \setminus \{1\}$ and $R = \mathbb{Z}[T^\pm, \frac{1}{1-T}]$. Then $R^{(m)} = \mathbb{Z}[T^\pm, \frac{1}{1-T^m}]$ with the structure of a $\mathbb{Z}[\zeta_m, T^\pm]$ -algebra by $T \mapsto T^{1/m}$.

By uniqueness of deformation for étale algebras, we can deform from $q = \zeta_m$ to the completion at $\Phi_m(q)$, the m th cyclotomic polynomial, yielding a (formally) étale $\mathbb{Z}[q, \underline{T}^\pm]_{\Phi_m(q)}^\wedge$ -algebra R_m with lifts $\gamma_{i,m} : R_m \rightarrow R_m$. So for any m we can define the categories and complexes as before. While only defined at each m separately, we can use the fact that ζ_m and ζ_{pm} agree in characteristic p to glue the construction globally using the Frobenius, yielding a complex over the Habiro ring $\mathcal{H}_{(R, \square)}$.

REFERENCES

- [And01] Yves André. “Noncommutative differentials and Galois theory for differential or difference equations”. French. In: *Ann. Sci. Éc. Norm. Supér. (4)* 34.5 (2001), pp. 685–739. ISSN: 0012-9593. DOI: 10.1016/S0012-9593(01)01074-6. URL: <https://eudml.org/doc/82555>.
- [Aom90] Kazuhiko Aomoto. “ q -analogue of de Rham cohomology associated with Jackson integrals. I”. English. In: *Proc. Japan Acad., Ser. A* 66.7 (1990), pp. 161–164. ISSN: 0386-2194. DOI: 10.3792/pjaa.66.161.
- [Bha22] Bhargav Bhatt. *Prismatic F -gauges*. Lecture notes from a course at Princeton University, Fall 2022. 2022. URL: <https://www.math.ias.edu/~bhatt/teaching/mat549f22/lectures.pdf>.
- [BL22] Bhargav Bhatt and Jacob Lurie. *Absolute prismatic cohomology*. 2022. arXiv: 2201.06120 [math.AG]. URL: <https://arxiv.org/abs/2201.06120>.
- [BS22] Bhargav Bhatt and Peter Scholze. “Prisms and prismatic cohomology”. English. In: *Ann. Math. (2)* 196.3 (2022), pp. 1135–1275. ISSN: 0003-486X. DOI: 10.4007/annals.2022.196.3.5.
- [CS23] Dustin Clausen and Peter Scholze. *Lectures on Analytic Stacks*. Lecture notes from a course at the Universität Bonn, Winter 2023–24. 2023. URL: https://youtube.com/playlist?list=PLx5f8Ie1FRgGmu6gmL-Kf_Rl_6Mm7juZ0&si=aEs5GIFUAu4PW_i0.
- [Dri20] Vladimir Drinfeld. *Prismatization*. 2024. arXiv: 2005.04746 [math.AG]. URL: <https://arxiv.org/abs/2005.04746>.
- [GS+24] Stavros Garoufalidis et al. *The Habiro ring of a number field*. 2024. arXiv: 2412.04241 [math.NT]. URL: <https://arxiv.org/abs/2412.04241>.
- [Jac10] F. H. Jackson. “ q -difference equations.” English. In: *Am. J. Math.* 32 (1910), pp. 305–314. ISSN: 0002-9327. DOI: 10.2307/2370183.
- [MW24] Samuel Meyer and Ferdinand Wagner. *Derived q -Hodge complexes and refined TC^-* . 2024. arXiv: 2410.23115 [math.AT]. URL: <https://arxiv.org/abs/2410.23115>.
- [MT21] Matthew Morrow and Takeshi Tsuji. *Generalised representations as q -connections in integral p -adic Hodge theory*. 2021. arXiv: 2010.04059 [math.NT]. URL: <https://arxiv.org/abs/2010.04059>.
- [nLab-a] nLab authors. *Picard-Fuchs equation*. <https://ncatlab.org/nlab/show/Picard-Fuchs+equation>. Revision 2. Apr. 2025.
- [Sch17] Peter Scholze. “Canonical q -deformations in arithmetic geometry”. English. In: *Ann. Fac. Sci. Toulouse, Math. (6)* 26.5 (2017), pp. 1163–1192. ISSN: 0240-2963. DOI: 10.5802/afst.1563.
- [Sch24] Peter Scholze. *The Habiro Ring of a Number Field*. Lectures from a course at the Universität Bonn, Winter 2025–25. 2024. URL: <https://archive.mpim-bonn.mpg.de/id/eprint/5132/>.
- [Shi20] Ryotaro Shirai. *q -deformation with (φ, Γ) structure of the de Rham cohomology of the Legendre family of elliptic curves*. 2020. arXiv: 2006.12310 [math.AG]. URL: <https://arxiv.org/abs/2006.12310>.
- [Stacks] The Stacks project authors. *The Stacks project*. <https://stacks.math.columbia.edu>. 2023.
- [Wag24] Ferdinand Wagner. *q -Witt vectors and q -Hodge complexes*. 2024. arXiv: 2410.23078 [math.NT]. URL: <https://arxiv.org/abs/2410.23078>.

UNIVERSITÄT BONN, BONN, D-53113
Email address: `wgabrielong@uni-bonn.de`
URL: `https://wgabrielong.github.io/`