

## DEVELOPMENT OF THE MARA-LEB FACILITY

J. Romero<sup>1,2,\*</sup>, W. Gins<sup>1</sup>, I. D. Moore<sup>1</sup>, P. Papadakis<sup>3</sup>, J. Sarén<sup>1</sup>, J. Uusitalo<sup>1</sup> and A. Zadvornaya<sup>1</sup>.

<sup>1</sup>*Department of Physics, University of Jyväskylä, Finland*

<sup>2</sup>*Department of Physics, University of Liverpool, United Kingdom*

<sup>3</sup>*Nuclear Physics Group, STFC Daresbury Laboratory, United Kingdom*

\*Email: joromero@jyu.fi

MARA-LEB (Mass Analysing Recoil Apparatus - Low Energy Branch) is a facility currently in the final stages of design as an extension to the MARA separator at the Accelerator Laboratory of the University of Jyväskylä [1]. The aim of this new facility is to study exotic ions close to the proton drip line and the region near the  $N=Z=50$  doubly-magic shell closure. This region, which includes the heaviest self-conjugate ions, such as  $^{80}\text{Zr}$ ,  $^{94}\text{Ag}$  and  $^{100}\text{Sn}$ , is optimal for the testing of nuclear models and their predictions [2]. In addition to this, because many nuclei play a significant role in the astrophysical rapid proton (rp) capture process, this region of the nuclear chart is of importance in astrophysical nucleosynthesis models [3].

The combination with the existing MARA facility is ideal due to the high mass selectivity of the separator [4]. Mass-selected recoils from MARA enter the first part of the facility, the gas cell, through a thin window that separates a buffer gas region from MARA's high-vacuum environment. Incoming recoils will be stopped and neutralised by a buffer gas within the cell. The neutralisation of the recoils allows for subsequent in-gas-cell or in-gas-jet laser ionisation and spectroscopy via a state-of-the-art Ti:Sapphire laser system [5].

The ions emerging from the gas cell will be transported and focused towards the experimental stations by the MARA-LEB ion transport system [6]. This will be achieved by a 90°-bent radio-frequency quadrupole (RFQ), followed by a straight, segmented RFQ, which will transport the ions to acceleration electrodes. These will accelerate the ions to 30 keV and feed them into a magnetic dipole which will provide further mass separation for contaminant suppression.

A recent experiment has been carried out using the MARA separator to investigate the production of isotopes in MARA-LEB's region of interest close to  $^{96}\text{Ag}$ . The charge state distribution of  $^{96}\text{Pd}$  recoils at the focal plane of MARA will help to finalise design aspects concerning the size of the gas cell window. In this contribution, a general overview of the MARA-LEB facility along with preliminary results of the experiment will be given.

- [1] J. Sarén, et al., *Nucl. Inst. Methods Phys. Res. B* **266**, 4196 (2008).
- [2] T. Faestermann, M. Gorska, and H. Grawe, *Prog. Part. Nucl. Phys.* **69**, 85 (2013).
- [3] R. K. Wallace, and S. E. Woosley, *Astrophys. J. Suppl. Ser.*, **45**, 398 (1981).
- [4] J. Uusitalo, J. Sarén, J. Partanen, and J. Hilton, *Acta Phys. Pol. B* **50**, 319 (2019).
- [5] P. Papadakis, et al., *AIP Conf. Proc.* **2011**, 070013 (2018).
- [6] P. Papadakis, et al., *Nucl. Inst. Methods Phys. Res. B* **463**, 286 (2020).