

Automated Reasoning

Week 12. Confluence and Completion

Cynthia Kop

Fall 2024

Recall: some nice properties

- \mathcal{R} is **weakly normalising** (WN):
every term has a normal form
- \mathcal{R} is **terminating** (= strongly normalising, SN):
no infinite sequence of terms t_1, t_2, t_3, \dots exists such that $t_i \rightarrow_{\mathcal{R}} t_{i+1}$ for all i
- \mathcal{R} is **confluent** (= Church-Rosser, CR):
if $s \rightarrow_{\mathcal{R}}^ t$ and $s \rightarrow_{\mathcal{R}}^* q$ then a term u exists satisfying $t \rightarrow_{\mathcal{R}}^* u$ and $q \rightarrow_{\mathcal{R}}^* u$*
- \mathcal{R} is **locally confluent** (= weak Church-Rosser, WCR):
if $s \rightarrow_{\mathcal{R}} t$ and $s \rightarrow_{\mathcal{R}} q$ then a term u exists satisfying $t \rightarrow_{\mathcal{R}}^ u$ and $q \rightarrow_{\mathcal{R}}^* u$*

Recall: some nice properties

- \mathcal{R} is **weakly normalising** (WN):
every term has a normal form
- \mathcal{R} is **terminating** (= strongly normalising, SN):
no infinite sequence of terms t_1, t_2, t_3, \dots exists such that $t_i \rightarrow_{\mathcal{R}} t_{i+1}$ for all i
- \mathcal{R} is **confluent** (= Church-Rosser, CR):
if $s \rightarrow_{\mathcal{R}}^ t$ and $s \rightarrow_{\mathcal{R}}^* q$ then a term u exists satisfying $t \rightarrow_{\mathcal{R}}^* u$ and $q \rightarrow_{\mathcal{R}}^* u$*
- \mathcal{R} is **locally confluent** (= weak Church-Rosser, WCR):
if $s \rightarrow_{\mathcal{R}} t$ and $s \rightarrow_{\mathcal{R}} q$ then a term u exists satisfying $t \rightarrow_{\mathcal{R}}^ u$ and $q \rightarrow_{\mathcal{R}}^* u$*

Confluence implies uniqueness of normal forms

Confluence implies uniqueness of normal forms

Theorem

If a TRS is confluent, then every term has at most one normal form.

Confluence implies uniqueness of normal forms

Theorem

If a TRS is confluent, then every term has at most one normal form.

Proof: Assume t has two normal forms u, u' .

Then by confluence there is a v such that $u \rightarrow_{\mathcal{R}}^* v$ and $u' \rightarrow_{\mathcal{R}}^* v$.

Since u, u' are normal forms we have $u = v = u'$. \square

Termination + confluence implies existence of unique normal forms

We conclude:

Termination + confluence implies existence of unique normal forms

We conclude:

Theorem

If a TRS is terminating and confluent, then every term has exactly one normal form.

Termination + confluence implies existence of unique normal forms

We conclude:

Theorem

If a TRS is terminating and confluent, then every term has exactly one normal form.

This normal form can be found just by rewriting the term until no further rewriting step is possible.

Termination + confluence implies existence of unique normal forms

We conclude:

Theorem

If a TRS is terminating and confluent, then every term has exactly one normal form.

This normal form can be found just by rewriting the term until no further rewriting step is possible.

This is a very useful combination!

This week

This week

- confluence versus local confluence

This week

- confluence versus local confluence
- critical pairs

This week


- confluence versus local confluence
- critical pairs
- solving the **word problem**

Confluence (Church-Rosser property)

If $s \rightarrow^ t$ and $s \rightarrow^* q$
then exists u such that
 $t \rightarrow^* u$ and $q \rightarrow^* u$.*


Confluence (Church-Rosser property)

If $s \rightarrow^ t$ and $s \rightarrow^* q$
then exists u such that
 $t \rightarrow^* u$ and $q \rightarrow^* u$.*



Local confluence (Weak Church-Rosser property):

*If $s \rightarrow t$ and $s \rightarrow q$
then exists u such that
 $t \rightarrow^* u$ and $q \rightarrow^* u$.*



Local versus general confluence

Local versus general confluence

$$R = \{ \textcolor{red}{a} \rightarrow \textcolor{red}{b}, \textcolor{red}{b} \rightarrow \textcolor{red}{a}, \textcolor{red}{a} \rightarrow \textcolor{red}{c}, \textcolor{red}{b} \rightarrow \textcolor{red}{d} \}$$

Local versus general confluence

$$R = \{ a \rightarrow b, b \rightarrow a, a \rightarrow c, b \rightarrow d \}$$



Local versus general confluence

$$R = \{ \textcolor{red}{a} \rightarrow \textcolor{red}{b}, \textcolor{red}{b} \rightarrow \textcolor{red}{a}, \textcolor{red}{a} \rightarrow \textcolor{red}{c}, \textcolor{red}{b} \rightarrow \textcolor{red}{d} \}$$



This counterexample relies on the TRS being **non-terminating**.

Newman's lemma (1942)

Theorem

For terminating TRSs the properties confluence and local confluence are equivalent.

Principle of well-founded induction

Theorem

Let $\text{SN}(\rightarrow)$ and

$$\forall t [\underbrace{\forall u [t \rightarrow^+ u \Rightarrow P(u)]}_{\text{Induction Hypothesis}} \Rightarrow P(t)]$$

Then $P(t)$ holds for all t .

Principle of well-founded induction

Theorem

Let $\text{SN}(\rightarrow)$ and

$$\forall t [\underbrace{\forall u [t \rightarrow^+ u \Rightarrow P(u)]}_{\text{Induction Hypothesis}} \Rightarrow P(t)]$$

Then $P(t)$ holds for all t .

(Think of $t \rightarrow^+ u$ as $t > u$ as in well-known induction.)

Principle of well-founded induction

Theorem

Let $\text{SN}(\rightarrow)$ and

$$\forall t [\underbrace{\forall u [t \rightarrow^+ u \Rightarrow P(u)]}_{\text{Induction Hypothesis}} \Rightarrow P(t)]$$

Then $P(t)$ holds for all t .

(Think of $t \rightarrow^+ u$ as $t > u$ as in well-known induction.)

Proof: by contradiction! Assume $\neg P(t)$

Proof of Newman's Lemma

Proof of Newman's Lemma

Assume $SN(\rightarrow)$ and $WCR(\rightarrow)$. We have to prove $CR(\rightarrow)$.

Proof of Newman's Lemma

Assume $\text{SN}(\rightarrow)$ and $\text{WCR}(\rightarrow)$. We have to prove $\text{CR}(\rightarrow)$.

We apply the principle of well-founded induction for $P(s)$ being

$$\forall t, q [\text{if } s \rightarrow^* t \wedge s \rightarrow^* q \text{ then } \exists u [t \rightarrow^* u \wedge q \rightarrow^* u]]$$

Proof of Newman's Lemma


Assume $\text{SN}(\rightarrow)$ and $\text{WCR}(\rightarrow)$. We have to prove $\text{CR}(\rightarrow)$.

We apply the principle of well-founded induction for $P(s)$ being


$$\forall t, q [\text{if } s \rightarrow^* t \wedge s \rightarrow^* q \text{ then } \exists u [t \rightarrow^* u \wedge q \rightarrow^* u]]$$

Suppose $s \rightarrow^* t$ and $s \rightarrow^* q$.

Proof of Newman's Lemma




Proof of Newman's Lemma



We must find u such that $t \rightarrow^* u$ and $q \rightarrow^* u$.


Proof of Newman's Lemma



We must find u such that $t \rightarrow^* u$ and $q \rightarrow^* u$.

If $s = t$ we can choose $u := q$.

Proof of Newman's Lemma




We must find u such that $t \rightarrow^* u$ and $q \rightarrow^* u$.

If $s = t$ we can choose $u := q$.

If $s = q$ we can choose $u := t$.

Proof of Newman's Lemma




We must find u such that $t \rightarrow^* u$ and $q \rightarrow^* u$.

If $s = t$ we can choose $u := q$.


If $s = q$ we can choose $u := t$.

In the remaining case we have $s \rightarrow^+ t$ and $s \rightarrow^+ q$.

Proof of Newman's Lemma




Proof of Newman's Lemma




By $\text{WCR}(\rightarrow)$ we can find w such that $t_1 \rightarrow^* w$ and $q_1 \rightarrow^* w$.


Proof of Newman's Lemma



Proof of Newman's Lemma




Proof of Newman's Lemma




By the induction hypothesis on t_1 , there is a common reduct for t and w .


Proof of Newman's Lemma



Proof of Newman's Lemma




Proof of Newman's Lemma




By the induction hypothesis on q_1 , there is a common reduct for w and q .

Proof of Newman's Lemma




Proof of Newman's Lemma




Again by the induction hypothesis on q_1 , there is a common reduct for a and b .

Proof of Newman's Lemma



Proof of Newman's Lemma



Hence, t and q indeed have a common reduct u !

Deciding confluence

Bad news: both undecidable

Deciding confluence

Bad news: both undecidable

Good news: decidable for **terminating** TRSs!

Deciding confluence

Bad news: both undecidable

Good news: decidable for **terminating** TRSs!

Idea:

*analyze overlapping patterns in left hand sides of the rules, yielding **critical pairs**.*

Deciding confluence

Bad news: both undecidable

Good news: decidable for **terminating** TRSs!

Idea:

*analyze overlapping patterns in left hand sides of the rules, yielding **critical pairs**.*

In our example for addition of natural numbers,

$$\begin{aligned}\text{add}(0, y) &\rightarrow y \\ \text{add}(s(x), y) &\rightarrow s(\text{add}(x, y))\end{aligned}$$

there is no overlap. Hence it is locally confluent.

Deciding confluence

Bad news: both undecidable

Good news: decidable for **terminating** TRSs!

Idea:

*analyze overlapping patterns in left hand sides of the rules, yielding **critical pairs**.*

In our example for addition of natural numbers,

$$\begin{aligned}\text{add}(0, y) &\rightarrow y \\ \text{add}(s(x), y) &\rightarrow s(\text{add}(x, y))\end{aligned}$$

there is no overlap. Hence it is locally confluent.

Termination: for instance by LPO.

Critical pairs

Definition

Let $\ell_1 \rightarrow r_1$ and $\ell_2 \rightarrow r_2$ be two (possibly equal) rewrite rules, which have no variables in common.

Critical pairs

Definition

Let $\ell_1 \rightarrow r_1$ and $\ell_2 \rightarrow r_2$ be two (possibly equal) rewrite rules, which have no variables in common.

Suppose $\ell_2 = C[t]$, where C is a (possibly trivial) context, and t not a variable.

Critical pairs

Definition

Let $\ell_1 \rightarrow r_1$ and $\ell_2 \rightarrow r_2$ be two (possibly equal) rewrite rules, which have no variables in common.

Suppose $\ell_2 = C[t]$, where C is a (possibly trivial) context, and t not a variable.

Suppose t and ℓ_1 unify with mgu σ : $t\sigma = \ell_1\sigma$.

Critical pairs

Definition

Let $\ell_1 \rightarrow r_1$ and $\ell_2 \rightarrow r_2$ be two (possibly equal) rewrite rules, which have no variables in common.

Suppose $\ell_2 = C[t]$, where C is a (possibly trivial) context, and t not a variable.

Suppose t and ℓ_1 unify with mgu σ : $t\sigma = \ell_1\sigma$.

Then $\langle C[r_1]\sigma, r_2\sigma \rangle$ is a **critical pair** of $\ell_1 \rightarrow r_1$ and $\ell_2 \rightarrow r_2$.

Critical pairs

Definition

Let $\ell_1 \rightarrow r_1$ and $\ell_2 \rightarrow r_2$ be two (possibly equal) rewrite rules, which have no variables in common.

Suppose $\ell_2 = C[t]$, where C is a (possibly trivial) context, and t not a variable.

Suppose t and ℓ_1 unify with mgu σ : $t\sigma = \ell_1\sigma$.

Then $\langle C[r_1]\sigma, r_2\sigma \rangle$ is a **critical pair** of $\ell_1 \rightarrow r_1$ and $\ell_2 \rightarrow r_2$.

Note that $\ell_2\sigma = C[t]\sigma = C\sigma[\ell_1\sigma]$ can be rewritten in two ways:

- with $\ell_2 \rightarrow r_2$
- with $\ell_1 \rightarrow r_1$

Example

Example

Assume we have rules for arithmetic including:

$$x - x \rightarrow 0$$

$$s(x) - y \rightarrow s(x - y)$$

Example

Assume we have rules for arithmetic including:

$$\begin{aligned}x - x &\rightarrow 0 \\ s(x) - y &\rightarrow s(x - y)\end{aligned}$$

Then $s(x) - s(x)$ can be rewritten in two ways:

- to 0 by the first rule;
- to $s(x - s(x))$ by the second rule.

Example

Assume we have rules for arithmetic including:

$$\begin{aligned}x - x &\rightarrow 0 \\ s(x) - y &\rightarrow s(x - y)\end{aligned}$$

Then $s(x) - s(x)$ can be rewritten in two ways:

- to 0 by the first rule;
- to $s(x - s(x))$ by the second rule.

Now $\langle 0, s(x - s(x)) \rangle$ is a **critical pair**.

Example

Assume we have rules for arithmetic including:

$$\begin{aligned}x - x &\rightarrow 0 \\ s(x) - y &\rightarrow s(x - y)\end{aligned}$$

Then $s(x) - s(x)$ can be rewritten in two ways:

- to 0 by the first rule;
- to $s(x - s(x))$ by the second rule.

Now $\langle 0, s(x - s(x)) \rangle$ is a **critical pair**.

Notation:

- $\ell_1 \rightarrow r_1$ to be the rule $z - z \rightarrow 0$
- $\ell_2 \rightarrow r_2$ to be the rule $s(x) - y \rightarrow s(x - y)$
- C to be the trivial context \square
- $t = \ell_2 = s(x) - y$

Example

Assume we have rules for arithmetic including:

$$\begin{aligned}x - x &\rightarrow 0 \\ s(x) - y &\rightarrow s(x - y)\end{aligned}$$

Then $s(x) - s(x)$ can be rewritten in two ways:

- to 0 by the first rule;
- to $s(x - s(x))$ by the second rule.

Now $\langle 0, s(x - s(x)) \rangle$ is a **critical pair**.

Notation:

- $\ell_1 \rightarrow r_1$ to be the rule $z - z \rightarrow 0$
- $\ell_2 \rightarrow r_2$ to be the rule $s(x) - y \rightarrow s(x - y)$
- C to be the trivial context \square
- $t = \ell_2 = s(x) - y$

Indeed t, ℓ_1 unify, with mgu σ :

Example

Assume we have rules for arithmetic including:

$$\begin{aligned}x - x &\rightarrow 0 \\ s(x) - y &\rightarrow s(x - y)\end{aligned}$$

Then $s(x) - s(x)$ can be rewritten in two ways:

- to 0 by the first rule;
- to $s(x - s(x))$ by the second rule.

Now $\langle 0, s(x - s(x)) \rangle$ is a **critical pair**.

Notation:

- $\ell_1 \rightarrow r_1$ to be the rule $z - z \rightarrow 0$
- $\ell_2 \rightarrow r_2$ to be the rule $s(x) - y \rightarrow s(x - y)$
- C to be the trivial context \square
- $t = \ell_2 = s(x) - y$

Indeed t, ℓ_1 unify, with mgu $\sigma: \sigma(x) = x, \sigma(y) = \sigma(z) = s(x)$

Example

Let R consist of the single rule

$$f(f(x)) \rightarrow g(x)$$

Example

Let R consist of the single rule

$$f(f(x)) \rightarrow g(x)$$

By choosing

- $\ell_1 \rightarrow r_1$ to be the rule $f(f(x)) \rightarrow g(x)$
- $\ell_2 \rightarrow r_2$ to be the rule $f(f(y)) \rightarrow g(y)$
- $C = f(\square)$
- $t = f(y)$

Example

Let R consist of the single rule

$$f(f(x)) \rightarrow g(x)$$

By choosing

- $\ell_1 \rightarrow r_1$ to be the rule $f(f(x)) \rightarrow g(x)$
- $\ell_2 \rightarrow r_2$ to be the rule $f(f(y)) \rightarrow g(y)$
- $C = f(\square)$
- $t = f(y)$

we see that t, ℓ_1 unify, with mgu σ :

$$\sigma(x) = x, \quad \sigma(y) = f(x)$$

Example

Let R consist of the single rule

$$f(f(x)) \rightarrow g(x)$$

By choosing

- $\ell_1 \rightarrow r_1$ to be the rule $f(f(x)) \rightarrow g(x)$
- $\ell_2 \rightarrow r_2$ to be the rule $f(f(y)) \rightarrow g(y)$
- $C = f(\square)$
- $t = f(y)$

we see that t, ℓ_1 unify, with mgu σ :

$$\sigma(x) = x, \quad \sigma(y) = f(x)$$

This yields the critical pair $\langle f(g(x)), g(f(x)) \rangle$.

Convergence

A critical pair $\langle t, q \rangle$ **converges** if $t \rightarrow_R^* u$ and $q \rightarrow_R^* u$ for some u .

Convergence

A critical pair $\langle t, q \rangle$ **converges** if $t \rightarrow_R^* u$ and $q \rightarrow_R^* u$ for some u .

Example

$$x + s(y) \rightarrow s(x + y)$$

$$s(x) + y \rightarrow s(x + y)$$

Convergence

A critical pair $\langle t, q \rangle$ **converges** if $t \rightarrow_R^* u$ and $q \rightarrow_R^* u$ for some u .

Example

$$x + s(y) \rightarrow s(x + y) \qquad s(x) + y \rightarrow s(x + y)$$

Critical pair: $\langle s(x + s(y)), s(s(x) + y) \rangle$

Convergence

A critical pair $\langle t, q \rangle$ **converges** if $t \rightarrow_R^* u$ and $q \rightarrow_R^* u$ for some u .

Example

$$x + s(y) \rightarrow s(x + y) \qquad s(x) + y \rightarrow s(x + y)$$

Critical pair: $\langle s(x + s(y)), s(s(x) + y) \rangle$

We have: $s(x + s(y)) \rightarrow_{\mathcal{R}} s(s(x + y)) \leftarrow_{\mathcal{R}} s(s(x) + y)$.

Convergence

A critical pair $\langle t, q \rangle$ **converges** if $t \rightarrow_R^* u$ and $q \rightarrow_R^* u$ for some u .

Example

$$x + s(y) \rightarrow s(x + y) \qquad s(x) + y \rightarrow s(x + y)$$

Critical pair: $\langle s(x + s(y)), s(s(x) + y) \rangle$

We have: $s(x + s(y)) \rightarrow_{\mathcal{R}} s(s(x + y)) \leftarrow_{\mathcal{R}} s(s(x) + y)$.

Example

$$\text{not}(\top) \rightarrow \perp \qquad \text{not}(\perp) \rightarrow \top \qquad \text{not}(\text{not}(x)) \rightarrow x$$

Convergence

A critical pair $\langle t, q \rangle$ **converges** if $t \rightarrow_R^* u$ and $q \rightarrow_R^* u$ for some u .

Example

$$x + s(y) \rightarrow s(x + y) \qquad s(x) + y \rightarrow s(x + y)$$

Critical pair: $\langle s(x + s(y)), s(s(x) + y) \rangle$

We have: $s(x + s(y)) \rightarrow_{\mathcal{R}} s(s(x + y)) \leftarrow_{\mathcal{R}} s(s(x) + y)$.

Example

$$\text{not}(\top) \rightarrow \perp \qquad \text{not}(\perp) \rightarrow \top \qquad \text{not}(\text{not}(x)) \rightarrow x$$

Critical pair: $\langle \text{not}(\perp), \top \rangle$

Convergence

A critical pair $\langle t, q \rangle$ **converges** if $t \rightarrow_R^* u$ and $q \rightarrow_R^* u$ for some u .

Example

$$x + s(y) \rightarrow s(x + y) \qquad s(x) + y \rightarrow s(x + y)$$

Critical pair: $\langle s(x + s(y)), s(s(x) + y) \rangle$

We have: $s(x + s(y)) \rightarrow_{\mathcal{R}} s(s(x + y)) \leftarrow_{\mathcal{R}} s(s(x) + y)$.

Example

$$\text{not}(\top) \rightarrow \perp \qquad \text{not}(\perp) \rightarrow \top \qquad \text{not}(\text{not}(x)) \rightarrow x$$

Critical pair: $\langle \text{not}(\perp), \top \rangle$

We have: $\text{not}(\perp) \rightarrow \top \leftarrow_R^* \top$.

Critical pair lemma

Theorem

A TRS R is locally confluent if and only if all its critical pairs converge.

Critical pair lemma

Theorem

A TRS R is locally confluent if and only if all its critical pairs converge.

Example

Critical pair lemma

Theorem

A TRS R is locally confluent if and only if all its critical pairs converge.

Example

$$\mathbf{f}(\mathbf{f}(x)) \rightarrow \mathbf{g}(x)$$

Critical pair lemma

Theorem

A TRS R is locally confluent if and only if all its critical pairs converge.

Example

$$\mathbf{f}(\mathbf{f}(x)) \rightarrow \mathbf{g}(x)$$

Proof idea: if $s \rightarrow t$ and $s \rightarrow q$ there are three cases:

Critical pair lemma

Theorem

A TRS R is locally confluent if and only if all its critical pairs converge.

Example

$$\mathbf{f}(\mathbf{f}(x)) \rightarrow \mathbf{g}(x)$$

Proof idea: if $s \rightarrow t$ and $s \rightarrow q$ there are three cases:

- the reductions happen at distinct places in s

Critical pair lemma

Theorem

A TRS R is locally confluent if and only if all its critical pairs converge.

Example

$$\mathbf{f}(\mathbf{f}(x)) \rightarrow \mathbf{g}(x)$$

Proof idea: if $s \rightarrow t$ and $s \rightarrow q$ there are three cases:

- the reductions happen at distinct places in s
- one reduction occurs above the other, but without an overlap

Critical pair lemma

Theorem

A TRS R is locally confluent if and only if all its critical pairs converge.

Example

$$\mathbf{f}(\mathbf{f}(x)) \rightarrow \mathbf{g}(x)$$

Proof idea: if $s \rightarrow t$ and $s \rightarrow q$ there are three cases:

- the reductions happen at distinct places in s
- one reduction occurs above the other, but without an overlap
- the two reductions are an instance of a critical pair

Decision procedure

To decide $\text{WCR}(\mathcal{R})$ for **terminating** rules:

Decision procedure

To decide $\text{WCR}(\mathcal{R})$ for **terminating** rules:

- Compute all critical pairs $\langle t, q \rangle$

Decision procedure

To decide $\text{WCR}(\mathcal{R})$ for **terminating** rules:

- Compute all critical pairs $\langle t, q \rangle$
(by unifying left-hand sides with subterms of left-hand sides)

Decision procedure

To decide $\text{WCR}(\mathcal{R})$ for **terminating** rules:

- Compute all critical pairs $\langle t, q \rangle$
(by unifying left-hand sides with subterms of left-hand sides)
- For all critical pairs $\langle t, q \rangle$ compute

$$\{v \mid t \rightarrow^* v\} \cap \{u \mid q \rightarrow^* u\}$$

Decision procedure

To decide $\text{WCR}(\mathcal{R})$ for **terminating** rules:

- Compute all critical pairs $\langle t, q \rangle$
(by unifying left-hand sides with subterms of left-hand sides)

- For all critical pairs $\langle t, q \rangle$ compute

$$\{v \mid t \rightarrow^* v\} \cap \{u \mid q \rightarrow^* u\}$$

- If one of these sets is empty then $\text{WCR}(\mathcal{R})$ does not hold.

Decision procedure

To decide $\text{WCR}(\mathcal{R})$ for **terminating** rules:

- Compute all critical pairs $\langle t, q \rangle$
(by unifying left-hand sides with subterms of left-hand sides)

- For all critical pairs $\langle t, q \rangle$ compute

$$\{v \mid t \rightarrow^* v\} \cap \{u \mid q \rightarrow^* u\}$$

- If one of these sets is empty then $\text{WCR}(\mathcal{R})$ does not hold.
- If all of these sets are non-empty then $\text{WCR}(\mathcal{R})$ does hold.

Non-overlapping systems

A TRS is said to have **no overlap** if there are only **trivial** critical pairs.

Non-overlapping systems

A TRS is said to have **no overlap** if there are only **trivial** critical pairs.

Overlap-free TRSs are locally confluent.

Non-overlapping systems

A TRS is said to have **no overlap** if there are only **trivial** critical pairs.

Overlap-free TRSs are locally confluent.

Not every non-overlapping TRS is confluent:

$$\begin{array}{lcl} d(x, x) & \rightarrow & b \\ c(x) & \rightarrow & d(x, c(x)) \\ a & \rightarrow & c(a) \end{array}$$

Non-overlapping systems

A TRS is said to have **no overlap** if there are only **trivial** critical pairs.

Overlap-free TRSs are locally confluent.

Not every non-overlapping TRS is confluent:

$$\begin{array}{lcl} d(x, x) & \rightarrow & b \\ c(x) & \rightarrow & d(x, c(x)) \\ a & \rightarrow & c(a) \end{array}$$

$$c(a) \rightarrow d(a, c(a)) \rightarrow d(c(a), c(a)) \rightarrow b$$

$$c(a) \rightarrow c(c(a)) \rightarrow^+ c(b)$$

Orthogonality

A rule $\ell \rightarrow r$ is **left-linear** if no variable occurs more than once in ℓ .

Orthogonality

A rule $\ell \rightarrow r$ is **left-linear** if no variable occurs more than once in ℓ .

A TRS is **left-linear** if all its rules are.

Orthogonality

A rule $\ell \rightarrow r$ is **left-linear** if no variable occurs more than once in ℓ .

A TRS is **left-linear** if all its rules are.

A TRS is **orthogonal** if it is left-linear and has no overlap.

Orthogonality

A rule $\ell \rightarrow r$ is **left-linear** if no variable occurs more than once in ℓ .

A TRS is **left-linear** if all its rules are.

A TRS is **orthogonal** if it is left-linear and has no overlap.

Theorem

Any orthogonal TRS is confluent.

Orthogonality

A rule $\ell \rightarrow r$ is **left-linear** if no variable occurs more than once in ℓ .

A TRS is **left-linear** if all its rules are.

A TRS is **orthogonal** if it is left-linear and has no overlap.

Theorem

Any orthogonal TRS is confluent.

Proof idea: this uses that if $s \twoheadrightarrow t$ and $s \twoheadrightarrow q$, there exists u such that $t \twoheadrightarrow u$ and $q \twoheadrightarrow u$ (where \twoheadrightarrow indicates a **parallel** move).

Orthogonality

A rule $\ell \rightarrow r$ is **left-linear** if no variable occurs more than once in ℓ .

A TRS is **left-linear** if all its rules are.

A TRS is **orthogonal** if it is left-linear and has no overlap.

Theorem

Any orthogonal TRS is confluent.

Proof idea: this uses that if $s \twoheadrightarrow t$ and $s \twoheadrightarrow q$, there exists u such that $t \twoheadrightarrow u$ and $q \twoheadrightarrow u$ (where \twoheadrightarrow indicates a **parallel** move).

A TRS is **weakly orthogonal** if it is left-linear and all its critical pairs have a form $\langle t, t \rangle$.

Orthogonality

A rule $\ell \rightarrow r$ is **left-linear** if no variable occurs more than once in ℓ .

A TRS is **left-linear** if all its rules are.

A TRS is **orthogonal** if it is left-linear and has no overlap.

Theorem

Any **weakly** orthogonal TRS is confluent.

Proof idea: this uses that if $s \twoheadrightarrow t$ and $s \twoheadrightarrow q$, there exists u such that $t \twoheadrightarrow u$ and $q \twoheadrightarrow u$ (where \twoheadrightarrow indicates a **parallel** move).

A TRS is **weakly orthogonal** if it is left-linear and all its critical pairs have a form $\langle t, t \rangle$.

The Word Problem

Write $\leftrightarrow_{\mathcal{R}}^*$ for the reflexive symmetric transitive closure of $\rightarrow_{\mathcal{R}}$.

The Word Problem

Write $\leftrightarrow_{\mathcal{R}}^*$ for the reflexive symmetric transitive closure of $\rightarrow_{\mathcal{R}}$.

Question: given \mathcal{R}, s, t , does $s \leftrightarrow_{\mathcal{R}}^* t$ hold?

The Word Problem

Write $\leftrightarrow_{\mathcal{R}}^*$ for the reflexive symmetric transitive closure of $\rightarrow_{\mathcal{R}}$.

Question: given \mathcal{R}, s, t , does $s \leftrightarrow_{\mathcal{R}}^* t$ hold?

This is called the **word problem**.

The Word Problem

Write $\leftrightarrow_{\mathcal{R}}^*$ for the reflexive symmetric transitive closure of $\rightarrow_{\mathcal{R}}$.

Question: given \mathcal{R}, s, t , does $s \leftrightarrow_{\mathcal{R}}^* t$ hold?

This is called the **word problem**.

Decidability:

The Word Problem

Write $\leftrightarrow_{\mathcal{R}}^*$ for the reflexive symmetric transitive closure of $\rightarrow_{\mathcal{R}}$.

Question: given \mathcal{R}, s, t , does $s \leftrightarrow_{\mathcal{R}}^* t$ hold?

This is called the **word problem**.

Decidability: if \mathcal{R} is terminating and confluent!

The Word Problem

Write $\leftrightarrow_{\mathcal{R}}^*$ for the reflexive symmetric transitive closure of $\rightarrow_{\mathcal{R}}$.

Question: given \mathcal{R}, s, t , does $s \leftrightarrow_{\mathcal{R}}^* t$ hold?

This is called the **word problem**.

Decidability: if \mathcal{R} is terminating and confluent!

A terminating and confluent TRS is called **complete**.

The Word Problem

Write $\leftrightarrow_{\mathcal{R}}^*$ for the reflexive symmetric transitive closure of $\rightarrow_{\mathcal{R}}$.

Question: given \mathcal{R}, s, t , does $s \leftrightarrow_{\mathcal{R}}^* t$ hold?

This is called the **word problem**.

Decidability: if \mathcal{R} is terminating and confluent!

A terminating and confluent TRS is called **complete**.

Now we give a decision procedure for the word problem for complete TRSs.

The Word Problem

Write $\leftrightarrow_{\mathcal{R}}^*$ for the reflexive symmetric transitive closure of $\rightarrow_{\mathcal{R}}$.

Question: given \mathcal{R}, s, t , does $s \leftrightarrow_{\mathcal{R}}^* t$ hold?

This is called the **word problem**.

Decidability: if \mathcal{R} is terminating and confluent!

A terminating and confluent TRS is called **complete**.

Now we give a decision procedure for the word problem for complete TRSs.

Theorem

If \mathcal{R} is a complete TRS
and s', t' are normal forms of s, t ,
then $s \leftrightarrow_{\mathcal{R}}^* t$ if and only if $s' = t'$.

Proof (deciding the word lemma)

We need a lemma (easy by induction on $\leftrightarrow_{\mathcal{R}}^*$):

Proof (deciding the word lemma)

We need a lemma (easy by induction on $\leftrightarrow_{\mathcal{R}}^*$):

Lemma:

If \mathcal{R} is confluent and $s \leftrightarrow_{\mathcal{R}}^ t$*

then a term q exists such that $s \rightarrow_{\mathcal{R}}^ q$ and $t \rightarrow_{\mathcal{R}}^* q$.*

Proof (deciding the word lemma)

We need a lemma (easy by induction on $\leftrightarrow_{\mathcal{R}}^*$):

Lemma:

If \mathcal{R} is confluent and $s \leftrightarrow_{\mathcal{R}}^ t$*

then a term q exists such that $s \rightarrow_{\mathcal{R}}^ q$ and $t \rightarrow_{\mathcal{R}}^* q$.*

Proof of the theorem:

Proof (deciding the word lemma)

We need a lemma (easy by induction on $\leftrightarrow_{\mathcal{R}}^*$):

Lemma:

If \mathcal{R} is confluent and $s \leftrightarrow_{\mathcal{R}}^ t$*

then a term q exists such that $s \rightarrow_{\mathcal{R}}^ q$ and $t \rightarrow_{\mathcal{R}}^* q$.*

Proof of the theorem:

If $s' = t'$ then $s \rightarrow_{\mathcal{R}}^* s' = t' \leftarrow_{\mathcal{R}}^* t$, hence $s \leftrightarrow_{\mathcal{R}}^* t$.

Proof (deciding the word lemma)

We need a lemma (easy by induction on $\leftrightarrow_{\mathcal{R}}^*$):

Lemma:

If \mathcal{R} is confluent and $s \leftrightarrow_{\mathcal{R}}^ t$*

then a term q exists such that $s \rightarrow_{\mathcal{R}}^ q$ and $t \rightarrow_{\mathcal{R}}^* q$.*

Proof of the theorem:

If $s' = t'$ then $s \rightarrow_{\mathcal{R}}^* s' = t' \leftarrow_{\mathcal{R}}^* t$, hence $s \leftrightarrow_{\mathcal{R}}^* t$.

Conversely assume $s \leftrightarrow_{\mathcal{R}}^* t$.

Proof (deciding the word lemma)

We need a lemma (easy by induction on $\leftrightarrow_{\mathcal{R}}^*$):

Lemma:

If \mathcal{R} is confluent and $s \leftrightarrow_{\mathcal{R}}^ t$
then a term q exists such that $s \rightarrow_{\mathcal{R}}^* q$ and $t \rightarrow_{\mathcal{R}}^* q$.*

Proof of the theorem:

If $s' = t'$ then $s \rightarrow_{\mathcal{R}}^* s' = t' \leftarrow_{\mathcal{R}}^* t$, hence $s \leftrightarrow_{\mathcal{R}}^* t$.

Conversely assume $s \leftrightarrow_{\mathcal{R}}^* t$.

Then $s' \leftarrow_{\mathcal{R}}^* s \leftrightarrow_{\mathcal{R}}^* t \rightarrow_{\mathcal{R}}^* t'$, hence $s' \leftrightarrow_{\mathcal{R}}^* t'$.

Proof (deciding the word lemma)

We need a lemma (easy by induction on $\leftrightarrow_{\mathcal{R}}^*$):

Lemma:

If \mathcal{R} is confluent and $s \leftrightarrow_{\mathcal{R}}^ t$
then a term q exists such that $s \rightarrow_{\mathcal{R}}^* q$ and $t \rightarrow_{\mathcal{R}}^* q$.*

Proof of the theorem:

If $s' = t'$ then $s \rightarrow_{\mathcal{R}}^* s' = t' \leftarrow_{\mathcal{R}}^* t$, hence $s \leftrightarrow_{\mathcal{R}}^* t$.

Conversely assume $s \leftrightarrow_{\mathcal{R}}^* t$.

Then $s' \leftarrow_{\mathcal{R}}^* s \leftrightarrow_{\mathcal{R}}^* t \rightarrow_{\mathcal{R}}^* t'$, hence $s' \leftrightarrow_{\mathcal{R}}^* t'$.

According to the lemma a term q exists such that $s' \rightarrow_{\mathcal{R}}^* q$ and $t' \rightarrow_{\mathcal{R}}^* q$.

Proof (deciding the word lemma)

We need a lemma (easy by induction on $\leftrightarrow_{\mathcal{R}}^*$):

Lemma:

If \mathcal{R} is confluent and $s \leftrightarrow_{\mathcal{R}}^ t$
then a term q exists such that $s \rightarrow_{\mathcal{R}}^* q$ and $t \rightarrow_{\mathcal{R}}^* q$.*

Proof of the theorem:

If $s' = t'$ then $s \rightarrow_{\mathcal{R}}^* s' = t' \leftarrow_{\mathcal{R}}^* t$, hence $s \leftrightarrow_{\mathcal{R}}^* t$.

Conversely assume $s \leftrightarrow_{\mathcal{R}}^* t$.

Then $s' \leftarrow_{\mathcal{R}}^* s \leftrightarrow_{\mathcal{R}}^* t \rightarrow_{\mathcal{R}}^* t'$, hence $s' \leftrightarrow_{\mathcal{R}}^* t'$.

According to the lemma a term q exists such that $s' \rightarrow_{\mathcal{R}}^* q$ and $t' \rightarrow_{\mathcal{R}}^* q$.

Since s', t' are normal forms we have $s' = q = t'$. \square

Example

\mathcal{R} consists of the rule $\textcolor{red}{f}(f(f(x))) \rightarrow x$

Example

\mathcal{R} consists of the rule $f(f(f(x))) \rightarrow x$

Does $f^{17}(0) \leftrightarrow_{\mathcal{R}}^* f^{10}(0)$ hold?

Example

\mathcal{R} consists of the rule $\mathbf{f}(\mathbf{f}(\mathbf{f}(x))) \rightarrow x$

Does $\mathbf{f}^{17}(0) \leftrightarrow_{\mathcal{R}}^* \mathbf{f}^{10}(0)$ hold?

We can establish fully automatically that this is not the case:

Example

\mathcal{R} consists of the rule $f(f(f(x))) \rightarrow x$

Does $f^{17}(0) \leftrightarrow_{\mathcal{R}}^* f^{10}(0)$ hold?

We can establish fully automatically that this is not the case:

- check that \mathcal{R} is terminating

Example

\mathcal{R} consists of the rule $f(f(f(x))) \rightarrow x$

Does $f^{17}(0) \leftrightarrow_{\mathcal{R}}^* f^{10}(0)$ hold?

We can establish fully automatically that this is not the case:

- check that \mathcal{R} is terminating
- check that \mathcal{R} is locally confluent

Example

\mathcal{R} consists of the rule $f(f(f(x))) \rightarrow x$

Does $f^{17}(0) \leftrightarrow_{\mathcal{R}}^* f^{10}(0)$ hold?

We can establish fully automatically that this is not the case:

- check that \mathcal{R} is terminating
- check that \mathcal{R} is locally confluent
- compute the normal form $f(f(0))$ of $f^{17}(0)$

Example

\mathcal{R} consists of the rule $f(f(f(x))) \rightarrow x$

Does $f^{17}(0) \leftrightarrow_{\mathcal{R}}^* f^{10}(0)$ hold?

We can establish fully automatically that this is not the case:

- check that \mathcal{R} is terminating
- check that \mathcal{R} is locally confluent
- compute the normal form $f(f(0))$ of $f^{17}(0)$
- compute the normal form $f(0)$ of $f^{10}(0)$

Example

\mathcal{R} consists of the rule $f(f(f(x))) \rightarrow x$

Does $f^{17}(0) \leftrightarrow_{\mathcal{R}}^* f^{10}(0)$ hold?

We can establish fully automatically that this is not the case:

- check that \mathcal{R} is terminating
- check that \mathcal{R} is locally confluent
- compute the normal form $f(f(0))$ of $f^{17}(0)$
- compute the normal form $f(0)$ of $f^{10}(0)$
- these are different, hence the answer is **No**.

Completing a TRS

Often a TRS \mathcal{R} is not complete, but a complete TRS \mathcal{R}' satisfying

Completing a TRS

Often a TRS \mathcal{R} is not complete, but a complete TRS \mathcal{R}' satisfying

$$\leftrightarrow_{\mathcal{R}'}^* = \leftrightarrow_{\mathcal{R}}^*$$

Completing a TRS

Often a TRS \mathcal{R} is not complete, but a complete TRS \mathcal{R}' satisfying

$$\leftrightarrow_{\mathcal{R}'}^* = \leftrightarrow_{\mathcal{R}}^*$$

can be found in a systematic way.

Completing a TRS

Often a TRS \mathcal{R} is not complete, but a complete TRS \mathcal{R}' satisfying

$$\leftrightarrow_{\mathcal{R}'}^* = \leftrightarrow_{\mathcal{R}}^*$$

can be found in a systematic way.

Finding such a complete TRS is called **(Knuth-Bendix) completion**.

Completing a TRS

Often a TRS \mathcal{R} is not complete, but a complete TRS \mathcal{R}' satisfying

$$\leftrightarrow_{\mathcal{R}'}^* = \leftrightarrow_{\mathcal{R}}^*$$

can be found in a systematic way.

Finding such a complete TRS is called **(Knuth-Bendix) completion**.

The new complete TRS can be used for the word problem.

Completing a TRS

Often a TRS \mathcal{R} is not complete, but a complete TRS \mathcal{R}' satisfying

$$\leftrightarrow_{\mathcal{R}'}^* = \leftrightarrow_{\mathcal{R}}^*$$

can be found in a systematic way.

Finding such a complete TRS is called **(Knuth-Bendix) completion**.

The new complete TRS can be used for the word problem.

Often, instead of an original TRS we only have a set of (unoriented) equations.

Completing a TRS

Often a TRS \mathcal{R} is not complete, but a complete TRS \mathcal{R}' satisfying

$$\leftrightarrow_{\mathcal{R}'}^* = \leftrightarrow_{\mathcal{R}}^*$$

can be found in a systematic way.

Finding such a complete TRS is called **(Knuth-Bendix) completion**.

The new complete TRS can be used for the word problem.

Often, instead of an original TRS we only have a set of (unoriented) equations.

Class example: $f(f(x)) = g(x)$

Knuth-Bendix Completion: ingredients

Knuth-Bendix Completion: ingredients

Reduction order: fix a reduction order \succ on terms, i.e.,

Knuth-Bendix Completion: ingredients

Reduction order: fix a reduction order \succ on terms, i.e.,

- \succ is well-founded
- \succ is stable (if $s \succ t$ then $s\sigma \succ t\sigma$)
- \succ is monotonic (if $s \succ t$ then $f(\dots, s, \dots) \succ f(\dots, t, \dots)$)

Knuth-Bendix Completion: ingredients

Reduction order: fix a reduction order \succ on terms, i.e.,

- \succ is well-founded
- \succ is stable (if $s \succ t$ then $s\sigma \succ t\sigma$)
- \succ is monotonic (if $s \succ t$ then $\mathbf{f}(\dots, s, \dots) \succ \mathbf{f}(\dots, t, \dots)$)

Then:

If $\ell \succ r$ for every rule $\ell \rightarrow r$ in \mathcal{R} , then $SN(\mathcal{R})$.

Knuth-Bendix Completion: ingredients

Reduction order: fix a reduction order \succ on terms, i.e.,

- \succ is well-founded
- \succ is stable (if $s \succ t$ then $s\sigma \succ t\sigma$)
- \succ is monotonic (if $s \succ t$ then $f(\dots, s, \dots) \succ f(\dots, t, \dots)$)

Then:

If $\ell \succ r$ for every rule $\ell \rightarrow r$ in \mathcal{R} , then $SN(\mathcal{R})$.

Equations and rules:

- Initially, E contains the equations we want to complete.
- Initially, $\mathcal{R} = \emptyset$.

Knuth-Bendix Completion algorithm

Repeat until E is empty:

Knuth-Bendix Completion algorithm

Repeat until E is empty:

1. Remove an equation $s = t$ from E , and
 - add $s \rightarrow t$ to \mathcal{R} if $s \succ t$
 - add $t \rightarrow s$ to \mathcal{R} if $t \succ s$
 - give up otherwise

Knuth-Bendix Completion algorithm

Repeat until E is empty:

1. Remove an equation $s = t$ from E , and

- add $s \rightarrow t$ to \mathcal{R} if $s \succ t$
- add $t \rightarrow s$ to \mathcal{R} if $t \succ s$
- give up otherwise

2. After adding a new rule $\ell \rightarrow r$ to \mathcal{R} :

- compute all critical pairs between it and existing rules of \mathcal{R}
- compute all critical pairs between the new rule and itself

Knuth-Bendix Completion algorithm

Repeat until E is empty:

1. Remove an equation $s = t$ from E , and
 - add $s \rightarrow t$ to \mathcal{R} if $s \succ t$
 - add $t \rightarrow s$ to \mathcal{R} if $t \succ s$
 - give up otherwise
2. After adding a new rule $\ell \rightarrow r$ to \mathcal{R} :
 - compute all critical pairs between it and existing rules of \mathcal{R}
 - compute all critical pairs between the new rule and itself
3. For every such critical pair $\langle u, v \rangle$:
 - \mathcal{R} -rewrite u to a normal form u'
 - \mathcal{R} -rewrite v to a normal form v'
 - if $u' \neq v'$, then add $u' = v'$ as an equation to the set E

Three cases

What can happen in this Knuth-Bendix procedure?

Three cases

What can happen in this Knuth-Bendix procedure?

- it fails due to an equation $s = t$ in E for which neither $s \succ t$ nor $t \succ s$ holds;

Three cases

What can happen in this Knuth-Bendix procedure?

- it fails due to an equation $s = t$ in E for which neither $s \succ t$ nor $t \succ s$ holds;
- it fails since the procedure goes on forever: E gets larger and is never empty;

Three cases

What can happen in this Knuth-Bendix procedure?

- it fails due to an equation $s = t$ in E for which neither $s \succ t$ nor $t \succ s$ holds;
- it fails since the procedure goes on forever: E gets larger and is never empty;
- it ends with E being empty.

Completing the procedure

Suppose: the procedure ends with E empty.

Completing the procedure

Suppose: the procedure ends with E empty.

Then:

- \mathcal{R} is terminating since it only contains rules $\ell \rightarrow r$ satisfying $\ell \succ r$.

Completing the procedure

Suppose: the procedure ends with E empty.

Then:

- \mathcal{R} is terminating since it only contains rules $\ell \rightarrow r$ satisfying $\ell \succ r$.
- \mathcal{R} is locally confluent since all critical pairs converge, so \mathcal{R} is complete.

Completing the procedure

Suppose: the procedure ends with E empty.

Then:

- \mathcal{R} is terminating since it only contains rules $\ell \rightarrow r$ satisfying $\ell \succ r$.
- \mathcal{R} is locally confluent since all critical pairs converge, so \mathcal{R} is complete.
- Convertibility $\leftrightarrow_{\mathcal{R}}^*$ of the resulting \mathcal{R} is equivalent to convertibility of the original E since in the whole procedure $\leftrightarrow_{\mathcal{R} \cup E}^*$ remains invariant.

Observation

The relation $\leftrightarrow_{\mathcal{R}}^*$ is an equivalence relation.

Observation

The relation $\leftrightarrow_{\mathcal{R}}^*$ is an equivalence relation.

In a complete TRS, the normal form is a unique representation for the corresponding equivalence class.

Observation

The relation $\leftrightarrow_{\mathcal{R}}^*$ is an equivalence relation.

In a complete TRS, the normal form is a unique representation for the corresponding equivalence class.

This is used for instance in **superposition** (next week).

Quiz

1. Give an example why local confluence does not imply confluence.
2. Determine, using critical pairs, whether the following system is locally confluent:

$$\begin{array}{lcl} f(g(x), g(b)) & \rightarrow & f(x, x) \\ g(a) & \rightarrow & b \\ b & \rightarrow & a \end{array}$$

3. Use Knuth-Bendix completion to find a complete TRS with the same \leftrightarrow_R relation as the above TRS.