

GNG 1105 A
ENGINEERING MECHANICS

Mid-term Exam

October 24, 2013

Times: 80 minutes


Page 1 of 1

Prof. A. Skaff

Closed book Examination. Only non- programmable calculators are allowed.

1- (15 marks)

The steel plate shown is supported by a frictionless pin joint at A and a frictionless slide at B. A force $P = 50\text{ N}$ acts at C, and another force $Q = 10\text{ N}$ acts at G. Determine all reactions at points A and B. All dimensions are in mm.




2- (15 marks)

A 3m rod AB is supported by a ball and socket joint at A. The weight of this rod is 200N and acts at its mid-point C.

This rod is being held in equilibrium by 2 cables BD & BE which lie in the x - z plane as shown. BD is parallel to the x -axis and BE is parallel to the z -axis.

a) Draw the FBD of rod AB.

b) Determine the tension in cables BD and BE.



GNG 1105 A
ENGINEERING MECHANICS

Mid-term Exam

Oct 24, 2013


SOLUTIONS

1- FBD- Plate

$$\uparrow \sum M_A = 0$$

$$\begin{aligned} P \times \frac{3}{5} \times 60 \text{ mm} - P \times \frac{4}{5} \times 25 \text{ mm} + Q \times 40 \text{ mm} \\ + B \sin 20^\circ \times 25 \text{ mm} - B \cos 20^\circ \times 60 \text{ mm} &= 0 \\ 50 \times \frac{3}{5} \times 60 - 50 \times \frac{4}{5} \times 25 + 10 \times 40 \\ + B \sin 20^\circ \times 25 - B \cos 20^\circ \times 60 &= 0 \\ 1800 - 1000 + 400 + 8.55B - 56.38B &= 0 \\ 47.838 = 1200 \\ \therefore B = 251 \text{ N} \end{aligned}$$

ANS. $P = 50 \text{ N}$



$$\rightarrow \sum F_x = 0$$

$$\begin{aligned} P \times \frac{3}{5} - B \cos 20^\circ + A_x + Q &= 0 \\ 50 \times \frac{3}{5} - 251 \cos 20^\circ + A_x + 10 &= 0 \\ \therefore A_x = 23.4 - 30 - 10 &= -16.4 \text{ N}, \quad \therefore A_x = 16.4 \text{ N} \leftarrow \end{aligned}$$

$$\uparrow \sum F_y = 0$$

$$\begin{aligned} A_y + P \times \frac{4}{5} + B \sin 20^\circ &= 0 \\ A_y + 50 \times \frac{4}{5} + 251 \sin 20^\circ &= 0 \\ \therefore A_y = -10 - 38.6 &= -48.6 \text{ N}, \quad \therefore A_y = 48.6 \text{ N} \downarrow \end{aligned}$$

$$A = \sqrt{(16.4)^2 + (48.6)^2} = 51.29 \text{ N}$$

$$\alpha = \tan^{-1} \frac{48.6}{16.4} = 71.35^\circ$$

$$\therefore A = 51.29 \text{ N}$$



ANS


3.

a) FBD - Rod AB

See diagram.

b) $\sum M_A = 0$

$$\vec{T}_{BE} \times \vec{r}_{B/A} + \vec{T}_{BD} \times \vec{r}_{B/A} + W \times \vec{r}_{C/A} = 0 \quad (1)$$

Where, $\vec{T}_{BE} = T_{BE} \vec{k}$ $\vec{T}_{BD} = T_{BD} \vec{i}$ $\vec{r}_{B/A} = 2\vec{i} - 2\vec{j} + 1\vec{k}$ $\vec{r}_{C/A} = 1\vec{i} - 1\vec{j} + 0.5\vec{k}$ $W = -200\vec{j}$ 

Insert in (1):

$$T_{BE} \vec{k} (2\vec{i} - 2\vec{j} + 1\vec{k}) + T_{BD} \vec{i} (2\vec{i} - 2\vec{j} + 1\vec{k}) - 200\vec{j} (1\vec{i} - 1\vec{j} + 0.5\vec{k}) = 0$$

$$2T_{BE} \vec{j} + 2T_{BE} \vec{i} - 2T_{BD} \vec{k} - T_{BD} \vec{j} + 200\vec{k} - 100\vec{i} = 0$$

Equate the coefficients of \vec{i} , \vec{j} and \vec{k} to zero:

$$\begin{vmatrix} \vec{k} & \vec{j} & \vec{i} \end{vmatrix}$$

$$(\vec{i}) : 2T_{BE} - 100 = 0;$$

$$\therefore T_{BE} = \frac{100}{2} = \underline{\underline{50N}}$$

ANS.

$$(\vec{j}) : 2T_{BE} - T_{BD} = 0$$

$$2 \times 50 - T_{BD} = 0;$$

$$\therefore T_{BD} = \underline{\underline{100N}}$$

ANS.

$$(\vec{k}) : -2T_{BD} + 200 = 0;$$

$$\therefore T_{BD} = 100N \checkmark \text{ check.}$$

- End -

Another Method

$$3.) \begin{vmatrix} (+) & (-) & (+) \\ \bar{i} & \bar{j} & \bar{k} \\ 0 & 0 & T_{BE} \\ 2 & -2 & +1 \end{vmatrix} + \begin{vmatrix} (+) & (-) & (+) \\ \bar{i} & \bar{j} & \bar{k} \\ T_{BD} & 0 & 0 \\ 2 & -2 & 1 \end{vmatrix} + \begin{vmatrix} (+) & (-) & (+) \\ \bar{i} & \bar{j} & \bar{k} \\ 0 & -200 & 0 \\ 1 & -1 & 0.5 \end{vmatrix} = 0$$

$$(+2T_{BE}\bar{i} + 2T_{BE}\bar{j}) + (-T_{BD}\bar{j} - 2T_{BD}\bar{k}) + (-200 \times 0.5\bar{i} + 200\bar{k}) = 0$$

$$2T_{BE}\bar{i} + 2T_{BE}\bar{j} - T_{BD}\bar{j} - 2T_{BD}\bar{k} - 100\bar{i} + 200\bar{k} = 0$$

Equate \bar{i} , \bar{j} & \bar{k} to zero:

$$\textcircled{\bar{i}}: 2T_{BE} - 100 = 0; \quad \therefore T_{BE} = \frac{100}{2} = \underline{\underline{50 \text{ N}}} \quad \text{ANS.}$$

$$\textcircled{\bar{j}}: 2T_{BE} - T_{BD} = 0; \quad \therefore T_{BD} = 100 \text{ N} \quad \text{ANS.}$$

$$2 \times 50 - T_{BD} = 0;$$

$$\textcircled{\bar{k}}: -2T_{BD} + 200 = 0; \quad \therefore T_{BD} = \frac{200}{2} = 100 \text{ N} \quad \text{check.}$$