


# FUZZY STATISTICAL INFERENCES BASED ON FUZZY RANDOM VARIABLES



GHOLAMREZA HESAMIAN



CRC Press  
Taylor & Francis Group

A CHAPMAN & HALL BOOK

# Fuzzy Statistical Inferences Based on Fuzzy Random Variables



Taylor & Francis  
Taylor & Francis Group  
<http://taylorandfrancis.com>

# Fuzzy Statistical Inferences Based on Fuzzy Random Variables

Gholamreza Hesamian



CRC Press

Taylor & Francis Group

Boca Raton London New York

---

CRC Press is an imprint of the  
Taylor & Francis Group, an **informa** business  
A CHAPMAN & HALL BOOK

MATLAB® is a trademark of The MathWorks, Inc. and is used with permission. The MathWorks does not warrant the accuracy of the text or exercises in this book. This book's use or discussion of MATLAB® software or related products does not constitute endorsement or sponsorship by The MathWorks of a particular pedagogical approach or particular use of the MATLAB® software.

First edition published 2022  
by CRC Press  
6000 Broken Sound Parkway NW, Suite 300, Boca Raton, FL 33487-2742

and by CRC Press  
2 Park Square, Milton Park, Abingdon, Oxon, OX14 4RN

© 2022 Gholamreza Hesamian

CRC Press is an imprint of Taylor & Francis Group, LLC

Reasonable efforts have been made to publish reliable data and information, but the author and publisher cannot assume responsibility for the validity of all materials or the consequences of their use. The authors and publishers have attempted to trace the copyright holders of all material reproduced in this publication and apologize to copyright holders if permission to publish in this form has not been obtained. If any copyright material has not been acknowledged please write and let us know so we may rectify in any future reprint.

Except as permitted under U.S. Copyright Law, no part of this book may be reprinted, reproduced, transmitted, or utilized in any form by any electronic, mechanical, or other means, now known or hereafter invented, including photocopying, microfilming, and recording, or in any information storage or retrieval system, without written permission from the publishers.

For permission to photocopy or use material electronically from this work, access [www.copyright.com](http://www.copyright.com) or contact the Copyright Clearance Center, Inc. (CCC), 222 Rosewood Drive, Danvers, MA 01923, 978-750-8400. For works that are not available on CCC please contact [mpkbookspermissions@tandf.co.uk](mailto:mpkbookspermissions@tandf.co.uk)

*Trademark notice:* Product or corporate names may be trademarks or registered trademarks and are used only for identification and explanation without intent to infringe.

---

#### Library of Congress Cataloging-in-Publication Data

---

Names: Hesamian, Gholamreza, author.

Title: Fuzzy statistical inferences based on fuzzy random variables /  
Gholamreza Hesamian.

Description: First edition. | Boca Raton : CRC Press, 2022. | Includes  
bibliographical references and index.

Identifiers: LCCN 2021041840 (print) | LCCN 2021041841 (ebook) | ISBN  
9781032162225 (hardback) | ISBN 9781032164533 (paperback) | ISBN  
9781003248644 (ebook)

Subjects: LCSH: Fuzzy statistics. | Random variables. | Inference.

Classification: LCC QA276.5 .H42 2022 (print) | LCC QA276.5 (ebook) | DDC  
519.5--dc23/eng/20211112

LC record available at <https://lccn.loc.gov/2021041840>

LC ebook record available at <https://lccn.loc.gov/2021041841>

---

ISBN: 978-1-032-16222-5 (hbk)

ISBN: 978-1-032-16453-3 (pbk)

ISBN: 978-1-003-24864-4 (ebk)

DOI: [10.1201/9781003248644](https://doi.org/10.1201/9781003248644)

Typeset in CMR10  
by KnowledgeWorks Global Ltd.

*To my parents,  
Sedigheh and Mohammad*

*And special thanks to my wife, Samin*



Taylor & Francis  
Taylor & Francis Group  
<http://taylorandfrancis.com>

---

---

# **Contents**

---

<b>Foreword</b>	<b>xi</b>
<b>Preface</b>	<b>xiii</b>
<b>List of Figures</b>	<b>xv</b>
<b>List of Tables</b>	<b>xvii</b>
<b>Symbols</b>	<b>xxi</b>
<b>I Fuzzy Statistical Inferences Based on Fuzzy Random Variables</b>	<b>1</b>
<b>1 Introduction</b>	<b>3</b>
1.1 Fuzzy Set . . . . .	3
1.1.1 Operations of fuzzy sets . . . . .	6
1.1.2 Cartesian product of fuzzy sets . . . . .	12
1.2 Fuzzy Numbers . . . . .	13
1.2.1 A similarity measure . . . . .	16
1.2.2 A criteria for ranking FNs . . . . .	18
1.2.3 Distance measure . . . . .	22
1.3 $\alpha$ -values of FNs . . . . .	25
1.3.1 A generalized difference for FNs . . . . .	31
1.3.2 Maximum and minimum of the two FNs . . . . .	34
1.3.3 Absolute value of a FN . . . . .	36
1.4 Exercise . . . . .	37
1.5 Glossary . . . . .	39
<b>2 Probability of a Fuzzy Event</b>	<b>41</b>
2.1 Non-fuzzy Probability of a Fuzzy Event . . . . .	42
2.2 Probability of a Fuzzy Product Event . . . . .	49
2.3 Conditional Probability, Independence, and Bayes' Theorem	52
2.4 Fuzzy Probability of a FE and Its Properties . . . . .	55
2.5 Exercise . . . . .	67
2.6 Glossary . . . . .	71

<b>3 Descriptive Statistics Based on Fuzzy Data</b>	<b>73</b>
3.1 Central Tendency . . . . .	73
3.2 Fuzzy Deviation Criteria . . . . .	77
3.3 Box Plot Based on Fuzzy Data . . . . .	81
3.4 Descriptive Statistics Based on Grouped Fuzzy Data . . . . .	84
3.5 Measure of Spread Based on Fuzzy Data . . . . .	90
3.6 Exercise . . . . .	94
3.7 Glossary . . . . .	96
<b>4 Probability Reasoning Based on Fuzzy Random Variable (FRV)</b>	<b>97</b>
4.1 Fuzzy Random Variables . . . . .	97
4.1.1 Fuzzy expectation and exact variance . . . . .	99
4.1.2 Correlation between two <b>FRVs</b> . . . . .	102
4.1.3 Probability of an interval based on <b>FRVs</b> . . . . .	105
4.1.4 Liminf and limsup for a sequence of <b>FRVs</b> . . . . .	107
4.1.5 Fuzzy cumulative distribution function . . . . .	108
4.1.6 Location-Scale <b>FRVs</b> . . . . .	113
4.2 Probabilistic Inequalities for Fuzzy Random Variables . . . . .	123
4.3 Limit Theorems Based on <b>FRVs</b> . . . . .	126
4.4 Exercise . . . . .	134
4.5 Glossary . . . . .	136
<b>5 Fuzzy Hypothesis Testing for FRVs</b>	<b>137</b>
5.1 Testing Fuzzy Hypotheses Based on Normal <b>FRVs</b> . . . . .	138
5.1.1 One-sample testing fuzzy hypotheses based on normal <b>FRVs</b> . . . . .	140
5.1.2 Two-samples testing fuzzy hypotheses based on normal <b>FRVs</b> . . . . .	148
5.2 Statistical Non-parametric Tests Based on <b>FRVs</b> . . . . .	155
5.2.1 Sign test . . . . .	155
5.2.2 Wilcoxon test . . . . .	159
5.2.3 Kolmogorov-Smirnov test based on <b>FRVs</b> . . . . .	162
5.2.3.1 One-sample Kolmogorov-Smirnov test . . . . .	162
5.2.3.2 Two-sample Kolmogorov-Smirnov test for <b>TFRVs</b> . . . . .	166
5.2.4 Kruskal-Wallis test . . . . .	170
5.3 One-Way ANOVA Based on <b>FRVs</b> . . . . .	178
5.4 Exercise . . . . .	183
5.5 Glossary . . . . .	190
<b>6 Fuzzy Quality Control and Reliability Systems for FRVs</b>	<b>193</b>
6.1 Fuzzy Control Charts and Capability Indices Based on <b>NFRVs</b> . . . . .	193
6.1.1 Fuzzy $\bar{x}$ -chart and $s$ -chart for <b>NFRVs</b> . . . . .	194

6.1.2	Fuzzy EWMA chart based on <b>NFRVs</b> . . . . .	197
6.1.3	Process capability indices based on <b>NFRVs</b> . . . . .	202
6.1.4	Testing hypotheses for fuzzy process capability $\tilde{C}_p$ . .	209
6.2	Reliability Evaluation Using <b>FRVs</b> . . . . .	210
6.2.1	Components of a reliability system based on <b>LTFRVs</b> . . . . .	212
6.3	Exercise . . . . .	220
6.4	Glossary . . . . .	226
<b>7</b>	<b>Regression and Time Series Models for FRVs</b>	<b>227</b>
7.1	Regression Analysis Based on <b>FRVs</b> . . . . .	227
7.1.1	A multivariate regression analysis based on <b>FRVs</b> . .	228
7.1.2	Fuzzy non-parametric regression model . . . . .	235
7.2	Time Series Models for <b>FRVs</b> . . . . .	241
7.2.1	Autocorrelation criterion based on a fuzzy time series . . . . .	242
7.2.2	Multivariate time series model based on fuzzy time series data . . . . .	246
7.2.3	Non-parametric time series model based on fuzzy time series data . . . . .	253
7.3	Exercise . . . . .	257
7.4	Glossary . . . . .	261
<b>Bibliography</b>		<b>263</b>
<b>Index</b>		<b>279</b>



Taylor & Francis  
Taylor & Francis Group  
<http://taylorandfrancis.com>

---

## **Foreword**

---

Probability and statistics are concerned with the chance to analyze events and analysis of data and stochastic models. Therefore, the quantitative description of events and data is essential for statistical inferences.

This book tries to extend some common statistical inferences adopted with new ideas and simple techniques as much as possible for the probability of fuzzy events and statistics with fuzzy data. For simplicity in theoretic and calculations, all fuzzy statistical procedures were conducted on triangular fuzzy numbers. These methods essentially rely on some fuzzy statistical methods recently published by the author. Many of these methods also modified and developed as much as possible.

Compared to similar fuzzy statistics books, we tried to cover a set of frequently used statistical techniques used in applied sciences. Unlike other existing fuzzy statistics books, this book tried to gather almost all essential statistical methods such as exact and fuzzy probabilities, probabilistic inequalities, limit theorems, hypothesis tests, quality control, reliability theory, analysis of variance, and descriptive statistics for fuzzy data. In each section, the proposed techniques were also compared with other existing ones. Each section was then ended via some theoretical and applied exercises. The required optimization procedure can be done by any mathematical software. However, in this book the Mathematica and MATLAB software were utilized in calculations and plotting the figures. The available programs can be obtained from the author.

It is my hope that this book will serve as a solid starting point for analyzing fuzzy quantities and thus enlarging the domain of applicability of the field of probability and statistics.



Taylor & Francis  
Taylor & Francis Group  
<http://taylorandfrancis.com>

---

## Preface

---

Probability and statistics are two fascinating topics on the relationship between mathematics and real-life applications. The probabilistic methods provide a rigorous framework for modeling uncertainty due to randomness. Statistics can be classified into two different categories namely descriptive statistics and statistical inference. Descriptive statistics describe the data, whereas statistical inference consists in the use of statistics to draw conclusions about some unknown aspect of a population based on a random sample from that population. It also helps to assess the relationship between the dependent and independent variables. In statistical inference, the data are taken from the sample and allow you to generalize the population.

However, the methods of probability theory and statistics typically involve processing imprecision of two distinct types arising from a lack of knowledge due to inherent vagueness in concepts themselves which, in the sense of classical probability and statistical inferences, may be well defined. Therefore, the conventional methods are usually inadequate for dealing with certain kinds of imprecision called fuzzy sets. There are many situations in the conventional probability theory that an event and/or a parameter are fuzzy quantities. Moreover, there are many practical problems that require dealing with observations that represent inherently imprecise or linguistic characteristics. In such cases, fuzzy sets may be more effective in encoding such quantities rather than precise ones.

It is now generally accepted that statistical theory and the theory of fuzzy sets are both used to study uncertainties where both types of uncertainties including randomness and fuzziness have occurred. Thus, to produce suitable statistical inferences dealing with imprecise information, we need to model the imprecise information and extend the usual probability theory and statistics to imprecise environments.

This book is an honest attempt to produce an unified techniques for the elementary probability theory based on fuzzy events induced by a family of density functions induced by fuzzy parameters. The main attempt of this book is to introduce some new techniques for the most commonly used statistical inferences based on fuzzy data. This book is designed as a practical guide for scientists to help them solve statistical problems involving fuzzy data and fuzzy events. Consequently, the book has a practical bent and contains lots of examples. A familiarity with probability and statistics is assumed for this book. The book is intended for a statistical scientist at the Bachelor of Science

level or above for a specific course on probability and statistics. It also has applications to other applied sciences such as engineering and social sciences.

Payame Noor University, Department of Statistics, Tehran 19395-3697, Iran.

*Gholamreza Hesamian*

September 2021

---

---

## *List of Figures*

---

1.1	Plot of $g(\tilde{A})$ in Example 1.9. . . . .	17
1.2	Membership functions of $\tilde{A}$ and $\tilde{B}$ in three cases in Example 1.12. . . . .	20
1.3	Membership functions of $\tilde{A}_i$ , $i = 1, 2, \dots, 5$ in Example 1.13. . . . .	21
1.4	Membership function of $\exp(\tilde{A})$ in Example 1.18. . . . .	28
1.5	Membership functions of $\widetilde{\min}(\tilde{A}, \tilde{B})$ and $\widetilde{\max}(\tilde{A}, \tilde{B})$ in Example 1.21. . . . .	35
2.1	Membership function of $\tilde{P}_X^\theta(\tilde{A})$ in Example 2.14. . . . .	59
2.2	Membership function of $\tilde{P}_X(\tilde{A})$ in Example 2.16. . . . .	65
2.3	Membership function of $\tilde{P}_X(\tilde{A})$ in Example 2.17. . . . .	65
2.4	Membership functions of $(\text{NVL}, \text{NL}, \dots, \text{PVL})$ in Exercise 2.8. . . . .	67
3.1	A typical Boxplot for fuzzy data. . . . .	83
3.2	A typical fuzzy frequency distribution. . . . .	87
4.1	Fuzzy quantities of the ‘perception of the registration fee’ . . . . .	98
4.2	Plot of $\tilde{P}(\tilde{X} \in (35000, \infty))$ in Example 4.6. . . . .	107
4.3	Plot of $\tilde{F}_{\tilde{X}}(x)$ at $x = 188, 190, 192, 194$ in Example 4.8. . . . .	111
4.4	Plot of $\tilde{F}_{\tilde{X}}(x)$ on $[185, 200]$ in Example 4.8. . . . .	111
4.5	Plot of $\tilde{F}_{\tilde{X}}(x)$ at $x = -1, +1$ in Example 4.9. . . . .	112
4.6	Plot of $\tilde{F}_{\tilde{X}}(z)$ at three points of 400, 500, and 600 in Example 4.10. . . . .	118
4.7	Plot of $\tilde{P}(\tilde{X} \in (200, 700))$ in Example 4.10. . . . .	119
4.8	Plot of $\tilde{F}_{\tilde{X}}(z)$ at three points of $-1, 0$ , and 2 in Example 4.11. . . . .	120
4.9	Plot of $\tilde{P}(\tilde{X} \in (7, 10))$ in Example 4.11. . . . .	120
6.1	Fuzzy control limits for each fuzzy sample means in Example 6.1. . . . .	199
6.2	The system reliability in Example 6.6. . . . .	217
6.3	Diagram of five unit bridge network in Exercise 6.6. . . . .	224
7.1	ACF plot in Example 7.6. . . . .	247



Taylor & Francis  
Taylor & Francis Group  
<http://taylorandfrancis.com>

---

---

# List of Tables

---

1.1	Preference degrees $P_d(\tilde{A}_i \succeq \tilde{A}_j)$ in Example 1.13. . . . .	21
2.1	Some $\alpha$ -cuts of $\tilde{P}_X^\theta(\tilde{A})$ in Example 2.14. . . . .	58
2.2	Some $\alpha$ -cuts of $\tilde{P}_X^p(\tilde{A})$ in Example 2.16. . . . .	64
2.3	Some $\alpha$ -cuts of $\tilde{P}_X^\theta(\tilde{A})$ in Example 2.17. . . . .	66
3.1	Calculation of the elements of fuzzy geometric mean in Example 3.1. . . . .	75
3.2	Calculation of the elements of fuzzy harmonic mean in Example 3.2. . . . .	75
3.3	Preference degrees in Example 3.3. . . . .	77
3.4	Preference degrees in Example 3.4. . . . .	78
3.5	Data set in Example 3.6. . . . .	78
3.6	Preference degrees in Example 3.6. . . . .	79
3.7	Computing the elements of $S_D$ in Example 3.7 . . . . .	80
3.8	Computing the elements of $s^2$ in Example 3.8. . . . .	81
3.9	Data set in Example 3.9. . . . .	82
3.10	Preference degrees in Example 3.9. . . . .	85
3.11	Outlier detecting in Example 3.9. . . . .	86
3.12	A typical frequency table based on fuzzy data. . . . .	87
3.13	Data set in Example 3.10. . . . .	89
3.14	Preference degrees for the first seven fuzzy data in Example 3.10. . . . .	90
3.15	Preference degrees for the last seven fuzzy data in Example 3.10. . . . .	90
3.16	The values of $D_{ij} = \{P_d(\tilde{x}_j \succeq \tilde{a}_i), P_d(\tilde{x}_j \preceq \tilde{a}_{i+1})\}$ in Example 3.10. . . . .	91
3.17	Frequency table in Example 3.10. . . . .	92
3.18	Frequency table in Exercise 3.2. . . . .	95
3.19	The center values of fuzzy life of laptop batteries in Exercise 3.4 . . . . .	95
4.1	Some $\alpha$ -cuts of $\tilde{P}(\tilde{X} \in (35000, \infty))$ in Example 4.6. . . . .	106
4.2	Some $\alpha$ -cuts of $\tilde{P}(\tilde{X} \in (35000, \infty))$ in Example 4.8. . . . .	110
4.3	Some $\alpha$ -cuts of $\tilde{F}_{\tilde{X}}(x)$ at $x = -1, 1$ in Example 4.9. . . . .	112
4.4	Data set in Example 4.14. . . . .	122

5.1	Data set in Example 5.1. . . . .	144
5.2	The fuzzy failure time observations (in projection hours) for the 31 lamps in Example 5.2. . . . .	144
5.3	Computing the elements of $s_n^2$ in Example 5.2. . . . .	146
5.4	Data set in Example 5.3. . . . .	147
5.5	Data set in Example 5.4. . . . .	153
5.6	Data set in Example 5.5 . . . . .	153
5.7	Data set in Example 5.6. . . . .	154
5.8	Computing the elements of $s^2$ in Example 5.6. . . . .	156
5.9	The critical values for testing $H_0$ versus $H_1$ for the Sign test.	157
5.10	Preference degrees of observed <b>FRS</b> over $\tilde{M}_0$ in Example 5.8.	159
5.11	Preference degrees in Example 5.9. . . . .	161
5.12	Calculation the elements of <b>FECDF</b> in Example 5.10. . . . .	167
5.13	Data set in Example 5.11. . . . .	169
5.14	Calculation the elements of <b>FECDFs</b> in Example 5.11. . . . .	171
5.15	The judges ( $\tilde{x}_{ij}$ ) for each presentation in each group in Example 5.12. . . . .	174
5.16	The values of $P_d(\tilde{x}_{1l} \succ \tilde{w}_j)$ in Example 5.12. . . . .	174
5.17	The values of $P_d(\tilde{x}_{2l} \succ \tilde{w}_j)$ in Example 5.9. . . . .	175
5.18	The values of $P_d(\tilde{x}_{3l} \succ \tilde{w}_j)$ in Example 5.9. . . . .	175
5.19	Fuzzy values of tensile strength in Example 5.13. . . . .	182
5.20	Computing the elements of <b>SST</b> in Example 5.13. . . . .	183
5.21	Computing the elements of <b>SSR</b> in Example 5.13. . . . .	184
5.22	Values of $d_2^2(\bar{x}_{i.}, \bar{x}_{j.})$ in Example 5.13. . . . .	184
5.23	Multiple comparison of fuzzy means in Example 5.13. . . . .	184
5.24	Data set in Exercise 5.1. . . . .	185
5.25	Data set in Exercise 5.2. . . . .	185
5.26	Data set in Exercise 5.3. . . . .	186
5.27	Data set in Exercise 5.4 . . . . .	186
5.28	Data set in Exercise 5.5. . . . .	187
5.29	Data set in Exercise 5.6. . . . .	188
5.30	Data set in Exercise 5.7. . . . .	188
5.31	Data set in Exercise 5.8. . . . .	189
5.32	Data set in Exercise 5.9. . . . .	189
5.33	Fuzzy data in Exercise 5.10. . . . .	191
6.1	A linguistic interpretation of degree of violence $D_v$ . . . . .	196
6.2	Data set in Example 6.1. . . . .	197
6.3	$\tilde{\bar{x}}_i$ , $s_i^2$ and $(D_v)_i$ in Example 6.1. . . . .	198
6.4	Data set in Example 6.2. . . . .	203
6.5	Degrees of violence in Example 6.2. . . . .	204
6.6	Quality conditions for $\mathcal{C}_p$ . . . . .	205
6.7	Fuzzy data in Example 6.3. . . . .	208
6.8	Degrees to which $\hat{\mathcal{C}}_p$ belongs to $I_i$ , $i = 1, 2, \dots, 6$ in Example 6.3. . . . .	209

6.9 Some specified values of $\tilde{R}_S(t)$ and $\tilde{M}_S(t)$ in Example 6.5 . . . . .	217
6.10 Some specified values of $\tilde{R}_S(t)$ , $\tilde{M}_S(t)$ , and $\tilde{C}_S(2t, t)$ in Example 6.6 . . . . .	220
6.11 Data set in Exercise 6.1 . . . . .	221
6.12 Data set in Exercise 6.2 . . . . .	222
6.13 Data set in Exercise 6.3 . . . . .	223
7.1 Data set in Example 7.1. . . . .	232
7.2 Values of $d_2^2(\tilde{y}_i, \hat{\tilde{y}}_i)$ and $S(\tilde{y}_i, \hat{\tilde{y}}_i)$ in Example 7.1. . . . .	233
7.3 Data set in Example 7.2. . . . .	234
7.4 Values of $d_2^2(\tilde{y}_i, \hat{\tilde{y}}_i)$ and $S(\tilde{y}_i, \hat{\tilde{y}}_i)$ in Example 7.2. . . . .	235
7.5 Data set in Example 7.3. . . . .	237
7.6 Values of $d_2^2(\tilde{y}_i, \hat{\tilde{y}}_i)$ and $S(\tilde{y}_i, \hat{\tilde{y}}_i)$ in Example 7.3. . . . .	239
7.7 Data set in Example 7.4. . . . .	240
7.8 Values of $d_2^2(\tilde{y}_i, \hat{\tilde{y}}_i)$ and $S(\tilde{y}_i, \hat{\tilde{y}}_i)$ in Example 7.4. . . . .	241
7.9 Data set in Example 7.6. . . . .	245
7.10 Calculation the elements of $\hat{c}_1$ in Example 7.6. . . . .	246
7.11 Values of $\hat{\rho}_k$ for some $k$ in Example 7.6. . . . .	246
7.12 The center values of fuzzy data in Example 7.7. . . . .	249
7.13 Preference measures and estimated values of coefficients in Example 7.7. . . . .	250
7.14 Values of $d_2^2(\tilde{x}_t, \hat{\tilde{x}}_t^5)$ and $S(\tilde{x}_t, \hat{\tilde{x}}_t^5)$ in Example 7.7. . . . .	251
7.15 Fuzzy time series data in Example 7.8. . . . .	251
7.16 Performance measures and estimated values of coefficients in Example 7.7. . . . .	252
7.17 The values of $d_2^2(\tilde{x}_t, \hat{\tilde{x}}_t^3)$ and $S(\tilde{x}_t, \hat{\tilde{x}}_t^3)$ for $p = 3$ in Example 7.7.	252
7.18 Performance measures and estimated values of bandwidths in Example 7.9. . . . .	255
7.19 Values of $d_2^2(\tilde{x}_t, \hat{\tilde{x}}_t^2)$ and $S(\tilde{x}_t, \hat{\tilde{x}}_t^2)$ in Example 7.9. . . . .	256
7.20 Performance measures and estimated values of bandwidths in Example 7.10. . . . .	256
7.21 Values of $d_2^2(\tilde{x}_t, \hat{\tilde{x}}_t^1)$ and $S(\tilde{x}_t, \hat{\tilde{x}}_t^1)$ in Example 7.10. . . . .	257
7.22 Performance measures and estimated values of coefficients for some $p$ based on the fuzzy linear time series model in Example 7.10. . . . .	257
7.23 Data set in Exercise 7.1. . . . .	258
7.24 Data set in Exercise 7.2. . . . .	259
7.25 Data set in Exercise 7.3. . . . .	260
7.26 Kernel functions in Exercise 7.4. . . . .	260
7.27 Data set in Exercise 7.9. . . . .	261



Taylor & Francis  
Taylor & Francis Group  
<http://taylorandfrancis.com>

# Symbols

---

## Symbol Description

$\tilde{A}$	Crisp set	$ \tilde{A} $	Cardinal number of a fuzzy set $\tilde{A}$
$\tilde{A}$	Fuzzy set		
<b>FN</b>	Fuzzy number	$\tilde{A} \cup \tilde{B}$	Union of fuzzy sets $\tilde{A}$ and $\tilde{B}$
$\tilde{A}$	Fuzzy set		
<b>TFN</b>	Triangular fuzzy number	$\tilde{A} \cap \tilde{B}$	Intersection of fuzzy sets $\tilde{A}$ and $\tilde{B}$
<b>STFN</b>	Symmetric triangular fuzzy number	$\tilde{A} - \tilde{B}$	Difference between two fuzzy sets $\tilde{A}$ and $\tilde{B}$
$\tilde{A}[\alpha]$	$\alpha$ -cut of a fuzzy number $\tilde{A}$	$\tilde{A} \odot \tilde{B}$	Product of two fuzzy sets $\tilde{A}$ and $\tilde{B}$
$\tilde{A}_{\alpha}^L$	Lower bound of the $\alpha$ -cut of a fuzzy number $\tilde{A}$	$\tilde{A} \Delta \tilde{B}$	Symmetric difference of $\tilde{A}$ and $\tilde{B}$
$\tilde{A}_{\alpha}^U$	Upper bound of the $\alpha$ -cut of a fuzzy number $\tilde{A}$	$\tilde{A}^c$	Complement of $\tilde{A}$
$\{x_1, x_2, \dots\}$	Set of elements $x_1, x_2, \dots$	$\tilde{A} \oplus \tilde{B}$	Summation of two fuzzy sets $\tilde{A}$ and $\tilde{B}$
$\mathbb{R}$	Set of real numbers	$\tilde{A} \ominus \tilde{B}$	Difference of two fuzzy sets $\tilde{A}$ and $\tilde{B}$
$\mathbb{X}$	Universal set		
$\mathbb{Z}$	Set of all integer numbers	$\tilde{A} \otimes \tilde{B}$	Multiply of two fuzzy sets $\tilde{A}$ and $\tilde{B}$
$\mathbb{N}$	Set of all natural numbers	$\tilde{A} \oslash \tilde{B}$	Deviation of two fuzzy sets $\tilde{A}$ and $\tilde{B}$
$\mathbb{F}(\mathbb{X})$	Set of all fuzzy sets on $\mathbb{X}$		
$\mathbb{F}(\mathbb{R})$	Set of all fuzzy numbers on $\mathbb{R}$	$g(\tilde{A})$	Image of $\tilde{A}$ under the mapping $g(.)$
$\tilde{A}_{\alpha}$	$\alpha$ -value of a fuzzy number $\tilde{A}$	$\liminf \tilde{A}_n$	Limes inferior of a countable collection of fuzzy sets $\{\tilde{A}_n\}_{n \in \mathbb{N}}$
$\tilde{A}(x)$	Membership degree of $\tilde{A}$ at $x$		

$\limsup \tilde{A}_n$	Limes superior of a countable collection of fuzzy sets $\{\tilde{A}_n\}_{n \in \mathbb{N}}$	$P_X$ $(\mathbb{R}, \mathcal{F}_m(\mathbb{R}))$	Probability measure induced by $X$ Fuzzy measurable space
$\tilde{A}_1 \times \tilde{A}_2 \times \dots \times \tilde{A}_n$	Cartesian product of the fuzzy sets $\tilde{A}_1, \tilde{A}_2, \dots, \tilde{A}_n$	$P_X(\tilde{A})$	Exact probability of a fuzzy event of $\tilde{A}$
$S(\tilde{A}, \tilde{B})$	Similarity measure between two triangular fuzzy numbers of $\tilde{A}$ and $\tilde{B}$	$f_X^\theta$ $=^d$ $E_X$	Parametric density function of $X$ identical distribution function Expectation with respect to $X$
$P_d(\tilde{A} \succ \tilde{B})$	Preference degree that ' $\tilde{A}$ ' is greater than ' $\tilde{B}$ '	$\tilde{\theta}$ $P_X^\theta(\tilde{A})$	Fuzzy parameter Exact probability of $\tilde{A}$ with fuzzy parameter
$d(\tilde{A}, \tilde{B})$	Exact distance between two fuzzy numbers of $\tilde{A}$ and $\tilde{B}$	$\tilde{P}_X^\theta(\tilde{A})$	Fuzzy probability of $\tilde{A}$ with fuzzy parameter
$\tilde{D}(\tilde{A}, \tilde{B})$	Fuzzy distance between two fuzzy numbers of $\tilde{A}$ and $\tilde{B}$	$P_X(\tilde{A})$	Probability of a fuzzy product event $\tilde{A}$
$\tilde{A} \ominus_G \tilde{B}$	Generalized difference between two fuzzy numbers of $\tilde{A}$ and $\tilde{B}$	$P_X(\tilde{A} \tilde{B})$	Conditional probability of $\tilde{A}$ given $\tilde{B}$
$\max(\tilde{A}, \tilde{B})$	Maximum of the two fuzzy numbers of $\tilde{A}$ and $\tilde{B}$	$\tilde{x}$	Fuzzy arithmetic mean
$\widetilde{\min}(\tilde{A}, \tilde{B})$	Minimum of the two fuzzy numbers of $\tilde{A}$ and $\tilde{B}$	$\tilde{x}_g$	Fuzzy geometric mean
$  \tilde{A}  $	absolute value a fuzzy number of $\tilde{A}$	$\tilde{m}$ $\tilde{H}_p$ $\tilde{R}$	Fuzzy median Fuzzy quantile Fuzzy sample range
$P$	Probability measure	$\widetilde{IQR}$	Fuzzy interquartile range
$\Omega$	Set of outcomes of an experiment	$S_D$	Mean absolute deviation from the fuzzy mean
$\mathcal{A}$	A $\sigma$ -filed of $\Omega$		Standard deviation of a set of fuzzy data
$(\Omega, \mathcal{A}, P)$	Probability space		
$\mathcal{B}(\mathbb{R})$	Set of all Borel subsets of $\mathbb{R}$	$S^2$	
$X$	Random variable		

$\tilde{S}_k$	Fuzzy (Pearson) measure of skewness	$\tilde{X}_n \rightarrow^d \tilde{X}$	Converges in distribution
$m_t$	Exact measure of skewness	$\tilde{X}_n \rightarrow^{\tilde{L}^p} \tilde{X}$	converges in $\tilde{L}^p$
$g$	Measure of kurtosis of fuzzy data	<b>SLLN</b>	Strong Law of Large Numbers
<b>CDF</b>	Cumulative distribution function	<b>GFRV</b>	Generalized fuzzy random variable
$\tilde{X}$	Fuzzy random variable ( <b>FRV</b> )	<b>TFRV</b>	Triangular fuzzy random variable
$\tilde{E}(\tilde{X})$	Fuzzy expectation of $\tilde{X}$	$\tilde{H}_0$	Triangular normal fuzzy random variable
<b>FRS</b>	Ruzzy random sample	$\tilde{H}_1$	Fuzzy null hypothesis
$\text{var}(\tilde{X})$	Exact variance of $\tilde{X}$	$\tilde{T}$	Fuzzy alternative hypothesis
$\text{Cov}(\tilde{X}, \tilde{Y})$	Covariance of two <b>FRVs</b> of $\tilde{X}$ and $\tilde{Y}$	$D_R$	Fuzzy sign-test Statistics
$\rho(\tilde{X}, \tilde{Y})$	Covariance of two <b>FRVs</b> of $\tilde{X}$ and $\tilde{Y}$	$D_A$	Degree of rejection of $\tilde{H}_0$
$\tilde{P}(\tilde{X} \in (a, b))$	Fuzzy probability that $\tilde{X} \in (a, b)$	$\delta$	Degree of acceptance of $\tilde{H}_0$
$\liminf_{n \rightarrow \infty} \tilde{X}_n$	$\liminf$ of a sequence of <b>FRVs</b>	$\tilde{M}$	Significance level
$\limsup_{n \rightarrow \infty} \tilde{X}_n$	$\limsup$ of a sequence of <b>FRVs</b>	$\tilde{W}$	Fuzzy median
$\tilde{F}_{\tilde{X}}(x)$	$\{\tilde{X}_n\}_{n \geq 1}$	$\sqrt{n}\tilde{D}_n$	Fuzzy Wilcoxon test statistics
<b>LSFRV</b>	Fuzzy cumulative distribution function of $\tilde{X}$ at $x$	<b>FECDF</b>	Fuzzy one-sample Kolmogorov-Smirnov test
<b>SFRV</b>	( <b>FCDF</b> ) Location-scale fuzzy random variable	$\sqrt{\frac{mn}{m+n}}\tilde{D}_{mn}$	Fuzzy empirical cumulative distribution function
$\tilde{X}_n \rightarrow^c \tilde{X}$	Converges completely	$\tilde{H}$	Fuzzy Two-sample
$\tilde{X}_n \rightarrow^p \tilde{X}$	Converges in probability	$Cr$	Kolmogorov-Smirnov test
$\tilde{X}_n \rightarrow^{a.s} \tilde{X}$	Converges almost surely	ANOVA	Fuzzy Kruskal-Wallis test
		SST	Credibility measure
		SSR	Analysis of variance
			Sum of squares total of fuzzy data
			Sum of squares

	between variations of fuzzy data	$\tilde{d}_a(\tilde{A}, \tilde{B})$	Fuzzy triangular absolute error distance
SSE	Sum of squares within variations of fuzzy data	$\tilde{\bar{C}}(u, v)$	Fuzzy estimated value of $\tilde{C}(u, v)$
$\chi^2_r$	Chi-square distribution with $r$ degrees of freedom	<b>LTIFRV</b>	Life time fuzzy random variable
$t_r$	t-student distribution with $r$ degrees of freedom	<b>FRF</b>	Fuzzy reliability function
		<b>IMTTF</b>	Fuzzy mean time to failure
$F_{\nu_1, \nu_2}$	F-fisher distribution with $r$ degrees of freedom of $\nu_1$ and $\nu_2$	<b>FMRLF</b>	Fuzzy mean residual life function
$\chi^2_{r,\delta}$	$\delta$ -quantile of the chi-squared distribution with $r$ degrees of freedom	$\tilde{R}_S$	Fuzzy system reliability
$t_{r,\delta}$	$\delta$ -quantile of the t-distribution with $r$ degrees of freedom	MSE	Fuzzy intercept
		RMSE	Regression coefficient
$F_{\nu_1, \nu_2, \delta}$	$\delta$ -quantile of the f-distribution with $\nu_1$ and $\nu_2$ degrees of freedom	MSM	Mean square error
		$K(\cdot)$	Root mean square error
$\widetilde{LCL}$	Fuzzy lower central limit	$h$	Mean similarity measure
$\widetilde{CL}$	Fuzzy central limit	<b>ACF</b>	Kernel function
$\widetilde{UCL}$	Fuzzy upper central limit	<b>GCV</b>	Bandwidth
$\widetilde{x}$ -chart	Fuzzy $\bar{x}$ -chart	$\widetilde{\boldsymbol{X}}_T$	Generalized cross-validation criterion
s-chart	s-chart based on fuzzy random variables	$\text{cov}(\tilde{X}_t, \tilde{X}_{t-k})$	Autocorrelation function
$D_v$	Degree of violence	$\text{cov}(\tilde{X}_t, \tilde{X}_{t-k})$	Fuzzy time series data
<b>EWMA</b>	Exponentially-weighted moving average	$\hat{\boldsymbol{c}}_k$	Autocovariance of the generating fuzzy time series
$\widetilde{C}(u, v)$	Fuzzy process capability		Autocorrelation of the generating fuzzy time series sample autocovariance function of the generating fuzzy time series

## Part I

# Fuzzy Statistical Inferences Based on Fuzzy Random Variables



Taylor & Francis  
Taylor & Francis Group  
<http://taylorandfrancis.com>

# 1

---

## Introduction

---

In this chapter, the basic definitions and properties from fuzzy sets and fuzzy numbers needed for the book are recalled and discussed. We utilize such definitions and properties to produce some probabilistic reasoning and statistical inferences in the fuzzy domain the next chapter.

---

### 1.1 Fuzzy Set

Let  $\mathbb{X}$  be an universal in the universe of discourse and it includes all possible elements related with the given problem. The classical set theory is built on the fundamental concept of ‘set’. A set is defined as collection of objects, which share certain characteristics. A set  $A$  (of  $\mathbb{X}$ ) can be represented a set by enumerating its elements as  $A = \{x_1, x_2, \dots, x_n\}$ . There exists a membership function, which may be also used to define a set as a mapping of  $\mathbb{X}$  into  $\{0, 1\}$ . Such membership function for a set  $A$  can be represents by

$$A(x) = \begin{cases} 1, & x \in A, \\ 0, & x \notin A. \end{cases} \quad (1.1)$$

Therefore, each individual element in a set is a member or an element of the set. However, in real-life we come across many situations where membership and non-membership in a set are not clearly defined. For instance, consider the set of  $A = \{\text{all real numbers near to } 1\}$  in the context ‘positive values less than 5’. In classical set theory, one may imagine a membership function to describe  $A$  as

$$A(x) = \begin{cases} 1, & x \in [1 - a, 1 + a], \\ 0, & x \notin (1 - a, 1 + a). \end{cases} \quad (1.2)$$

where  $a \in (0, 1]$ . As a result, a value which is exactly  $a$  belongs to  $A$  and is considered to be ‘all real numbers near to 1’, but the values of  $[0, a - 0.001]$  do not belong to  $A$ . In addition, all values in  $[1 - a, 1 + a]$  may provide different memberships to describe ‘near to 1’. The value of 1 is completely consistence to ‘near to 1’ but the other values partially inherit the imprecise concept of ‘all real numbers near to 1’. These distinction are mathematically correct, but practically unreasonable. Therefore,  $A$  is not well defined in the sense

of classical set theory and cannot be precisely represented by 0 or 1. The ambiguity in this situation follows from that the boundaries of  $A$  is imprecise so as to characterize the  $A$  in a precise and rigorous way. For this purpose, instead of a sharp cut at the exact values of 0.9 and 1.1 (as well as all values in  $[1 - a, 1 + a]$ ), different memberships of ‘near to 1’ can be given via a membership function on  $[0, 1]$  instead of  $\{0, 1\}$ . In this regard, one may utilize the following membership function to describe ‘All numbers near to 1’ as

$$A(x) = \begin{cases} x, & 0 \leq x \leq 1, \\ 1 - x, & 1 \leq x \leq 2, \\ 0, & \mathbb{R} - [1, 2]. \end{cases} \quad (1.3)$$

For example, the value of 0.25 is considered to be ‘all real numbers near to 1’ with ‘degree 0.25’ and with ‘degree 0.95’ for the value of 0.25 and 0.95 according to the membership function of  $f$ . Therefore, we cannot exclude the all values in  $[0, 1]$  from  $A$ , nor include them completely. Thus  $A$  is a generalization of the classical characteristic function, which can be used to conclude that a value ‘is’ or ‘is not’ a member of ‘all real numbers near to 1’. Thus, such membership functions can provide a suitable mathematical tool for modeling such imprecision situations. Unlike the classical sets, the fuzzy set allows flexible membership degrees to each element of sets. That means that a fuzzy set contains elements that have varying degrees of membership in the set. Therefore, it provides a potential procedure to appreciate how uncertainty originating from human thinking can affect scientific problems.

Fuzzy sets were introduced by Zadeh [219] for the first time as an extension of classical sets. During the last two decades, fuzzy set theory has been successfully employed in working with numerous practical applications (for instance, see [12, 157, 185, 187, 223, 215]).

**Definition 1.1** ([41, 157, 187]) A fuzzy set of  $\mathbb{X}$  (universe of discourse) is a mapping  $\tilde{A} : \mathbb{X} \rightarrow [0, 1]$ , which assigns to each  $x \in \mathbb{X}$  a degree of membership  $0 \leq \tilde{A}(x) \leq 1$ . The set of all fuzzy sets on  $\mathbb{X}$  is denoted by  $\mathbb{F}(\mathbb{X})$ .

**Definition 1.2** ([41, 114, 157])  $\tilde{A}$  is said to be a convex fuzzy set on  $\mathbb{X}$  if and only if for all  $x, y \in \mathbb{X}$  and  $\lambda \in [0, 1]$ ,  $\tilde{A}(\lambda x + (1 - \lambda)y) \geq \min\{\tilde{A}(x), \tilde{A}(y)\}$ . A fuzzy set  $\tilde{A}$  is called normal fuzzy set if there exist  $x_0 \in \mathbb{X}$  such that  $\tilde{A}(x_0) = 1$ .

**Definition 1.3** ([41, 114, 157]) For each  $\alpha \in (0, 1]$ ,  $\{x \in \mathbb{X} \mid \tilde{A}(x) \geq \alpha\}$  of  $\mathbb{X}$  is called the  $\alpha$ -cut of  $\tilde{A}$  and is denoted by  $\tilde{A}[\alpha]$ . In addition,  $\tilde{A}[0] = Cl\{x \in \mathbb{X} : \tilde{A}(x) > 0\}$  is called the support of  $\tilde{A}$  where  $Cl(A)$  is the closure of  $A$  [140].

For every  $\alpha \in (0, 1]$ , the lower and upper bounds of  $\tilde{A}[\alpha]$  are denoted by  $\tilde{A}_\alpha^L$  and  $\tilde{A}_\alpha^U$ , respectively.

**Lemma 1.1** A fuzzy set  $\tilde{A}$  is convex if and only if  $\{\tilde{A}[\alpha]\}_{\alpha \in [0, 1]}$  is a collection of non-empty nested sets on  $\mathbb{X}$ .

**Proof** For a proof see [114].

Having a collection of non-empty nested sets  $\{\tilde{A}[\alpha]\}_{\alpha \in [0,1]}$  on  $\mathbb{X}$ , the membership function of  $\tilde{A}(x)$  can be evaluated by the following lemma.

**Lemma 1.2** Let  $\{\tilde{A}[\alpha]\}_{\alpha \in [0,1]}$  be a collection of non-empty nested sets on  $\mathbb{X}$ . Then:

$$\tilde{A}(x) = \sup\{\alpha \in [0, 1] : x \in \tilde{A}[\alpha]\}. \quad (1.4)$$

**Proof** See [114, 123, 157, 187] to prove this result.

**Definition 1.4** We say  $\tilde{A}$  is a discrete fuzzy set if it is convex, normal and  $\tilde{A}(x)$  is not zero only at a finite number of  $x \in \mathbb{X} = \{x_1, x_2, \dots, x_n\}$ . For such cases,  $\tilde{A}$  is typically denoted by  $\tilde{A} = \{\frac{a_1}{x_1}, \frac{a_2}{x_2}, \dots, \frac{a_n}{x_n}\}$  where  $\tilde{A}(x_i) = a_i \in (0, 1]$  are the membership degrees of  $x_i$ .

**Definition 1.5** A fuzzy set  $\tilde{A}$  is called positive fuzzy set ( $\tilde{A} > 0$ ) if  $\inf \tilde{A}[0] > 0$ , and it is negative fuzzy set ( $\tilde{A} < 0$ ) if  $\sup \tilde{A}[0] < 0$ .

**Example 1.1** Consider the set ‘small’ in  $\mathbb{X}$  consisting of natural numbers less than 10. Such an imprecise set can be described via the following fuzzy set:

$$\tilde{A} = \left\{ \frac{1}{1}, \frac{0.9}{2}, \frac{0.8}{3}, \frac{0.5}{4}, \frac{0.3}{5}, \frac{0.1}{6}, \frac{0}{7, 8, 9} \right\}.$$

For all  $x, y \in \mathbb{X}$  and  $\lambda \in [0, 1]$ , it is easy to check that  $\tilde{A}(\lambda x + (1 - \lambda)y) \geq \min\{\tilde{A}(x), \tilde{A}(y)\}$ .  $\tilde{A}$  is also a normal fuzzy set. Accordingly,  $\tilde{A}$  is a (discrete) fuzzy set.

Since an element can partially belong to a fuzzy set, a natural generalization of the classical notion of cardinality is to weigh each element by its membership degree, which resulted in the following definition for cardinality of a fuzzy set.

**Definition 1.6** [41, 114] For any fuzzy set  $\tilde{A}$  defined on a finite universal set  $\mathbb{X}$ , the cardinality of  $\tilde{A}$  is defined by  $|\tilde{A}| = \sum_{x \in \mathbb{X}} \tilde{A}(x)$ .

**Example 1.2** Let  $\mathbb{X} = \{1, 2, 3, 4, 5\}$  and  $\tilde{A} = \{\frac{0.3}{1}, \frac{0.5}{2}, \frac{0.7}{3}, \frac{1}{4}, \frac{0.8}{5}\}$ . Then  $|\tilde{A}| = \sum_x \tilde{A}(x) = 0.3 + 0.5 + 0.7 + 1 + 0.8 = 3.3$ .

The inclusion measure is an important concept in the area of fuzzy sets and indicates the degree to which a given fuzzy set belongs to a non-fuzzy set [34].

**Definition 1.7** Let  $\tilde{A} \in \mathbb{F}(\mathbb{X})$  be a fuzzy set and  $I$  be a subset of  $\mathbb{R}$ . Then, the degree to which  $\tilde{A}$  belongs to  $I$  is defined by  $d(\tilde{A} \in I) = \frac{\sum_{x \in I} \tilde{A}(x)}{|\tilde{A}|}$ .

Therefore,  $d(\tilde{A} \in I)$  is a relation between fuzzy sets  $\tilde{A}$  and classical set of  $I$  that indicates the degree to which  $\tilde{A}$  belongs to  $I$ .

**Lemma 1.3** Let  $\tilde{A} \in \mathbb{F}(\mathbb{X})$  and  $I \subseteq \mathbb{R}$ . Then:

- 1) If  $\{I_j\}_{j=1}^k$  is a collection of disjoint sets on  $\mathbb{X}$ , then  $\sum_{j=1}^k d(\tilde{A} \in I_j) = 1$ .
- 2)  $d(\tilde{A} \in I) = 1$  if and only if  $\tilde{A}[0] \subseteq I$ .

**Proof** Assuming  $\{I_j\}_{j=1}^k$  is a sequence of disjoint sets on  $\mathbb{X}$ , implies that

$$\begin{aligned} \sum_{j=1}^k d(\tilde{A} \in I_j) &= \frac{1}{|\tilde{A}|} \sum_{j=1}^k \sum_{x \in I_j} \tilde{A}(x) \\ &= \frac{1}{|\tilde{A}|} \sum_{x \in \bigcup_{j=1}^k I_j} \tilde{A}(x) \\ &= \frac{1}{|\tilde{A}|} |\tilde{A}| = 1. \end{aligned}$$

Now, if  $\tilde{A}[0] \subseteq I$  then:

$$d(\tilde{A} \in I) = \frac{\sum_{x \in \tilde{A}[0]} \tilde{A}(x) + \sum_{x \in (I - \tilde{A}[0])} \tilde{A}(x)}{|\tilde{A}|} = \frac{|\tilde{A}|}{|\tilde{A}|} = 1.$$

Conversely,  $d(\tilde{A} \in I) = 1$  yields  $\sum_{x \in I^c} \tilde{A}(x) = 0$  where  $I^c$  shows the complement set of  $I$ . This simply concludes that  $\tilde{A}[0] \subseteq I$  and verifies the proof.

**Remark 1.1** It should be pointed out that  $d(\tilde{A} \in I)$  may be interpreted as the probability that  $\tilde{A}$  belongs to  $I$ . Moreover, based on a given collection of disjoint sets  $\{I_j\}_{j=1}^k$ , from Lemma 1.3, it is natural to say that  $\tilde{A} \in I_*$  if  $d(\tilde{A} \in I_*) = \max_{j=1}^k d(\tilde{A} \in I_j)$ . In particular, assume that  $I$  and  $J$  are two sets of  $\mathbb{X}$  in which  $I \cup J = \mathbb{X}$  and  $I \cap J = \emptyset$ . Therefore, if  $d(\tilde{A} \in I) \geq 0.5$  then  $d(\tilde{A} \in J) < 0.5$ . In this case, one can say  $\tilde{A}$  belongs to  $I$ .

**Example 1.3** Let  $\tilde{A} = \{\frac{0.4}{3}, \frac{0.6}{4}, \frac{1}{5}, \frac{0.8}{6}, \frac{0.5}{7}\}$  be a fuzzy set on  $\mathbb{X} = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$ . Then, assuming  $I_1 = \{1, 2, 3, 4\}$ ,  $I_2 = \{5, 6\}$ , and  $I_3 = \{7, 8, 9, 10\}$  implies that  $d(\tilde{A} \in I_1) = 0.303$ ,  $d(\tilde{A} \in I_2) = 0.545$ , and  $d(\tilde{A} \in I_3) = 0.152$ . Since  $d(\tilde{A} \in I_2) \geq 0.5$ , we get that  $\tilde{A} \in I_2$ .

### 1.1.1 Operations of fuzzy sets

Although the set-theoretic operations (union, intersection, and complement) possess some rigorous axiomatic properties, the conventional operations of sets can be extended for fuzzy sets [12, 157, 185, 187, 215, 223]. In this section, some well-established operations of fuzzy sets are recalled and discussed.

**Definition 1.8** For two fuzzy sets  $\tilde{A}$  and  $\tilde{B}$ , we say:

1.  $\tilde{A}$  is included in  $\tilde{B}$ , say  $\tilde{A} \subseteq \tilde{B}$ , if  $\tilde{A}(x) \leq \tilde{B}(x)$ , for all  $x \in \mathbb{X}$ .
2.  $\emptyset$  is the empty (fuzzy) set if  $\emptyset(x) = 0$  for any  $x \in \mathbb{X}$ .
3.  $\tilde{A}^c$  is the complement of fuzzy set  $\tilde{A}$  with the membership function  $\tilde{A}^c(x) = 1 - \tilde{A}(x)$ , for all  $x \in \mathbb{X}$ .
4.  $\tilde{A} \cup \tilde{B}$  is the union of fuzzy sets  $\tilde{A}$  and  $\tilde{B}$  with the membership function  $(\tilde{A} \cup \tilde{B})(x) = \max\{\tilde{A}(x), \tilde{B}(x)\}$ , for all  $x \in \mathbb{X}$ .
5.  $\tilde{A} \cap \tilde{B}$  is the intersection of fuzzy sets  $\tilde{A}$  and  $\tilde{B}$  with the membership function  $(\tilde{A} \cap \tilde{B})(x) = \min\{\tilde{A}(x), \tilde{B}(x)\}$ , for all  $x \in \mathbb{X}$ .
6.  $\tilde{A} - \tilde{B}$  is the difference between two fuzzy sets  $\tilde{A}$  and  $\tilde{B}$  with the membership function of  $(\tilde{A} - \tilde{B})(x) = \max\{0, \tilde{A}(x) - \tilde{B}(x)\}$ .
7.  $\tilde{A} \odot \tilde{B}$  is the product of two fuzzy sets  $\tilde{A}$  and  $\tilde{B}$  with the membership function of  $(\tilde{A} \odot \tilde{B})(x) = \tilde{A}(x)\tilde{B}(x)$ .
8.  $\tilde{A} \Delta \tilde{B}$  is the symmetric difference of  $\tilde{A}$  and  $\tilde{B}$  with the membership function of  $(\tilde{A} \Delta \tilde{B})(x) = |\tilde{A}(x) - \tilde{B}(x)|$ .

**Example 1.4** Let  $\mathbb{X} = \{1, 2, 3, 4\}$ ,  $\tilde{A} = \left\{ \frac{0.2}{1}, \frac{0.7}{2}, \frac{1}{3}, \frac{0.6}{4} \right\}$ , and  $\tilde{B} = \left\{ \frac{0.9}{1}, \frac{1}{2}, \frac{0.8}{3}, \frac{0.5}{4} \right\}$ . Then, it can be easy to verify that:

$$\begin{aligned}\tilde{A}^c &= \left\{ \frac{0.8}{1}, \frac{0.3}{2}, \frac{0}{3}, \frac{0.4}{4} \right\}, \\ \tilde{A} \cup \tilde{B} &= \left\{ \frac{\max\{0.2, 0.9\}}{1}, \frac{\max\{0.7, 1\}}{2}, \frac{\max\{1, 0.8\}}{3}, \frac{\max\{0.6, 0.5\}}{4} \right\} \\ &= \left\{ \frac{0.9}{1}, \frac{1}{2}, \frac{1}{3}, \frac{0.6}{4} \right\},\end{aligned}$$

$$\begin{aligned}\tilde{A} \cap \tilde{B} &= \left\{ \frac{\min\{0.2, 0.9\}}{1}, \frac{\min\{0.7, 1\}}{2}, \frac{\min\{1, 0.8\}}{3}, \frac{\min\{0.6, 0.5\}}{4} \right\} \\ &= \left\{ \frac{0.2}{1}, \frac{0.7}{2}, \frac{0.8}{3}, \frac{0.5}{4} \right\},\end{aligned}$$

$$\begin{aligned}\tilde{A} - \tilde{B} &= \left\{ \frac{\max\{0, 0.2 - 0.9\}}{1}, \frac{\max\{0, 0.7 - 1\}}{2}, \frac{\max\{0, 1 - 0.8\}}{3}, \right. \\ &\quad \left. \frac{\max\{0, 0.6 - 0.5\}}{4} \right\} \\ &= \left\{ \frac{0}{1}, \frac{0}{2}, \frac{0.2}{3}, \frac{0.1}{4} \right\},\end{aligned}$$

$$\begin{aligned}\tilde{A} \odot \tilde{B} &= \left\{ \frac{0.2 \times 0.9}{1}, \frac{0.7 \times 1}{2}, \frac{1 \times 0.8}{3}, \frac{0.6 \times 0.5}{4} \right\} \\ &= \left\{ \frac{0.18}{1}, \frac{0.7}{2}, \frac{0.8}{3}, \frac{0.03}{4} \right\},\end{aligned}$$

$$\begin{aligned}\tilde{A} \triangle \tilde{B} &= \left\{ \frac{|0.2 - 0.9|}{1}, \frac{|0.7 - 1|}{2}, \frac{|1 - 0.8|}{3}, \frac{|0.6 - 0.5|}{4} \right\} \\ &= \left\{ \frac{0.7}{1}, \frac{0.3}{2}, \frac{0.2}{3}, \frac{0.1}{4} \right\}.\end{aligned}$$

**Lemma 1.4** For three fuzzy sets  $\tilde{A}$ ,  $\tilde{B}$ , and  $\tilde{C}$ ,

1. Distributivity:

$$\tilde{A} \cap (\tilde{B} \cup \tilde{C}) = (\tilde{A} \cap \tilde{B}) \cup (\tilde{A} \cap \tilde{C}), \quad (1.5)$$

$$\tilde{A} \cup (\tilde{B} \cap \tilde{C}) = (\tilde{A} \cup \tilde{B}) \cap (\tilde{A} \cup \tilde{C}). \quad (1.6)$$

2. De Morgan Laws:

$$(\tilde{A} \cup \tilde{B})^c = \tilde{A}^c \cap \tilde{B}^c, \quad (1.7)$$

$$(\tilde{A} \cap \tilde{B})^c = \tilde{A}^c \cup \tilde{B}^c. \quad (1.8)$$

**Proof** To see a proof of these results see [41, 114].

**Theorem 1.1** For three fuzzy sets  $\tilde{A}$ ,  $\tilde{B}$ , and  $\tilde{C}$ , the following hold.

- 1)  $\tilde{A} \odot (\tilde{B} \cup \tilde{C}) = (\tilde{A} \odot \tilde{B}) \cup (\tilde{A} \odot \tilde{C})$ .
- 2)  $\tilde{A} \odot (\tilde{B} \cap \tilde{C}) = (\tilde{A} \odot \tilde{B}) \cap (\tilde{A} \odot \tilde{C})$ .
- 3)  $\tilde{A} \odot (\tilde{B} - \tilde{C}) = (\tilde{A} \odot \tilde{B}) - (\tilde{A} \odot \tilde{C})$ .

**Proof** Note that  $\tilde{A} \odot (\tilde{B} \cup \tilde{C})(x) = \tilde{A}(x) \max\{\tilde{B}(x), \tilde{C}(x)\} = \max\{\tilde{A}(x) \tilde{B}(x), \tilde{A}(x) \tilde{C}(x)\}$ . This verifies part (1). The second part can be shown similarly. Further, we have  $(\tilde{A} \odot (\tilde{B} - \tilde{C}))(x) = \tilde{A}(x) \max\{0, \tilde{B}(x) - \tilde{C}(x)\} = \max\{0, \tilde{A}(x) \tilde{B}(x) - \tilde{A}(x) \tilde{C}(x)\}$ . This concludes that  $\tilde{A} \odot (\tilde{B} - \tilde{C}) = (\tilde{A} \odot \tilde{B}) - (\tilde{A} \odot \tilde{C})$ .

**Theorem 1.2** For two fuzzy sets  $\tilde{A}$  and  $\tilde{B}$  and for every  $\alpha \in (0, 1]$ , the following results show the connections between fuzzy sets and their  $\alpha$ -cuts:

1. If  $\tilde{A} \subseteq \tilde{B}$  then  $\tilde{A}[\alpha] \subseteq \tilde{B}[\alpha]$ .
2.  $\tilde{A}^c[\alpha] = (\tilde{A}[1 - \alpha])^c$  where  $\tilde{A}[1 - \alpha] = \{x : \tilde{A}(x) > 1 - \alpha\}$ .

$$3. (\tilde{A} \cup \tilde{B})[\alpha] = \tilde{A}[\alpha] \cup \tilde{B}[\alpha].$$

$$4. (\tilde{A} \cap \tilde{B})[\alpha] = \tilde{A}[\alpha] \cap \tilde{B}[\alpha].$$

**Proof** For a comprehensive proof see [41, 114].

**Lemma 1.5** For two fuzzy sets  $\tilde{A}$  and  $\tilde{B}$ ,  $|\tilde{A} \cup \tilde{B}| = |\tilde{A}| + |\tilde{B}| - |\tilde{A} \cap \tilde{B}|$ .

**Proof** According to the definition of  $\tilde{A} \cup \tilde{B}$ , we easily have:

$$\begin{aligned} |\tilde{A} \cup \tilde{B}| &= \sum_{x \in \mathbb{R}} (\tilde{A} \cup \tilde{B})(x) \\ &= \sum_{x \in \mathbb{R}} (\tilde{A}(x) + \tilde{B}(x) - (\tilde{A} \cap \tilde{B})(x)) \\ &= |\tilde{A}| + |\tilde{B}| - |\tilde{A} \cap \tilde{B}|. \end{aligned}$$

**Definition 1.9** (Extension principle [41, 114]) Suppose that  $g : \mathbb{X} \rightarrow \mathbb{Y}$  is a real-valued function and  $\tilde{A}$  is a fuzzy set on  $\mathbb{F}(\mathbb{X})$ . The image of the fuzzy set  $\tilde{A}$  under the mapping  $g(.)$  can be expressed as a fuzzy set  $g(\tilde{A}) \in \mathbb{F}(\mathbb{Y})$  with the following membership function:

$$g(\tilde{A})(y) = \begin{cases} \sup_{x:y=g(x)} \tilde{A}(x), & g^{-1}(y) \neq \emptyset, \\ 0, & g^{-1}(y) = \emptyset. \end{cases} \quad (1.9)$$

**Example 1.5** Consider a fuzzy set on  $\mathbb{Z} = \{\dots, -2, -1, 0, 1, 2, \dots\}$  stands for ‘about 1’ with the membership function of  $\tilde{A} = \{\frac{0.4}{-2}, \frac{0.6}{-1}, \frac{0.8}{0}, \frac{1}{1}, \frac{0.8}{2}, \frac{0.6}{3}, \frac{0.4}{4}\}$ . Letting  $g(x) = x^4$ , we have  $y \in \{0, 1, 16, 81, 256\}$  and thus:

$$\begin{aligned} g(\tilde{A})(0) &= \tilde{A}(0) = 0.8, \\ g(\tilde{A})(1) &= \max\{\tilde{A}(-1), \tilde{A}(1)\} = \max\{0.6, 1\} = 1, \\ g(\tilde{A})(16) &= \max\{\tilde{A}(-2), \tilde{A}(2)\} = \max\{0.4, 0.8\} = 0.8, \\ g(\tilde{A})(81) &= \tilde{A}(3) = 0.6, \\ g(\tilde{A})(256) &= \tilde{A}(4) = 0.4. \end{aligned}$$

From above, the membership function of  $g(\tilde{A}) = (\tilde{A})^4$  can be obtained as:

$$(\tilde{A})^4 = \left\{ \frac{0.8}{0}, \frac{1}{1}, \frac{0.8}{16}, \frac{0.6}{81}, \frac{0.4}{256} \right\}.$$

In order to use fuzzy sets in any real-life one should be able to perform arithmetic operations including addition, subtraction, multiplication, and division in computational process, which is called fuzzy arithmetic. The usual arithmetic operations on real numbers can be extended to the ones defined on fuzzy sets based on the Extension principle.

**Definition 1.10** [41, 114] The arithmetic operations between two fuzzy sets of  $\tilde{A}$  and  $\tilde{B}$  is defined as a fuzzy set  $\tilde{A} \circ \tilde{B}$  with the following membership function:

$$(\tilde{A} \circ \tilde{B})(z) = \sup_{x,y:z=x \bullet y} \min\{\tilde{A}(x), \tilde{B}(y)\}. \quad (1.10)$$

The symbols  $\circ = \oplus, \ominus, \otimes, \oslash$  denotes the extended arithmetic operations  $\bullet = +, -, \times, /$  in a fuzzy domain.

Next, a notion of convergence a collection of fuzzy sets is given.

**Definition 1.11** Let  $\{\tilde{A}_n\}_{n \in \mathbb{N}}$  be a countable collection of fuzzy sets and  $\tilde{A}$  be a fuzzy set. Then, we say:

1.  $\tilde{A}_n \rightarrow \tilde{A}$  if  $\lim_{n \rightarrow \infty} \tilde{A}_n(x) = \tilde{A}(x)$  for all  $x \in \mathbb{R}$ .
2.  $\tilde{A}_n \uparrow \tilde{A}$  if  $\{\tilde{A}_n\}_{n \in \mathbb{N}}$  is an increasing countable collection of fuzzy sets and  $\lim_{n \rightarrow \infty} \tilde{A}_n(x) = \tilde{A}(x)$  for all  $x \in \mathbb{R}$ .
3.  $\tilde{A}_n \downarrow \tilde{A}$  if  $\{\tilde{A}_n\}_{n \in \mathbb{N}}$  is a decreasing countable collection of fuzzy sets and  $\lim_{n \rightarrow \infty} \tilde{A}_n(x) = \tilde{A}(x)$  for all  $x \in \mathbb{R}$ .

**Example 1.6** Let  $\mathbb{X} = \{-3, -2, -1, 0, 1, 2, 3\}$  and consider a sequence of discrete fuzzy set on  $\mathbb{X}$  with  $\tilde{A}_n(x) = (1 - (1/nx^2))^n$ ,  $x \in \mathbb{X}$ . Since  $\lim_{n \rightarrow \infty} \tilde{A}_n(x) = e^{-x^2}$ , we have  $\lim_{n \rightarrow \infty} \tilde{A}_n(x) = \tilde{A}(x) = e^{-x^2}$  with the following membership function:

$$\tilde{A} = \left\{ \frac{-3}{e^{-9}}, \frac{-2}{e^{-4}}, \frac{-1}{e^{-1}}, \frac{0}{1}, \frac{1}{e^{-1}}, \frac{2}{e^{-4}}, \frac{3}{e^{-9}} \right\}.$$

**Proposition 1.1** For a countable collection of fuzzy sets  $\{\tilde{A}_n\}_{n \in \mathbb{N}}$ ,  $\tilde{A}_n \rightarrow \tilde{A}$  if and only if  $\tilde{A}_n[\alpha] \rightarrow \tilde{A}[\alpha]$  for any  $\alpha \in (0, 1]$ .

**Proof** The proof is left for the reader.

**Definition 1.12** The limes superior and limes inferior of a countable collection of fuzzy sets  $\{\tilde{A}_n\}_{n \in \mathbb{N}}$  are defined to be fuzzy sets  $\limsup \tilde{A}_n = \bigcap_{n \geq 1} \bigcup_{m \geq n} \tilde{A}_m$  and  $\liminf \tilde{A}_n = \bigcup_{n \geq 1} \bigcap_{m \geq n} \tilde{A}_m$ , respectively.

**Example 1.7** Let  $\mathbb{X} = \{-3/4, -2/4, -1/4, 0, 1/4, 2/4, 3/4\}$  and consider a sequence of discrete fuzzy set on  $\mathbb{X}$  with  $\tilde{A}_n(x) = \max\{0, 1 - (n/(n+1))|x|\}$ ,  $x \in \mathbb{X}$ . Then, for a fixed  $x \in \mathbb{X}$ , we have

$$\begin{aligned} (\bigcup_{m \geq n} \tilde{A}_m)(x) &= \max_{m \geq n} \tilde{A}_m(x) \\ &= \max\{0, \max_{m \geq n} (1 - (m/(m+1))|x|)\} \\ &= \max\{0, 1 - (n/(n+1))|x|\}. \end{aligned}$$

This concludes that

$$\begin{aligned}
 (\limsup \tilde{A}_n)(x) &= (\cap_{n \geq 1} \cup_{m \geq n} \tilde{A}_m)(x) \\
 &= \min_{n \geq 1} (\cup_{m \geq n} \tilde{A}_m)(x) \\
 &= \min_{n \geq 1} \max\{0, 1 - (n/(n+1))|x|\} \\
 &= \max\{0, 1 - |x|\} = \tilde{A}(x).
 \end{aligned}$$

Therefore,

$$\limsup \tilde{A}_n = \left\{ \frac{-3/4}{1/4}, \frac{-2/4}{2/4}, \frac{-1/4}{3/4}, \frac{0}{1}, \frac{1/4}{3/4}, \frac{2/4}{2/4}, \frac{3/4}{1/4} \right\}.$$

**Theorem 1.3** Let  $\{\tilde{A}_n\}_{n \in \mathbb{N}}$  be a countable collection of fuzzy sets. Then, the following are valid.

1.  $\liminf \tilde{A}_n \subseteq \limsup \tilde{A}_n$ .
2.  $(\liminf \tilde{A}_n)^c = \limsup \tilde{A}_n^c$ .
3.  $(\limsup \tilde{A}_n)^c = \liminf \tilde{A}_n^c$ .

**Proof** First note that  $(\liminf \tilde{A}_n)[\alpha] = \cup_{n \geq 1} \cap_{m \geq n} \tilde{A}_n[\alpha]$ . That concludes that  $(\liminf \tilde{A}_n)[\alpha] \subseteq \cap_{n \geq 1} \cup_{m \geq n} \tilde{A}_n[\alpha] = (\limsup \tilde{A}_n)[\alpha]$  for any  $\alpha \in [0, 1]$  or  $\liminf \tilde{A}_n \subseteq \limsup \tilde{A}_n$ . In addition, for any  $\alpha \in [0, 1]$ ,  $(\liminf \tilde{A}_n)^c[\alpha] = (\cup_{n \geq 1} \cap_{m \geq n} \tilde{A}_n[1-\alpha])^c = \cap_{n \geq 1} \cup_{m \geq n} (\tilde{A}_n[1-\alpha])^c = \cap_{n \geq 1} \cup_{m \geq n} \tilde{A}_n^c[\alpha]$ . This means  $(\liminf \tilde{A}_n)^c = \limsup \tilde{A}_n^c$ . Similarly, it can be shown that  $(\limsup \tilde{A}_n)^c = \liminf \tilde{A}_n^c$ .

**Theorem 1.4** Let  $\{\tilde{A}_n\}_{n \in \mathbb{N}}$  and  $\{\tilde{B}_n\}_{n \in \mathbb{N}}$  be two countable collections of fuzzy sets. Then:

- 1)  $\limsup(\tilde{A} \oplus \tilde{B})_n \subseteq (\limsup \tilde{A}_n \oplus \limsup \tilde{B}_n)$ .
- 2)  $\liminf(\tilde{A} \oplus \tilde{B})_n \supseteq (\liminf \tilde{A}_n \oplus \liminf \tilde{B}_n)$ .

**Proof** For any  $\alpha \in (0, 1]$ ,  $\limsup(\tilde{A}_n \oplus \tilde{B}_n)[\alpha] = \cap_{n \geq 1} \cup_{m \geq n} (\tilde{A}_n[\alpha] + \tilde{B}_n[\alpha])$ . Now, assume that  $x$  is an arbitrary element of  $\limsup(\tilde{A}_n \oplus \tilde{B}_n)[\alpha]$ . Therefore, for any  $n \geq 1$ , there exist a  $m \geq n$  such that  $x = y + z$  where  $y \in \tilde{A}_n[\alpha]$  and  $z \in \tilde{B}_n[\alpha]$ . This concludes that  $x \in (\limsup \tilde{A}_n \oplus \limsup \tilde{B}_n)[\alpha]$  which providing (1). The second part can be verified similarly.

**Proposition 1.2** For a countable collection of fuzzy sets  $\{\tilde{A}_n\}_{n \in \mathbb{N}}$ :

- 1)  $\lim_{n \rightarrow \infty} \cup_{k \geq n} \tilde{A}_k = \limsup \tilde{A}_n$ .
- 2)  $\lim_{n \rightarrow \infty} \cap_{k \geq n} \tilde{A}_k = \liminf \tilde{A}_n$ .

**Proof** Let  $\tilde{B}_k^n = \cup_{k \geq n} \tilde{A}_k$ , for any  $\alpha \in (0, 1]$ . For (1), since  $\tilde{B}_k^n[\alpha] \downarrow \cap_{n \geq 1} \cup_{k \geq n} \tilde{A}_k[\alpha] = \limsup \tilde{A}_n[\alpha]$ , we easily have  $\lim_{n \rightarrow \infty} \cup_{k \geq n} \tilde{A}_k = \limsup \tilde{A}_n$  by Lemma 1.1. The second part can be verified similarly.

**Proposition 1.3** Let  $\{\tilde{A}_n\}_{n \in \mathbb{N}}$  be a countable collection of fuzzy sets. Then  $\tilde{A}_n \rightarrow \tilde{A}$  if and only if  $\liminf \tilde{A}_n = \limsup \tilde{A}_n = \tilde{A}$ .

**Proof** First assume that  $\tilde{A}_n \rightarrow \tilde{A}$ . This simply concludes that  $\tilde{A}_n[\alpha] \rightarrow \tilde{A}[\alpha]$  for any  $\alpha \in (0, 1]$  which implies that  $\liminf \tilde{A}_n[\alpha] = \limsup \tilde{A}_n = \tilde{A}[\alpha]$  for any  $\alpha \in [0, 1]$ . Thus  $\liminf \tilde{A}_n = \limsup \tilde{A}_n = \tilde{A}$  and thus the ‘if part’ is verified. The ‘only if part’ can be shown similarly.

Next, a common notion of a real-valued function of a fuzzy set is given.

### 1.1.2 Cartesian product of fuzzy sets

The Cartesian products of some sets mean the product of some non-empty sets in a systematic way. Here, a common definition of the fuzzy product of (discrete) fuzzy is recalled, and some of its main properties are investigated. All such concepts were gathered from [100].

**Definition 1.13** Let  $\tilde{A}_i \in \mathbb{F}(\mathbb{X}_i)$ ,  $i = 1, 2, \dots, n$ . A fuzzy product set on  $\mathbb{X} = \mathbb{X}_1 \times \mathbb{X}_2 \times \dots \times \mathbb{X}_n$  is defined as a Cartesian product of the fuzzy set  $\tilde{\mathbf{A}} = \tilde{A}_1 \times \tilde{A}_2 \times \dots \times \tilde{A}_n$  with the following membership function:

$$\tilde{\mathbf{A}}(\mathbf{x}) = \min_{i=1}^n \{\tilde{A}_i(x_i)\}, \quad x_i \in \mathbb{X}_i. \quad (1.11)$$

**Example 1.8** Let  $\mathbb{X}_1 = \mathbb{X}_2 = \{2, 4, 6\}$ ,  $\tilde{A}_1 = \{\frac{0.7}{2}, \frac{1}{4}, \frac{0.6}{6}\}$  and  $\tilde{A}_2 = \{\frac{1}{2}, \frac{0.8}{4}\}$ . Then, the membership function of fuzzy product of  $\tilde{A}_1$  and  $\tilde{A}_2$  can be shown by:

$$\tilde{A}_1 \times \tilde{A}_2 = \left\{ \frac{0.7}{(2, 2)}, \frac{0.7}{(2, 4)}, \frac{1}{(4, 2)}, \frac{0.8}{(4, 4)}, \frac{0.6}{(6, 2)}, \frac{0.6}{(6, 4)} \right\}.$$

**Proposition 1.4** If  $\{\tilde{\mathbf{A}}_j = \tilde{A}_{1j} \times \tilde{A}_{2j} \times \dots \times \tilde{A}_{nj}\}_{j=1}^m$  is a sequence of fuzzy product sets then  $\bigcup_{j=1}^m \tilde{\mathbf{A}}_j = (\bigcup_{j=1}^m \tilde{A}_{1j}) \times \dots \times (\bigcup_{j=1}^m \tilde{A}_{nj})$ .

**Proof** For a fixed  $\mathbf{x} = (x_1, x_2, \dots, x_n) \in \mathbb{X}^n$ , we have:

$$\begin{aligned} \left( \bigcup_{j=1}^m \tilde{\mathbf{A}}_j \right) (\mathbf{x}) &= \max_{j=1}^m \min \{\tilde{A}_{1j}(x_1), \tilde{A}_{2j}(x_2), \dots, \tilde{A}_{nj}(x_n)\} \\ &= \min \left\{ \max_{j=1}^m \tilde{A}_{1j}(x_1), \max_{j=1}^m \tilde{A}_{2j}(x_2), \dots, \max_{j=1}^m \tilde{A}_{nj}(x_n) \right\} \\ &= \left( \bigcup_{j=1}^m \tilde{A}_{1j} \right) (x_1) \times \left( \bigcup_{j=1}^m \tilde{A}_{2j} \right) (x_2) \times \dots \times \left( \bigcup_{j=1}^m \tilde{A}_{nj} \right) (x_n). \end{aligned} \quad (1.12)$$

It completes the proof.

**Proposition 1.5** If  $\{\tilde{A}_j = \tilde{A}_{1j} \times \tilde{A}_{2j} \times \dots \times \tilde{A}_{nj}\}_{j=1}^m$  is a sequence of fuzzy product sets then  $\bigcap_{j=1}^m \tilde{A}_j = (\bigcap_{j=1}^m \tilde{A}_{1j}) \times \dots \times (\bigcap_{j=1}^m \tilde{A}_{nj})$ .

**Proof** For a fixed  $\mathbf{x} = (x_1, x_2, \dots, x_n) \in \mathbb{X}^n$ , it is easily seen that:

$$\begin{aligned} \left(\bigcap_{j=1}^m \tilde{A}_j\right)(\mathbf{x}) &= \min_{j=1}^m \min\{\tilde{A}_{1j}(x_1), \tilde{A}_{2j}(x_2), \dots, \tilde{A}_{nj}(x_n)\} = \\ &= \min\{\min_{j=1}^m \tilde{A}_{1j}(x_1), \min_{j=1}^m \tilde{A}_{2j}(x_2), \dots, \min_{j=1}^m \tilde{A}_{nj}(x_n)\} = \\ &= \left(\bigcap_{j=1}^m \tilde{A}_{1j}\right)(x_1) \times \left(\bigcap_{j=1}^m \tilde{A}_{2j}\right)(x_2) \times \dots \times \left(\bigcap_{j=1}^m \tilde{A}_{nj}\right)(x_n). \end{aligned}$$

**Lemma 1.6** Let  $\{\tilde{A}_i\}_{i=1}^n$  be a finite collection of fuzzy sets. For a given fuzzy set  $\tilde{B}$ , we have the following results:

1.  $(\cup_{i=1}^n \tilde{A}_i) \times \tilde{B} = \cup_{i=1}^n (\tilde{A}_i \times \tilde{B})$ .
2.  $\tilde{B} \times (\cup_{i=1}^n \tilde{A}_i) = \cup_{i=1}^n (\tilde{B} \times \tilde{A}_i)$ .
3.  $(\cap_{i=1}^n \tilde{A}_i) \times \tilde{B} = \cap_{i=1}^n (\tilde{A}_i \times \tilde{B})$ .
4.  $\tilde{B} \times (\cap_{i=1}^n \tilde{A}_i) = \cap_{i=1}^n (\tilde{B} \times \tilde{A}_i)$ .

**Proof** The first part is proved only. The others are left for readers. For this purpose, a fixed  $\mathbf{x} = (x_1, x_2, \dots, x_n) \in \mathbb{X}^n$  and  $x \in \mathbb{X}$ , we have:

$$\begin{aligned} ((\cup_{i=1}^n \tilde{A}_i) \times \tilde{B})(\mathbf{x}, x) &= \min\{\max\{\tilde{A}_1(x_1), \tilde{A}_2(x_2), \dots, \tilde{A}_n(x_n)\}, \tilde{B}(x)\} = \\ &= \max\{\min\{\tilde{A}_1(x_1), \tilde{B}(x)\}, \dots, \min\{\tilde{A}_n(x_n), \tilde{B}(x)\}\} = (\cup_{i=1}^n (\tilde{A}_i \times \tilde{B}))(\mathbf{x}, x). \end{aligned}$$

This confirm that  $(\cup_{i=1}^n \tilde{A}_i) \times \tilde{B} = \cup_{i=1}^n (\tilde{A}_i \times \tilde{B})$ .

---

## 1.2 Fuzzy Numbers

Fuzzy numbers generalize classical real numbers and roughly speaking a fuzzy number is a fuzzy set of the real line that has some additional properties. That is the domain of a fuzzy number is a specified set of real numbers and its range is the span of non-negative real numbers between 0 and 1. Each numerical value in the domain is assigned a specific degree of membership where 0 represents the smallest possible grade, and 1 is the largest possible degree. A unified definition of a fuzzy number is given below.

**Definition 1.14 ([41, 114, 157])** A fuzzy set  $\tilde{A}$  of  $\mathbb{R}$  is called a fuzzy number (**FN**) if it is normal and the set  $\tilde{A}[\alpha] = \{x \in \mathbb{R} : \tilde{A}(x) \geq \alpha\}$  is a non-empty nested closed interval in  $\mathbb{R}$ , for every  $\alpha \in [0, 1]$ .

This interval is denoted by  $\tilde{A}[\alpha] = [\tilde{A}_\alpha^L, \tilde{A}_\alpha^U]$ , where  $\tilde{A}_\alpha^L = \inf\{x : x \in \tilde{A}[\alpha]\}$  and  $\tilde{A}_\alpha^U = \sup\{x : x \in \tilde{A}[\alpha]\}$ . For this book, take  $\mathcal{F}(\mathbb{R})$  as denoting the set of all **FNs**.

The arithmetic operations between two **FNs**  $\tilde{A}$  and  $\tilde{B}$  can also be evaluated using arithmetic operations of their  $\alpha$ -cuts.

**Theorem 1.5** *Let  $\tilde{A}$  and  $\tilde{B}$  be two **FNs**. Then, for every  $\alpha \in (0, 1)$ :*

$$1. (\tilde{A} \oplus \tilde{B})[\alpha] = [\tilde{A}_\alpha^L + \tilde{B}_\alpha^L, \tilde{A}_\alpha^U + \tilde{B}_\alpha^U].$$

$$2. (\tilde{A} \ominus \tilde{B})[\alpha] = [\tilde{A}_\alpha^L - \tilde{B}_\alpha^U, \tilde{A}_\alpha^U - \tilde{B}_\alpha^L].$$

$$3. (\tilde{A} \otimes \tilde{B})[\alpha] = [(\tilde{A} \otimes \tilde{B})_\alpha^L, (\tilde{A} \otimes \tilde{B})_\alpha^U] \text{ where}$$

$$(\tilde{A} \otimes \tilde{B})_\alpha^L = \min\{\tilde{A}_\alpha^L \tilde{B}_\alpha^L, \tilde{A}_\alpha^L \tilde{B}_\alpha^U, \tilde{A}_\alpha^U \tilde{B}_\alpha^L, \tilde{A}_\alpha^U \tilde{B}_\alpha^U\}, \quad (1.13)$$

$$(\tilde{A} \otimes \tilde{B})_\alpha^U = \max\{\tilde{A}_\alpha^L \tilde{B}_\alpha^L, \tilde{A}_\alpha^L \tilde{B}_\alpha^U, \tilde{A}_\alpha^U \tilde{B}_\alpha^L, \tilde{A}_\alpha^U \tilde{B}_\alpha^U\}, \quad (1.14)$$

$$4. (\tilde{A} \oslash \tilde{B})[\alpha] = [(\tilde{A} \oslash \tilde{B})_\alpha^L, (\tilde{A} \oslash \tilde{B})_\alpha^U] \text{ where}$$

$$(\tilde{A} \oslash \tilde{B})_\alpha^L = \min\{\tilde{A}_\alpha^L / \tilde{B}_\alpha^L, \tilde{A}_\alpha^L / \tilde{B}_\alpha^U, \tilde{A}_\alpha^U / \tilde{B}_\alpha^L, \tilde{A}_\alpha^U / \tilde{B}_\alpha^U\}, \quad (1.15)$$

$$(\tilde{A} \oslash \tilde{B})_\alpha^U = \max\{\tilde{A}_\alpha^L / \tilde{B}_\alpha^L, \tilde{A}_\alpha^L / \tilde{B}_\alpha^U, \tilde{A}_\alpha^U / \tilde{B}_\alpha^L, \tilde{A}_\alpha^U / \tilde{B}_\alpha^U\}, \quad (1.16)$$

provided that  $0 \notin \tilde{B}[0]$ .

**Proof** See [41, 114, 157, 187], for instance.

By the above results and Lemma 1.2, the membership functions of  $(\tilde{A} \circ \tilde{B})$  can be evaluated as:

$$(\tilde{A} \circ \tilde{B})(x) = \sup\{\alpha \in [0, 1] : x \in (\tilde{A} \circ \tilde{B})[\alpha]\}. \quad (1.17)$$

where  $\circ = \oplus, \ominus, \otimes, \oslash$  denotes the extended arithmetic operations  $\bullet = +, -, \times, /$  in a fuzzy domain.

Several definitions have been proposed to capture the information contained in a (unimodal) **FN** to simplify the **FN**'s representation using a functional parametric form known as *LR*-fuzzy numbers (*LR-FN*)  $\tilde{A} = (a; l_a, r_a)_{LR}$  [41, 114, 123]. The membership function of a *LR-FN*  $\tilde{A}$  is defined by:

$$\tilde{A}(x) = \begin{cases} L\left(\frac{a-x}{l_a}\right), & a-l_a \leq x \leq a, \\ R\left(\frac{x-a}{r_a}\right), & a \leq x \leq a+r_a, \\ 0, & x \in \mathbb{R} - [a-l_a, a+r_a], \end{cases} \quad (1.18)$$

where  $l_a > 0$  is the left spread,  $r_a > 0$  is the right spread, and  $L$  and  $R$  represent reference functions defining the left and the right shapes of the **FN**, respectively. The left and the right shapes  $L, R : [0, 1] \rightarrow [0, 1]$  should also satisfy the following conditions:

1.  $L(1) = R(1) = 0$ ,
2.  $L(0) = R(0) = 1$ , and
3.  $L(x)$  and  $R(x)$  are continuous and monotone decreasing functions on  $[0, 1]$ .

Moreover, a ***LR-FN*** becomes an ***L-FN*** when  $L(x) = R(x)$ . It is easy to check that the  $\alpha$ -cut of a ***LR-FN***  $\tilde{A}$  is:

$$\tilde{A}[\alpha] = [\tilde{A}_\alpha^L, \tilde{A}_\alpha^U] = [a - L^{-1}(\alpha)l_a, a + R^{-1}(\alpha)r_a], \quad \alpha \in [0, 1].$$

Some common operations between two ***LR-FNs*** are given as follows.

**Theorem 1.6** *For two ***LR-FNs***  $\tilde{A} = (a; l_a, r_a)_{LR}$  and  $\tilde{B} = (b; l_b, r_b)_{LR}$ ,*

- 1) *Addition:  $\tilde{A} \oplus \tilde{B} = (a + b; l_a + l_b, r_a + r_b)_{LR}$ .*
- 2) *Difference:  $\tilde{A} \ominus \tilde{B} = (a - b; l_a + r_b, r_a + l_b)_{RL}$ .*
- 3) *The scalar multiplication:*

$$\lambda \otimes \tilde{A} = \begin{cases} (\lambda a; \lambda l_a, \lambda r_a)_{LR}, & \lambda > 0, \\ (\lambda a; -\lambda r_a, -\lambda l_a)_{RL}, & \lambda < 0. \end{cases}$$

**Proof** See [41, 114, 123] to have a proof.

In next the chapters, we employ the most commonly used ***LR-FNs*** so-called triangular fuzzy numbers (***TFNs***) to handle imprecision included in data set. Note that the membership function of a ***TFN***, denoted by  $\tilde{A} = (a; l_a, r_a)_T$ , is given by:

$$\tilde{A}(x) = \begin{cases} \frac{x-(a-l_a)}{l_a}, & a - l_a \leq x \leq a, \\ \frac{a+r_a-x}{r_a}, & a \leq x \leq a + r_a, \\ 0, & x \in \mathbb{R} - [a - l_a, a + r_a]. \end{cases} \quad (1.19)$$

It is worth noting that a ***TFN***  $\tilde{A} = (a; l_a, r_a)_T$  can also be rewritten as  $\tilde{A} = (a^L, a, a^U)_T$  where  $a^L = a - l_a$  and  $a^U = a + r_a$ . In this regards, the membership function of  $\tilde{A}$  is:

$$\tilde{A}(x) = \begin{cases} \frac{x-a^L}{a-a^L}, & a^L \leq x \leq a, \\ \frac{a^U-x}{a^U-a}, & a \leq x \leq a^U, \\ 0, & x \in \mathbb{R} - [a^L, a^U]. \end{cases} \quad (1.20)$$

For simplicity, a symmetric ***TFN*** is represented by ***STFN*** throughout this book.

Here, an example is provided to obtain the function of a ***LR-FN***.

**Example 1.9** Let  $g(x) = x^2$  and  $\tilde{A} = (0; 2, 1)_{LR}$  with  $L(x) = \max\{0, 1 - x\}$  and  $R(x) = \sqrt{1 - x^2}$ . First note that the membership function of  $\tilde{A}$  is:

$$\tilde{A}(x) = \begin{cases} \frac{x+2}{2}, & -2 \leq x \leq 0, \\ \sqrt{1-x^2}, & 0 \leq x \leq 1, \\ 0, & x \in \mathbb{R} - [-2, 1]. \end{cases} \quad (1.21)$$

Then, it is easy to check that:

$$\begin{aligned} g(\tilde{A})(y) &= \sup_{x:y=x^2} \tilde{A}(y) \\ &= \begin{cases} \max\{\tilde{A}(\sqrt{y}), \tilde{A}(-\sqrt{y})\}, & 0 \leq y \leq 1, \\ \tilde{A}(-\sqrt{y}), & 1 < y \leq 4, \end{cases} \\ &= \begin{cases} \max\{\frac{2-\sqrt{y}}{2}, \sqrt{1-y}\}, & 0 \leq y \leq 1, \\ \frac{2-\sqrt{y}}{2} & 1 < y \leq 4, \end{cases} \\ &= \begin{cases} \sqrt{1-y}, & 0 \leq y \leq 0.64, \\ \frac{2-\sqrt{y}}{2}, & 0.64 < y \leq 4. \end{cases} \end{aligned}$$

The membership function of  $g(\tilde{A}) = \tilde{A}^2$  is plotted in Fig. 1.1.

An example of investigating of limit superior of a sequence of **TFNs** is given by the following example


**Example 1.10** Consider a countable sequence of **TFNs** defined as  $\tilde{A}_n = (0; l_{a_n}, r_{a_n})_T$  where  $l_{a_n} = 1/n$  and  $r_{a_n} = 2 - 1/n$ . Based on Proposition 1.3, we have

$$\begin{aligned} \limsup \tilde{A}_n[\alpha] &= \cap_{n \geq 1} \cup_{m \geq n} [-(1/m)(1 - \alpha), (2 - 1/m)(1 - \alpha)] \\ &= \cap_{n \geq 1} [-(1/n)(1 - \alpha), 2(1 - \alpha)] \\ &= [0, 2(1 - \alpha)] \\ &= \liminf \tilde{A}_n[\alpha], \end{aligned}$$

which concludes that  $\lim_{n \rightarrow \infty} \tilde{A}_n = (0; 0, 2)_T$ .

### 1.2.1 A similarity measure

Similarity measures are very important and useful tool to determine the amount of similarity between two or more fuzzy quantities. In almost every field of science and engineering, the concept of similarity measure has important significance [8]. Such measures are used to compare different kinds of objects in real-life applications [16, 20, 26, 76, 124, 171, 176, 193]. Here, a similarity measure between two **TFNs** is introduced. We will employ such criteria as a performance measure in Chapter 7.

**FIGURE 1.1**

Plot of  $g(\tilde{A})$  in Example 1.9.

**Definition 1.15** Let  $\tilde{A} = (a^L, a, a^U)_T$  and  $\tilde{B} = (b^L, b, b^U)_T$  be two **TFNs**. The similarity measure between  $\tilde{A}$  and  $\tilde{B}$  is defined as:

$$S(\tilde{A}, \tilde{B}) = 1 - \frac{|a^L - b^L| + |a - b| + |a^U - b^U|}{3(\max\{a^U, b^U\} - \min\{a^L, b^L\})}. \quad (1.22)$$

**Proposition 1.6** Let  $\tilde{A} = (a^L, a, a^U)_T$  and  $\tilde{B} = (b^L, b, b^U)_T$  be two **TFNs**. Then:

- 1)  $S(\tilde{A}, \tilde{B}) \in [0, 1]$ .
- 2)  $S(\tilde{A}, \tilde{B}) = S(\tilde{B}, \tilde{A})$ .
- 3)  $S(\tilde{A}, \tilde{B}) = 1$  if and only if  $\tilde{A} = \tilde{B}$ .

**Proof** The proof is left for the reader.

According to Proposition 1.6, we can say that the values of  $S(\tilde{A}, \tilde{B}) < 0.5$  show the degree of dissimilarity while the values of  $S(\tilde{A}, \tilde{B}) \geq 0.5$  represent the degree of similarity. In this regard, one can say that:

- 1) If  $S(\tilde{A}, \tilde{B}) \in [0.5, 0.6)$  then  $\tilde{A}$  and  $\tilde{B}$  are more or less similar.
- 2) If  $S(\tilde{A}, \tilde{B}) \in [0.6, 0.7)$  then the similarity of  $\tilde{A}$  and  $\tilde{B}$  is moderate.

- 3) If  $S(\tilde{A}, \tilde{B}) \in [0.7, 0.8)$  then the similarity of  $\tilde{A}$  and  $\tilde{B}$  is strong.
- 4) If  $S(\tilde{A}, \tilde{B}) \in [0.8, 0.9)$  then the similarity of  $\tilde{A}$  and  $\tilde{B}$  is very strong.
- 3) If  $S(\tilde{A}, \tilde{B}) \in [0.9, 0.1]$  then the similarity of  $\tilde{A}$  and  $\tilde{B}$  is complete.

**Example 1.11** Let  $\tilde{A} = (2, 3, 4)_T$  and  $\tilde{B} = (2.5, 3.5, 3.8)_T$  be two **TFNs**. Then, the similarity measure between  $\tilde{A}$  and  $\tilde{B}$  is:

$$S(\tilde{A}, \tilde{B}) = 1 - \frac{|2.5 - 2| + |3.5 - 3| + |3.8 - 4|}{3(\max\{4, 3.7\} - \min\{2, 2.5\})} = 1 - \frac{1.2}{3(4 - 2)} = 0.8.$$

Therefore, the similarity of  $\tilde{A}$  and  $\tilde{B}$  is very strong.

**Lemma 1.7** If  $\{S_i\}_{i=1}^n$  is a sequence of similarity measures then  $S = (1/n) \sum_{i=1}^n S_i$  is also a similarity measure.

**Proof** The verification of this lemma is left as an exercise.

### 1.2.2 A criteria for ranking FNs

The problem of ordering **FNs** has received considerable attention in real-life applications. For instance, in many statistical hypothesis tests with fuzzy data, fuzzy hypotheses are based on comparing observed fuzzy test statistics and critical values. Therefore there is a need for a criterion to compare such quantities. Because of the nature of the measurement, many different strategies have been proposed for ranking of **FNs** [14, 22, 27, 42, 155, 178, 192, 203, 200, 214]. Another method of ranking **FNs** relies on preference degree. This method provides a decision-maker with a preference degree to measure how a **FN** is greater than another [216, 143, 87]. Here, a common preference criterion to rank **FNs** is recalled [143].

**Definition 1.16** For two **FNs**  $\tilde{A}$  and  $\tilde{B}$ , define:

$$\Delta_{\tilde{A}\tilde{B}} = \int_{\{\alpha: \tilde{A}_\alpha^L \geq \tilde{B}_\alpha^L\}} (\tilde{A}_\alpha^L - \tilde{B}_\alpha^L) d\alpha + \int_{\{\alpha: \tilde{A}_\alpha^U \geq \tilde{B}_\alpha^U\}} (\tilde{A}_\alpha^U - \tilde{B}_\alpha^U) d\alpha. \quad (1.23)$$

The preference degree that ‘ $\tilde{A}$  is greater than  $\tilde{B}$ ’ is defined by:

$$P_d(\tilde{A} \succ \tilde{B}) = \begin{cases} 0.5, & \tilde{A} = \tilde{B}, \\ \frac{\Delta_{\tilde{A}\tilde{B}}}{\Delta_{\tilde{A}\tilde{B}} + \Delta_{\tilde{B}\tilde{A}}}, & \tilde{A} \neq \tilde{B}. \end{cases} \quad (1.24)$$

**Definition 1.17** For two **FNs**  $\tilde{A}$  and  $\tilde{B}$ , it is said that:

1.  $\tilde{A}$  is greater than  $\tilde{B}$ , denoting by  $\tilde{A} \succ_{P_d} \tilde{B}$ , if  $P_d(\tilde{A} \succ \tilde{B}) > 0.5$ .
2.  $\tilde{A}$  is equivalent to  $\tilde{B}$ , denoting by  $\tilde{A} \simeq_{P_d} \tilde{B}$ , if  $P_d(\tilde{A} \succ \tilde{B}) = P_d(\tilde{B} \succ \tilde{A}) = 0.5$ .

3.  $\tilde{A}$  is greater than or equal to  $\tilde{B}$ , denoting by  $\tilde{A} \succeq_{P_d} \tilde{B}$ , if  $\tilde{A} \succ_{P_d} \tilde{B}$  or  $\tilde{A} \simeq_{P_d} \tilde{B}$ .

The preference criterion  $P_d$  meets the following properties.

**Proposition 1.7** let  $\tilde{A}$ ,  $\tilde{B}$ , and  $\tilde{C}$  be three FNs. Then:

- 1)  $P_d$  is reciprocal, i.e.,  $P_d(\tilde{A} \succ \tilde{B}) = 1 - P_d(\tilde{B} \succ \tilde{A})$ ,
- 2)  $P_d$  is reflexive, i.e.,  $\tilde{A} \succeq_{P_d} \tilde{A}$ ,
- 3)  $P_d$  is transitive, i.e.,  $\tilde{A} \succeq_{P_d} \tilde{B}$  and  $\tilde{B} \succeq_{P_d} \tilde{C}$  imply  $\tilde{A} \succeq_{P_d} \tilde{C}$ ,
- 4)  $P_d(\tilde{A} \succeq \tilde{B}) = 1$  if and only if  $\tilde{B}_\alpha^L \leq \tilde{A}_\alpha^L$  and  $\tilde{B}_\alpha^U \leq \tilde{A}_\alpha^U$  for all  $\alpha \in [0, 1]$ .

**Proof** The parts (1), (2), and (4) are immediately followed. To prove (3), note that  $P_d(\tilde{A} \succeq \tilde{B}) \geq 0.5$  if and only if  $\int_0^1 (M_{\tilde{A}}[\alpha] - M_{\tilde{B}}[\alpha])d\alpha \geq 0$  where  $M_{\tilde{A}}[\alpha] = (\tilde{A}_\alpha^L + \tilde{A}_\alpha^U)/2$  and  $M_{\tilde{B}}[\alpha] = (\tilde{B}_\alpha^L + \tilde{B}_\alpha^U)/2$ . Therefore,  $\tilde{A} \succeq_{P_d} \tilde{B}$  and  $\tilde{B} \succeq_{P_d} \tilde{C}$  concludes that:

$$\int_0^1 (M_{\tilde{A}}[\alpha] - M_{\tilde{C}}[\alpha])d\alpha = \int_0^1 (M_{\tilde{A}}[\alpha] - M_{\tilde{B}}[\alpha])d\alpha + \int_0^1 (M_{\tilde{B}}[\alpha] - M_{\tilde{C}}[\alpha])d\alpha \geq 0.$$


That is  $\tilde{A} \succeq_{P_d} \tilde{C}$ .

**Example 1.12** Consider two TFNs  $\tilde{A} = (a^L, a, a^U)_T$  and  $\tilde{B} = (b^L, b, b^U)_T$  for three specific cases as shown in Fig 1.2. First, assume that  $b^L < a^L < a < b < b^U < a^U$  ((A) in Fig. (1.2)). To compute  $P_d(\tilde{B} \succeq \tilde{A})$ ,  $\Delta_{\tilde{A}\tilde{B}}$  and  $\Delta_{\tilde{B}\tilde{A}}$  can evaluate as follows:

$$\begin{aligned} \Delta_{\tilde{A}\tilde{B}} &= \int_0^{\frac{a^L - b^L}{b - a + a^L - b^L}} (a - b + ((b - b^L) - (a - a^L))(1 - \alpha))d\alpha \\ &\quad + \int_0^{\frac{a^U - b^U}{b - a + a^U - b^U}} ((b - a) + ((b^U - b) - (a^U - a))(1 - \alpha))(1 - \alpha)d\alpha \\ &= \frac{(a^L - b^L)^2}{2(b - a + a^L - b^L)} + \frac{(a^U - b^U)^2}{2(b - a + a^U - b^U)}, \end{aligned}$$

and

$$\begin{aligned} \Delta_{\tilde{B}\tilde{A}} &= \int_{\frac{a^L - b^L}{b - a + a^L - b^L}}^1 (b - a + ((a - a^L) - (b - b^L))(1 - \alpha)d\alpha \\ &\quad + \int_{\frac{a^U - b^U}{b - a + a^U - b^U}}^1 ((b - a) + ((b^U - b) - (a^U - a))(1 - \alpha))(1 - \alpha)d\alpha \\ &= \frac{(a - b)^2}{2(b - a + a^L - b^L)} + \frac{(a - b)^2}{2(b - a + a^U - b^U)}. \end{aligned}$$

**FIGURE 1.2**

Membership functions of  $\tilde{A}$  and  $\tilde{B}$  in three cases in Example 1.12.

Therefore:

$$P_d(\tilde{B} \succeq \tilde{A}) = \frac{\Delta_{\tilde{B}\tilde{A}}}{\Delta_{\tilde{A}\tilde{B}} + \Delta_{\tilde{B}\tilde{A}}},$$

where

$$1) \Delta_{\tilde{B}\tilde{A}} = (a - b)^2(2(b - a) + a^L - b^L + a^U - b^U).$$

$$2) \Delta_{\tilde{A}\tilde{B}} = (b - a + a^U - b^U)(a^L - b^L)^2 + (b - a + a^L - b^L)(a^U - b^U)^2.$$

Similarly, the preference degree to which  $\tilde{A}$  is greater than  $\tilde{B}$  for cases (1) and (2) can be summarized as follows:

(1) If  $b^L < a^L < a < b < a^U < b^U$  ((B) in Fig. (1.2)), then:


$$P_d(\tilde{B} \succeq \tilde{A}) = \frac{\frac{(b-a+b^L-a^L)(b-a+a^U-b^U)+(a-b)^2}{2(b-a+a^U-b^U)}}{\frac{(b-a+b^L-a^L)(b-a+a^U-b^U)+(a-b)^2}{2(b-a+a^U-b^U)} + \frac{(a^U-b^U)^2}{2(b-a+a^U-b^U)}}.$$

(2) If  $a^L < b^L < a < b < b^U < a^U$  ((C) in Fig. (1.2)), then:

$$P_d(\tilde{B} \succeq \tilde{A}) = \frac{\frac{(b-a+b^U-a^U)(b-a+a^L-b^L)+(a-b)^2}{2(b-a+a^L-b^L)}}{\frac{(b-a+b^U-a^U)(b-a+a^L-b^L)+(a-b)^2}{2(b-a+a^L-b^L)} + \frac{(a^L-b^L)^2}{2(b-a+a^L-b^L)}},$$

For all cases (1), (2), and (3), it should be noted that  $P_d(\tilde{B} \succeq \tilde{A}) \in (0.5, 1)$ .

**Remark 1.2** Using the preference degree  $P_d$  defined for each ordered pair of FNs, it should be noted that we can sort a set of  $n$  FNs  $\tilde{A}_i$  by sorting  $\{\tilde{A}_1, \tilde{A}_2, \dots, \tilde{A}_n\}$  via calculating  $\binom{n}{2}$  preference degrees into  $\{\tilde{A}_{k_1}, \tilde{A}_{k_2}, \dots, \tilde{A}_{k_n}\}$  so that for any  $i < j$ ,  $P_d(\tilde{A}_j \succeq \tilde{A}_i) \geq 0.5$ . The feasibility of the sorting is guaranteed by Proposition 1.7. Based on the sorting results, therefore,  $\tilde{A}_{k_n}$  is the most preferred choice,  $\tilde{A}_{k_{(n-1)}}$  is the second, etc.

**FIGURE 1.3**

Membership functions of  $\tilde{A}_i$ ,  $i = 1, 2, \dots, 5$  in Example 1.13.

**Example 1.13** Consider five **TFNs** of  $\tilde{A}_1 = (-1, 0, 1)_T$ ,  $\tilde{A}_2 = (0.5, 1, 1.5)_T$ ,  $\tilde{A}_3 = (1, 2, 2.5)_T$ ,  $\tilde{A}_4 = (-0.5, 0.5, 2.5)_T$ , and  $\tilde{A}_5 = (0, 1.5, 3)_T$ . The plots of  $\tilde{A}_i$ ,  $i = 1, 2, \dots, 5$  are shown in Fig. 1.3. The preference degree  $P_d(\tilde{A}_j \succeq \tilde{A}_i)$ ,  $i, j = 1, 2, \dots, 5$  are listed in Table 1.1. Accordingly, we can get a sorted result as  $\tilde{A}_5 \succeq_{P_d} \tilde{A}_3 \succeq_{P_d} \tilde{A}_4 \succeq_{P_d} \tilde{A}_2 \succeq_{P_d} \tilde{A}_1$ .

**TABLE 1.1**

Preference degrees  $P_d(\tilde{A}_i \succeq \tilde{A}_j)$  in Example 1.13.

$j$	1	2	3	4	5
$P_d(\tilde{A}_1 \succeq \tilde{A}_j)$	1	0	0	0	0
$P_d(\tilde{A}_2 \succeq \tilde{A}_j)$	1	1	0	$\frac{3}{8}$	$\frac{1}{2}$
$P_d(\tilde{A}_3 \succeq \tilde{A}_j)$	1	1	1	1	$\frac{3}{8}$
$P_d(\tilde{A}_4 \succeq \tilde{A}_j)$	1	$\frac{5}{8}$	0	1	0
$P_d(\tilde{A}_5 \succeq \tilde{A}_j)$	1	$\frac{1}{2}$	$\frac{5}{8}$	1	1

**Remark 1.3** Let  $\tilde{A} = (a^L, a, a^U)_T$  and  $\tilde{B} = (b^L, b, b^U)_T$  be two **TFNs**. According to Proposition 1.7, we have  $P_d(\tilde{A} \succeq \tilde{B}) = 1$  if and only if  $a^L \leq b^L$ ,  $a \leq b$  and  $a^U \leq b^U$ .

Further, a different type of ranking for **TFNs** is introduced based on an exact criterion. We employed this criterion to construct a fuzzy hypothesis in Chapter 5.

**Definition 1.18** Let  $\tilde{A} = (a^L, a, a^U)_T$  and  $\tilde{B} = (b^L, b, b^U)_T$  be two **TFNs**. Then, we say that:

1.  $\tilde{A} \simeq_M \tilde{B}$ , if  $M_{\tilde{A}} = M_{\tilde{B}}$ .
2.  $\tilde{A} \succ_M \tilde{B}$ , if  $M_{\tilde{A}} > M_{\tilde{B}}$ .
3.  $\tilde{A} \succeq_M \tilde{B}$ , if  $M_{\tilde{A}} > M_{\tilde{B}}$  or  $M_{\tilde{A}} = M_{\tilde{B}}$ .  
where  $M_{\tilde{A}} = (a^L + a + a^U)/3$  and  $M_{\tilde{B}} = (b^L + b + b^U)/3$ .

**Lemma 1.8** Let  $\tilde{A}$ ,  $\tilde{B}$ , and  $\tilde{C}$  be three **TFNs**. Then:

- 1)  $\succeq_M$  is reflexive, i.e.,  $\tilde{A} \succeq_M \tilde{A}$ .
- 2)  $\succeq_M$  is transitive, i.e.,  $\tilde{A} \succeq_M \tilde{B}$  and  $\tilde{B} \succeq_M \tilde{C}$  imply  $\tilde{A} \succeq_M \tilde{C}$ .
- 3)  $\succeq_M$  is antisymmetric, i.e.,  $\tilde{A} \succeq_M \tilde{B}$  and  $\tilde{B} \succeq_M \tilde{A}$  imply  $\tilde{A} \simeq_M \tilde{B}$ .

**Proof** The proof is left as an exercise.

### 1.2.3 Distance measure

In several fields, the necessity to determine the distance that separates two points arises. When there is imprecision on the location of these points, the calculation of the distance has to take this uncertainty into consideration. The methods for measuring the distance between fuzzy sets have become important due to their significant applications in various areas [17, 49, 50, 79, 136, 192, 199]. Particularly, many statistical inferences such as hypothesis test, decision making, regression analysis, and non-parametric inferences rely on distance measures. Therefore, when we speak statistical inferences based on fuzzy data, there is a need to extend the classic distance measures where the observed data are reported by fuzzy quantities.

**Definition 1.19** It is said that  $d$  is a distance measure between two fuzzy sets if it meets the following conditions for any fuzzy sets  $\tilde{A}$ ,  $\tilde{B}$ , and  $\tilde{C}$ :

- 1)  $d(\tilde{A}, \tilde{B}) = 0$  if and only if  $\tilde{A} = \tilde{B}$ ,
- 2)  $d(\tilde{A}, \tilde{B}) = d(\tilde{B}, \tilde{A})$ ,
- 3)  $d(\tilde{A}, \tilde{C}) \leq d(\tilde{A}, \tilde{B}) + d(\tilde{B}, \tilde{C})$ .

Here, a non-fuzzy distance measures between two **TFNs** are introduced and discussed.

**Lemma 1.9** Consider two **TFNs**  $\tilde{A} = (a^L, a, a^U)_T$  and  $\tilde{B} = (b^L, b, b^U)_T$ . Let:

- 1)  $d_1(\tilde{A}, \tilde{B}) = \frac{|a^L - b^L| + |a - b| + |a^U - b^U|}{3}$ ,
- 2)  $d_2(\tilde{A}, \tilde{B}) = \max\{|a^L - b^L|, |a - b|, |a^U - b^U|\}$ ,

$$3) d_3(\tilde{A}, \tilde{B}) = \sqrt{|a^L - b^L|^2 + |a - b|^2 + |a^U - b^U|^2}.$$

Then,  $d_1$ ,  $d_2$ , and  $d_3$  are distance measures.

**Proof** The proof is left for the reader.

**Example 1.14** Consider two **TFNs**  $\tilde{A} = (-3, -1, 3)_T$  and  $\tilde{B} = (-2, 1, 4)_T$ . Then:

$$1) d_1(\tilde{A}, \tilde{B}) = \frac{|a^L - b^L| + |a - b| + |a^U - b^U|}{3} = 4/3,$$

$$2) d_2(\tilde{A}, \tilde{B}) = \sqrt{\frac{|a^L - b^L|^2 + |a - b|^2 + |a^U - b^U|^2}{3}} = \sqrt{2},$$

$$3) d_3(\tilde{A}, \tilde{B}) = \max\{|a^L - b^L|, |a - b|, |a^U - b^U|\} = 2.$$

The following definition extends the distance between two **FNs** of  $\tilde{A}$  and  $\tilde{B}$  as a **FN**.

**Definition 1.20** [199] For three fuzzy sets  $\tilde{A}$ ,  $\tilde{B}$ , and  $\tilde{C}$ , it is said that  $\tilde{D}$  is a fuzzy distance measure (**FDM**) if it meets the following conditions:

$$1) \tilde{D}(\tilde{A}, \tilde{B}) = I\{0\} \text{ if and only if } \tilde{A} = \tilde{B}.$$

$$2) \tilde{D}(\tilde{A}, \tilde{B}) = \tilde{D}(\tilde{B}, \tilde{A}).$$

$$3) \tilde{D}(\tilde{A}, \tilde{C}) \preceq_{P_d} \tilde{D}(\tilde{A}, \tilde{B}) \oplus \tilde{D}(\tilde{B}, \tilde{C}).$$

**Theorem 1.7** For two fuzzy sets  $\tilde{A}$  and  $\tilde{B}$  define:

$$\tilde{D}(\tilde{A}, \tilde{B})(d) = \sup_{x,y:|x-y|=d} \min\{\tilde{A}(x), \tilde{B}(y)\}. \quad (1.25)$$

Then,  $\tilde{D}$  is a **FDM**.

**Proof** To prove, see [199].

**Example 1.15** Let  $\mathbb{X} = \{-1, 0, 2, 3, 4\}$ ,  $\tilde{A} = \{\frac{0.3}{-1}, \frac{0.5}{0}, \frac{0.7}{2}, \frac{1}{3}, \frac{0.8}{4}\}$ , and  $\tilde{B} = \{\frac{0.8}{-1}, \frac{1}{0}, \frac{0.9}{2}, \frac{0.6}{3}, \frac{0.4}{4}\}$ . Then, the membership functions of  $\tilde{D}(\tilde{A}, \tilde{B})$  can be evaluated as follows:

$$\begin{aligned} \tilde{D}(\tilde{A}, \tilde{B})(0) &= \sup_{x,y:|x-y|=0} \min\{\tilde{A}(x), \tilde{B}(y)\} \\ &= \max \left\{ \min\{\tilde{A}(-1), \tilde{B}(-1)\}, \min\{\tilde{A}(0), \tilde{B}(0)\} \right. \\ &\quad , \min\{\tilde{A}(2), \tilde{B}(2)\}, \min\{\tilde{A}(3), \tilde{B}(3)\}, \min\{\tilde{A}(4), \tilde{B}(4)\} \left. \right\} \\ &= \max \left\{ \min\{0.3, 0.8\}, \min\{0.5, 1\}, \min\{0.7, 0.9\}, \min\{1, 0.6\}, \right. \\ &\quad \left. \min\{0.4, 0.8\} \right\} = 0.7. \end{aligned}$$

$$\begin{aligned}
\tilde{D}(\tilde{A}, \tilde{B})(1) &= \sup_{x,y:|x-y|=1} \min\{\tilde{A}(x), \tilde{B}(y)\} \\
&= \max \left\{ \min\{\tilde{A}(-1), \tilde{B}(0)\}, \min\{\tilde{A}(0), \tilde{B}(-1)\}, \min\{\tilde{A}(2), \right. \\
&\quad \left. \tilde{B}(3)\}, \min\{\tilde{A}(3), \tilde{B}(2)\}, \min\{\tilde{A}(4), \tilde{B}(3)\}, \min\{\tilde{A}(3), \tilde{B}(4)\} \right\} \\
&= \max \left\{ \min\{0.3, 1\}, \min\{0.8, 0.3\}, \min\{0.7, 0.6\}, \min\{0.6, 0.7\}, \right. \\
&\quad \left. \min\{1, 0.9\}, \min\{0.8, 0.6\}, \min\{0.4, 1\} \right\} = 0.9.
\end{aligned}$$

Similarly, it can be shown that  $\tilde{D}(\tilde{A}, \tilde{B})(3) = 1$ ,  $\tilde{D}(\tilde{A}, \tilde{B})(2) = \tilde{D}(\tilde{A}, \tilde{B})(4) = \tilde{D}(\tilde{A}, \tilde{B})(5) = 0.8$  and thus:

$$\tilde{D}(\tilde{A}, \tilde{B}) = \left\{ \frac{0.7}{0}, \frac{0.9}{1}, \frac{0.8}{2}, \frac{1}{3}, \frac{0.8}{\{4, 5\}} \right\}.$$

Here, the proposed fuzzy distance measures between two **TFNs** are developed in the fuzzy domain.

**Proposition 1.8** Consider two **TFNs**  $\tilde{A} = (a^L, a, a^U)_T$  and  $\tilde{B} = (b^L, b, b^U)_T$ . Let:

$$\tilde{d}_a(\tilde{A}, \tilde{B}) = (d_1(\tilde{A}, \tilde{B}), d_2(\tilde{A}, \tilde{B}), d_3(\tilde{A}, \tilde{B}))_T, \quad (1.26)$$

where  $d_1$ ,  $d_2$ , and  $d_3$  are as in Lemma 1.9. Then,  $\tilde{d}_a$  is **FDM**.

**Proof** First note that  $d_1$ ,  $d_2$ , and  $d_3$  are (exact) distance measures and  $d_1(\tilde{A}, \tilde{B}) \leq d_2(\tilde{A}, \tilde{B}) \leq d_3(\tilde{A}, \tilde{B})$  for any  $\tilde{A}, \tilde{B} \in \mathcal{F}(\mathbb{R})$ . Therefore, it can be checked that

$$1) \quad \tilde{d}_a(\tilde{A}, \tilde{B}) = I\{0\} \text{ if and only if } \tilde{A} = \tilde{B},$$

$$2) \quad \tilde{d}_a(\tilde{A}, \tilde{B}) = \tilde{d}_a(\tilde{B}, \tilde{A}),$$

and

$$\begin{aligned}
d_1^L(\tilde{A}, \tilde{C}) &\leq d_1^L(\tilde{A}, \tilde{B}) + d_1^L(\tilde{B}, \tilde{C}), \\
d_2(\tilde{A}, \tilde{C}) &\leq d_2(\tilde{A}, \tilde{B}) + d_2(\tilde{B}, \tilde{C}), \\
d_3^U(\tilde{A}, \tilde{C}) &\leq d_3^U(\tilde{A}, \tilde{B}) + d_3^U(\tilde{B}, \tilde{C}).
\end{aligned}$$

From Remark 1.3, these conclude that  $P_d(\tilde{d}_a(\tilde{A}, \tilde{C})) \preceq (\tilde{d}_a(\tilde{A}, \tilde{B}) \oplus \tilde{d}_a(\tilde{B}, \tilde{C})) = 1$  or  $\tilde{d}_a(\tilde{A}, \tilde{C}) \preceq_{P_d} \tilde{d}_a(\tilde{A}, \tilde{B}) \oplus \tilde{d}_a(\tilde{B}, \tilde{C})$ . Therefore,  $\tilde{d}_a$  is **FDM**.

**Example 1.16** Recall Example 1.14. Therefore, the fuzzy distance between  $\tilde{A}$  and  $\tilde{B}$  can be evaluated via a **TFN**  $\tilde{d}_a(\tilde{A}, \tilde{B}) = (4/3, \sqrt{2}, 2)_T$ .

### 1.3 $\alpha$ -values of FNs

Here, the notion of  $\alpha$ -values of FNs is recalled. We will utilize such a notion to rewrite a common notion of the fuzzy random variable in the next section [78].

**Definition 1.21** *The  $\alpha$ -values of a FN  $\tilde{A}$  is a mapping  $\tilde{A}_\alpha : [0, 1] \rightarrow \mathbb{R}$  defined by:*

$$\tilde{A}_\alpha = \begin{cases} \tilde{A}_{2\alpha}^L, & \alpha \in [0, 0.5], \\ \tilde{A}_{2(1-\alpha)}^U, & \alpha \in [0.5, 1]. \end{cases} \quad (1.27)$$

**Lemma 1.10** *The  $\alpha$ -values a FN  $\tilde{A}$  is an increasing function of  $\alpha$  on  $[0, 1]$ .*

**Proof** Since  $\tilde{A}_\alpha^L$  and  $\tilde{A}_\alpha^U$  are increasing and decreasing functions, for any  $\alpha_1 < \alpha_2$  in  $[0, 1]$ , we have  $\tilde{A}_{2\alpha_1}^L \leq \tilde{A}_{2\alpha_2}^L$  for  $\alpha \in [0, 0.5]$  and  $\tilde{A}_{2(1-\alpha_1)}^U \leq \tilde{A}_{2(1-\alpha_2)}^U$  for  $\alpha \in [0.5, 1]$ . These verify the proof.

**Example 1.17** Let  $\tilde{A} = (a; l_a, r_a)_{LR}$  be a LR-FN. From Definition 1.21, one finds that:

$$\tilde{A}_\alpha = \begin{cases} a - l_a L^{-1}(2\alpha), & 0 \leq \alpha \leq 0.5, \\ a + r_a R^{-1}(2(1 - \alpha)), & 0.5 \leq \alpha \leq 1. \end{cases}$$

For instance,

1. If  $\tilde{A} = (a; l_a, r_a)_T$  is a TFN, then:

$$\tilde{A}_\alpha = \begin{cases} (a - l_a) + 2l_a\alpha, & 0 \leq \alpha \leq 0.5, \\ a + r_a - 2r_a(1 - \alpha), & 0.5 \leq \alpha \leq 1. \end{cases}$$

2. Let  $\tilde{A} = (a; l_a, r_a)_{LR}$  with  $L(x) = \sqrt{1 - x^3}$  and  $R(x) = 1 - x^5$  then:

$$\tilde{A}_\alpha = \begin{cases} a - l_a \sqrt[3]{1 - 4\alpha^2}, & 0 \leq \alpha \leq 0.5, \\ a + r_a \sqrt[5]{2\alpha - 1}, & 0.5 \leq \alpha \leq 1. \end{cases}$$

The relationship between  $\alpha$ -values and  $\alpha$ -cuts of a FN is given below.

**Lemma 1.11** *Let  $\tilde{A}[\alpha]$  and  $\tilde{A}_\alpha$  be the  $\alpha$ -cuts and  $\alpha$ -values of a FN  $\tilde{A}$ , respectively. Then:*

$$\tilde{A}[\alpha] = [\tilde{A}_\alpha^L, \tilde{A}_\alpha^U] = [\tilde{A}_{\alpha/2}, \tilde{A}_{1-\alpha/2}]. \quad (1.28)$$

**Proof** The proof is a simple consequence of Eq. (1.27) since  $\tilde{A}_{\alpha/2} = \tilde{A}_\alpha^L$  and  $\tilde{A}_{1-\alpha/2} = \tilde{A}_\alpha^U$  for any  $\alpha \in [0, 1]$ .

Therefore, the membership degree of  $\tilde{A}$  at  $x \in \mathbb{R}$  can be written as:

$$\tilde{A}(x) = \sup\{\alpha \in [0, 1] : x \in [\tilde{A}_{\alpha/2}, \tilde{A}_{1-\alpha/2}]\}.$$

The following Lemma shows that the interval-valued arithmetic operations of **FNs** can be simplified using their  $\alpha$ -values.

**Lemma 1.12** Let  $\lambda \in \mathbb{R}$  and  $\tilde{A}$  and  $\tilde{B}$  be two **FNs**. Then, for any  $\alpha \in [0, 1]$ :

$$1. (\tilde{A} \oplus \tilde{B})_\alpha = \tilde{A}_\alpha + \tilde{B}_\alpha.$$

$$2. (\lambda \otimes \tilde{A})_\alpha = \begin{cases} \lambda \tilde{A}_\alpha, & \lambda \geq 0, \\ -\lambda \tilde{A}_{1-\alpha}, & \lambda < 0. \end{cases}$$

$$3. (\tilde{A} \otimes \tilde{B})_\alpha = \begin{cases} \tilde{A}_\alpha \tilde{B}_\alpha, & \text{if } \tilde{A}, \tilde{B} > 0, \\ \tilde{A}_{1-\alpha} \tilde{B}_{1-\alpha}, & \text{if } \tilde{A}, \tilde{B} < 0, \\ \tilde{A}_{1-\alpha} \tilde{B}_\alpha, & \text{if } \tilde{A} > 0, \tilde{B} < 0. \end{cases}$$

**Proof** From Definition 1.21 and the arithmetic operation of **FNs**, it is easy to check that:

$$\begin{aligned} (\tilde{A} \oplus \tilde{B})_\alpha &= \begin{cases} (\tilde{A} \oplus \tilde{B})_{2\alpha}^L, & 0 \leq \alpha \leq 0.5, \\ (\tilde{A} \oplus \tilde{B})_{2(1-\alpha)}^U, & 0.5 < \alpha \leq 1. \end{cases} \\ &= \begin{cases} \tilde{A}_{2\alpha}^L + \tilde{B}_{2\alpha}^L, & 0 \leq \alpha \leq 0.5, \\ \tilde{A}_{2(1-\alpha)}^U + \tilde{B}_{2(1-\alpha)}^U, & 0.5 < \alpha \leq 1. \end{cases} \\ &= \tilde{A}_\alpha + \tilde{B}_\alpha, \end{aligned} \tag{1.29}$$

and

$$\begin{aligned} (\lambda \otimes \tilde{A})_\alpha &= \begin{cases} (\lambda \otimes \tilde{A})_{2\alpha}^L, & 0 \leq \alpha \leq 0.5, \\ (\lambda \otimes \tilde{A})_{2(1-\alpha)}^U, & 0.5 < \alpha \leq 1. \end{cases} \\ &= \lambda \tilde{A}_\alpha I(\lambda \geq 0) - \lambda \tilde{A}_{1-\alpha} I(\lambda < 0). \end{aligned} \tag{1.30}$$

These verify (1) and (2). To prove (3), first assume that  $\tilde{A}$  and  $\tilde{B}$  are two positive **FNs**. Since  $(\tilde{A}^2)_\alpha = (\tilde{A}_\alpha)^2$  we get:

$$\begin{aligned} ((\tilde{A} \oplus \tilde{B})^2)_\alpha &= ((\tilde{A} \oplus \tilde{B}) \otimes (\tilde{A} \oplus \tilde{B}))_\alpha \\ &= (\tilde{A}_\alpha + \tilde{B}_\alpha)^2 \\ &= \tilde{A}_\alpha^2 + \tilde{B}_\alpha^2 + 2\tilde{A}_\alpha \tilde{B}_\alpha. \end{aligned}$$

By (1) and (2), therefore:

$$\begin{aligned} ((\tilde{A} \oplus \tilde{B})^2)_\alpha &= (\tilde{A}^2 \oplus \tilde{B}^2 \oplus 2\tilde{A} \otimes \tilde{B})_\alpha \\ &= \tilde{A}_\alpha^2 + \tilde{B}_\alpha^2 + 2(\tilde{A} \otimes \tilde{B})_\alpha. \end{aligned}$$

This shows that  $(\tilde{A} \otimes \tilde{B})_\alpha = \tilde{A}_\alpha \tilde{B}_\alpha$ . Now, if  $\tilde{A}$  and  $\tilde{B}$  are two negative FNs, then we have  $\tilde{A} = (-1) \otimes \tilde{A}'$  and  $\tilde{B} = (-1) \otimes \tilde{B}'$  for two positive FNs of  $\tilde{A}'$  and  $\tilde{B}'$ . So, it concludes that:

$$\begin{aligned} (\tilde{A} \otimes \tilde{B})_\alpha &= \left( (-1) \otimes \tilde{A}' \otimes (-1) \otimes \tilde{B}' \right)_\alpha \\ &= (\tilde{A}' \otimes \tilde{B}')_\alpha = \tilde{A}'_\alpha \tilde{B}'_\alpha = \tilde{A}_{1-\alpha} \tilde{B}_{1-\alpha}. \end{aligned}$$

This provides (3). These results can also be verified in cases where  $\tilde{A} > 0$  and  $\tilde{B} < 0$ .

According to Lemma 1.12, it should be noted that the arithmetic operations of FNs can be rewritten according to their  $\alpha$ -values.

**Lemma 1.13** Consider a mapping  $\tilde{A}_\alpha : [0, 1] \rightarrow \mathbb{R}$  such that: 1)  $\tilde{A}_\alpha$  strictly increasing with respect to  $\alpha$  for any  $\alpha \in [0, 1]$  and 2)  $\tilde{A}_{0.5}$  is a constant number. Then,  $\{\tilde{A}_\alpha\}_{\alpha \in [0, 1]}$  provides a FN  $\tilde{A}$  with the following  $\alpha$ -cut:

$$\tilde{A}[\alpha] = [\tilde{A}_{\alpha/2}, \tilde{A}_{1-\alpha/2}]. \quad (1.31)$$

**Proof** Let  $\tilde{A}_\alpha : [0, 1] \rightarrow \mathbb{R}$  be a mapping such that: 1)  $\tilde{A}_\alpha$  strictly increasing with respect to  $\alpha$  for any  $\alpha \in [0, 1]$  and 2)  $\tilde{A}_{0.5}$  is a constant number. It is easy to verify that  $[\tilde{A}_{1-\alpha_2/2}, \tilde{A}_{\alpha_2/2}] \subseteq [\tilde{A}_{1-\alpha_1/2}, \tilde{A}_{\alpha_1/2}]$  for any  $\alpha_1 < \alpha_2$ . Therefore,  $\tilde{A}[\alpha] = [\tilde{A}_{1-\alpha/2}, \tilde{A}_{\alpha/2}]$  construct a sequence of  $\alpha$ -cuts of a FN.


This is the reason why we focused on  $\alpha$ -values since a FN of  $\tilde{A}$  can be traceable via its  $\alpha$ -values ( $\{\tilde{A}_\alpha\}$ ) instead of its  $\alpha$ -cuts ( $\{\tilde{A}[\alpha]\}$ ).

**Lemma 1.14** Let  $\tilde{A} \in \mathcal{F}(\mathbb{R})$  and  $g$  be a continuous and strictly monotone real-valued function. Then, for every  $\alpha \in [0, 1]$

$$(g(\tilde{A}))_\alpha = \begin{cases} g(\tilde{A}_\alpha), & g \text{ is increasing,} \\ g(\tilde{A}_{1-\alpha}), & g \text{ is decreasing.} \end{cases} \quad (1.32)$$

**Proof** Without loss of generality, assume that  $g$  is a strictly increasing function. Regarding the extension principle and Definition 1.21, first note that  $g(\tilde{A})(y) = \tilde{A}(g^{-1}(y))$  and therefore:

$$\begin{aligned} g(\tilde{A})[\alpha] &= g(\tilde{A}[\alpha]) \\ &= [g(\tilde{A}_\alpha^L), g(\tilde{A}_\alpha^U)] \\ &= [g(\tilde{A}_{\alpha/2}), g(\tilde{A}_{1-\alpha/2})]. \end{aligned}$$

**FIGURE 1.4**

Membership function of  $\exp(\tilde{A})$  in Example 1.18.

Now, by the relationship between  $\alpha$ -values and  $\alpha$ -cuts given in Definition 1.21, it can be seen that  $(g(\tilde{A}))_\alpha = g(\tilde{A}_\alpha)$  for every  $\alpha \in [0, 1]$ .

**Remark 1.4** By the relationship between  $\alpha$ -values and  $\alpha$ -cuts of **FNs**, note that the preference degree that ' $\tilde{A}$  is greater than  $\tilde{B}$ ' can be rewritten as follows:

$$P_d(\tilde{A} \succ \tilde{B}) = \begin{cases} 0.5, & \tilde{A} = \tilde{B}, \\ \frac{\Delta_{\tilde{A}\tilde{B}}}{\Delta_{\tilde{A}\tilde{B}} + \Delta_{\tilde{B}\tilde{A}}}, & \tilde{A} \neq \tilde{B}. \end{cases}$$

where

$$\Delta_{\tilde{A}\tilde{B}} = \int_{\{\alpha: \tilde{A}_\alpha \geq \tilde{B}_\alpha\}} (\tilde{A}_\alpha - \tilde{B}_\alpha) d\alpha. \quad (1.33)$$

Therefore  $P_d(\tilde{A} \succeq \tilde{B}) = 1$  if and only if  $\tilde{A}_\alpha \geq \tilde{B}_\alpha$  for any  $\alpha \in [0, 1]$ .

From Lemma 1.14, we can observe that the conventional function of a **FN** (based on the extension principle) can also be rewritten based on the  $\alpha$ -values of the **FNs**.

**Example 1.18** Let  $g(x) = \exp(x)$  and  $\tilde{A} = (a; l_a, r_a)_T$  be a **TFN**. Therefore, the  $\alpha$ -cuts of  $g(\tilde{A})$  can be evaluated as:

$$(g(\tilde{A}))_\alpha = \begin{cases} \exp((a - l_a) + 2l_a\alpha), & 0 \leq \alpha \leq 0.5, \\ \exp(a + r_a - 2r_a(1 - \alpha)), & 0.5 < \alpha \leq 1. \end{cases}$$

## References

- S.S.G. Abhishek and S.R. Singh . A score function-based method of forecasting using intuitionistic fuzzy time series. *New Mathematics and Natural Computation*, 14:91–111, 2018.
- M.G. Akbari and G. Hesamian . A partial-robust-ridge-based regression model with fuzzy predictors-responses. *Journal of Computational and Applied Mathematics*, 351:290–301, 2019.
- M.G. Akbari and G. Hesamian . Elastic net oriented to fuzzy semi-parametric regression model with fuzzy explanatory variables and fuzzy responses. 1:1–11, 2019.
- Abbasi G. Al-Refaie, A. and Ghanim D. . Proposed  $\alpha$ -cut CUSUM and EWMA Control Charts for Fuzzy Response Observations. *International Journal of Reliability, Quality and Safety Engineering*, 28:21–28, 2021.
- I.M. Aliev and Z. Kara . Fuzzy system reliability analysis using time dependent fuzzy set. *Control and Cybernetics*, 33:653–662, 2004.
- B.F. Arnold . Testing fuzzy hypothesis with crisp data. *Fuzzy Sets and Systems*, 9:323–333, 1998.
- Z. Artstein and J.C. Hansen . Convexification in limit laws of random sets in Banach spaces. *Annals of Probability*, 13:307–309, 1985.
- S Ashraf and T Rashid . Fuzzy similarity measures. LAP LAMBERT Academic Publishing, 2010.
- Bantan R.A.R. Aslam, M. and Khan N. . Design of S 2-NNEWMA control chart for monitoring process having indeterminate production data. *Processes*, 7:36–44, 2019.
- Lawry J. Baldwin, J.F. and Martin T.P. . A mass assignment theory of the probability of fuzzy events. *Fuzzy Sets and Systems*, 83:353–367, 1996.
- Pedrycz W. Bargiela, A. and Nakashima T. . Multiple regression with fuzzy data. *Fuzzy sets and systems*, 158:2169–2188, 2007.
- B. Bede . Mathematics of Fuzzy Sets and Fuzzy Logic. Springer Heidelberg, Berlin, 2013.
- P Billingsley . Probability and measure. John Wiley and Sons, New York, 2008.
- G. Bortolan and R.A. Degani . A review of some methods for ranking fuzzy subsets. *Fuzzy Sets and Systems*, 15:1–19, 1985.
- G.E.P. Box and G.M. Jenkins . Time series analysis: forecasting and control. Holden-Day, San Francisco, 1976.
- Barrenechea E. Bustince, H. and Pagola M. . Relationship between restricted dissimilarity functions, restricted equivalence functions and normal EN-functions: Image thresholding invariant. *Pattern Recognition Letters*, 29:525–536, 2008.
- C. Chakraborty and D. Chakraborty . A theoretical development on a fuzzy distance measure for fuzzy numbers. *Mathematical and Computer Modeling*, 43:254–261, 2006.
- K.S. Chen and T.C. Chang . Construction and fuzzy hypothesis testing of Taguchi Six Sigma quality index. *International Journal of Production Research*, 58:3110–3125, 2020.
- L.H. Chen and S.H. Nien . A new approach to formulate fuzzy regression models. *Applied Soft Computing*, 86:23–36, 2020.
- S.J. Chen and S.M. Chen . Fuzzy risk analysis based on similarity measures of generalized fuzzy numbers. *IEEE Transactions on Fuzzy Systems*, 11:45–56, 2003.
- C.B. Cheng and E.S. Lee . Non-parametric fuzzy regression k-NN and kernel smoothing techniques. *Computers and Mathematics with Applications*, 38:239–251, 1999.
- C.H. Cheng . A new approach for ranking fuzzy numbers by distance method. *Fuzzy Sets and Systems*, 95:307–317, 1998.
- C.H. Cheng and C.H. Chen . Fuzzy time series model based on weighted association rule for financial market forecasting. *Expert Systems*, 35:23–30, 2018.
- S.H. Choi and J.J. Buckley . Fuzzy regression using least absolute deviation estimators. *Soft Computing*, 12:257–263, 2008.
- S.H. Choi and J.H. Yoon . General fuzzy regression using least squares method. *International Journal of Systems Science*, 41:477–485, 2010.
- C.c. Chou . A new similarity measure of fuzzy numbers. *Journal of Intelligent and Fuzzy Systems*, 26:287–294, 2014.
- T.C. Chu and C.T. Tsao . Ranking fuzzy numbers with an area between the centroid point and original point. *Computers and Mathematic Application*, 43:111–117, 2002.
- N. Chukhrova and A. Johannsen . Fuzzy regression analysis: Systemtic review and bibliography. *Applied Soft Computing*, 84:1–20, 2019.

- N. Chukhrova and A. Johannsen . Fuzzy hypothesis testing for a population proportion based on set-valued information. *Fuzzy Sets and Systems*, 387:127–157, 2020.
- N. Chukhrova and A. Johannsen . Fuzzy hypothesis testing: Systematic review and bibliography. *Applied Soft Computing*, page <https://doi.org/10.1016/j.asoc.2021.107331>, 2021.
- KL Chung . *A Course in Probability Theory ( 3rd Ed.)*. Academic Press, San Diego, 2001.
- Dominguez-Mencher J.S. Lopez-Diaz M. Colubi, A. and Ralescu D.A. . On the formalization of fuzzy random variables. *Information Science*, 133:3–6, 2001.
- D'Urso- P. Giordani P. Coppi, R. and A. Santoro . Least squares estimation of a linear regression model with *L<sub>R</sub>*-fuzzy response. *Computational Statistics and Data Analysis*, 51:267–286, 2006.
- Bustince- H. Fernndez J. Couso, I. and L. Snchez . The Null Space of Fuzzy Inclusion Measures. *IEEE Transactions on Fuzzy Systems*, 29:641–648, 2021.
- W.H. Woodall Champ- C.W. Cynthia , A. Lowry and E.R. Steven . A multivariate exponentially weighted moving average control chart. *Technometrics*, 34:46–53, 1992.
- J. de Andrés-Sánchez . Claim reserving with fuzzy regression and the two ways of ANOVA. *Applied Soft Computing*, 12, 2012.
- G Debara . *Measure theory and integration*. Ellis Horwood Limited, 1981.
- V.G. Deganibay . Behaviour of fuzzy ANOVA. *Kybernetes*, 16:107–112, 1987.
- Masson M.H. Denoeux, T. and Herbert P.H. . Non-parametric rank-based statistics and significance tests for fuzzy data. *Fuzzy Sets and Systems*, 153:1–28, 2005.
- P. Diamond and P. Kloeden . *Metric Spaces of Fuzzy Sets*. World Scientific, Singapore, 1994.
- D. Dubois and H. Prade . *Fuzzy Sets and Systems: Theory and Applications*. Academic, New York, 1980.
- D. Dubois and H. Prade . Ranking fuzzy numbers in the setting of possibility theory. *Information Sciences*, 30:183–224, 1983.
- D Dubois and H Prade . *Possibility Theory*. New York: Plenum, 1988.
- N Erginol and S Senturk . *Fuzzy EWMA and fuzzy CUSUM control charts* . In: Kahraman C. , Kabak O. (eds) *Fuzzy Statistical Decision-Making. Studies in Fuzziness and Soft Computing* , vol. 343 ., pp. 281 – 295. Springer, Cham, 2016.
- A. Faraz and A.F. Shapiro . An application of fuzzy random variables to control charts. *Fuzzy Sets and Systems*, 161:2684–2694, 2010.
- TS Ferguson . *A Course in Large Sample Theory*. London: Chapman & Hall, 1996.
- T. Flaminio and L. Godo . A logic for reasoning about the probability of fuzzy events. *Fuzzy Sets and Systems*, 158:625–638, 2007.
- S. Frühwirth-Schnatter . On statistical inference for fuzzy data with applications to descriptive statistics. *Fuzzy Sets and Systems*, 50:143–165, 1992.
- M. Ganbari and R. Nuraei . Revision of a fuzzy distance measure. *International Journal of Industrial Mathematics*, 5:143–147, 2013.
- A.H. Ganie and S. Singh . An innovative picture fuzzy distance measure and novel multi-attribute decision-making method. *Complex and Intelligent Systems*, 7:781–805, 2021.
- G.D. Garson . *Curve fitting and non-linear regression*. Asheboro, NC: Statistical Associates Publishers, 2012.
- S.S. Gautam and S. Singh . A refined method of forecasting based on high-order intuitionistic fuzzy time series data. *Progress in Artificial Intelligence*, 7:339–350, 2018.
- JD Gibbons and S Chakraborti . *Non-parametric Statistical Inference ( 4th ed.)*. Marcel Dekker, New York, 2003.
- JD Gibbons and S Chakraborti . *Non-parametric Statistical Inference ( 5th ed.)*. Marcel Dekker, New York, 2003.
- Lpez-Daz M. Gil, M.A. and Ralescu D.A. . Overview on the development of fuzzy random variables. *Fuzzy Sets and Systems*, 157:2546–2557, 2006.
- Montenegro- M. Rodríguez G. -Colubi A. Gil, M.A. and Casals M.R. . Bootstrap approach to the multi-sample test of means with imprecise data. *Computer Statistics and Data Analysis*, 51:148–162, 2006.
- Hahn- M.G. Gine, E. and J. Zinn . Limit theorems for random sets, an application of probability in Banach space results, *Probab. ACM International Conference Proceeding Series* , pages 112–135, 1983.
- Belyayev- YK Gnedenko, BV and AD Solov'yev . *Mathematical methods of reliability theory*. Academic Press, 2014.

- Xiang- L. Gong, Y. and G. Liu . Fuzzy regression model based on geometric centroid and incentre points and application to performance evaluation. International Journal of Uncertainty, Fuzziness and Knowledge-Based Systems, 28:269–288, 2020.
- P Grzegorzewski . *Distribution-free tests for vague data* , In: ( Lopez-Diaz M , et al. (Eds) ) Soft Methodology and Random Information Systems. Springer, Heidelberg, 2004.
- P. Grzegorzewski . K-sample median test for vague data. International Journal of Intelligent Systems, 24:529–539, 2009.
- P. Grzegorzewski . Two-sample dispersion problem for fuzzy data. Information Processing and Management of Uncertainty in Knowledge-Based Systems, 1239:82–96, 2020.
- Dai- Z. Zhao A. Guan, H. and J. He . A novel stock forecasting model based on High-order-fuzzy-fluctuation trends and back propagation neural network. PLoS ONE, 13:40–51, 2018.
- Jain- G. Tayal D.K. Gupta, C. and O. Castillo . ClusFuDE: Forecasting low dimensional numerical data using an improved method based on automatic clustering, fuzzy relationships and differential evolution. Engineering Applications of Artificial Intelligence, 71:175–189, 2018.
- E. Haktanir and C. Kahraman . Z-fuzzy hypothesis testing in statistical decision making. Journal of Intelligent and Fuzzy Systems, 37:6545–6555, 2019.
- Yabing- Z.H.A. Zhang R. -Sun Q. He, Q. and Liu T . Reliability analysis for multi-state system based on triangular fuzzy variety subset bayesian networks. Eksplotacja Niezawodnosci-Maintenance and Reliability, 19:152–165, 2017.
- S. Heilpern . Fuzzy subsets of the space of probability measures and expected value of fuzzy variable. Fuzzy Sets and Systems, 54:301–309, 1993.
- Akbari- M.G. Hesamian, G. and M. Asadollahi . Fuzzy semi-parametric partially linear model with fuzzy inputs and fuzzy outputs. Expert Systems with Applications, 71:230–239, 2017.
- Akbari- M.G. Hesamian, G. and M. Asadollahi . Fuzzy semi-parametric partially linear model with fuzzy inputs and fuzzy outputs. Expert Systems With Applications, 71:230–239, 2017.
- Akbari- M.G. Hesamian, G. and E. Ranjbar . Exponentially weighted moving average control chart based on normal fuzzy random variables. International Journal of Fuzzy Systems, 21:1187–1195, 2019.
- Akbari- M.G. Hesamian, G. and V. Ranjbar . Some inequalities and limit theorems for fuzzy random variables adopted with a-values of fuzzy numbers. Soft Computing, 24:3797–3807, 2020.
- Akbari- M.G. Hesamian, G. and R. Yaghoobpoor . Quality control process based on fuzzy random variables. IEEE Transactions on Fuzzy Systems, 27:671–685, 2019.
- Akbari- M.G. Hesamian, G. and R. Yaghoobpoor . Quality Control Process based on Fuzzy Random Variables. IEEE Transactions on Fuzzy Systems, 27:671–685, 2019.
- Akbari- M.G. Hesamian, G. and J. Zendehdel . Location and scale fuzzy random variables. International Journal of Systems Science, 51:229–241, 2020.
- Akbari- M.G. Hesamian, G. and J. Zendehdel . Location and scale fuzzy random variables. International Journal of Systems Science, 51:229–241, 2020.
- G. Hesamian . Fuzzy similarity measure based on fuzzy sets. Control and Cybernetics, 46:23–40, 2017.
- G. Hesamian and M.G. Akbari . A semi-parametric model for time series based on fuzzy data. IEEE Transactions on Fuzzy Systems, 26:2953–2966, 2018.
- G. Hesamian and M.G. Akbari . Fuzzy absolute error distance measure based on a generalised difference operation. International Journal of Systems Science, 49:2454–2462, 2018.
- G. Hesamian and M.G. Akbari . Fuzzy absolute error distance measure based on a generalized difference operation. International Journal of Systems Science, 49:2454–2462, 2018.
- G. Hesamian and M.G. Akbari . Fuzzy Lasso regression model with exact explanatory variables and fuzzy responses. International Journal of Approximate Reasoning, 115:290–300, 2019.
- G. Hesamian and M.G. Akbari . Fuzzy Quantile Linear Regression model adopted with a semi-parametric technique based on fuzzy predictors and fuzzy responses. Expert systems with applications, 118:585–597, 2019.
- G. Hesamian and M.G. Akbari . A robust varying coefficient approach to fuzzy multiple regression model. Journal of Computational and Applied Mathematics, 371:23–34, 2020.
- G. Hesamian and M.G. Akbari . Fuzzy spline univariate regression with exact predictors and fuzzy responses. Journal of Computational and Applied Mathematics, 375:31–44, 2020.
- G. Hesamian and M.G. Akbari . Testing hypotheses for multivariate normal distribution with fuzzy random variables. International Journal of Systems Science, page DOI:

- 10.1080/00207721.2021.1936274, 2021.
- G. Hesamian and J. Chachi . Two-sample kolmogorov-smirnov fuzzy test for fuzzy random variables. *Statistical Papers*, 56:61–82, 2015.
- G. Hesamian and J. Chachi . Two-sample Kolmogorov-Smirnov fuzzy test for fuzzy random variables. *Statistical Papers*, 56:61–82, 2015.
- G. Hesamian and Akbari. M.G. A preference index for ranking closed intervals and fuzzy numbers. *International Journal of Uncertainty, Fuzziness and Knowledge-Based Systems*, 25:741–757, 2017.
- G. Hesamian and M. Shams . A Note on Fuzzy Probability of a Fuzzy Event. *International Journal of Intelligent*, 32:676–685, 2017.
- G. Hesamian and S.M. Taheri . Fuzzy empirical distribution: properties and applications. *Kybernetika*, 49:962–982, 2013.
- D.H. Hong and H.J. Kim . Marcinkiewicz-type law of large numbers for fuzzy random variables. *Fuzzy Sets Systems*, 64, 1994.
- D.H. Hong and H.J. Kim . Weak law of large numbers for i.i.d. fuzzy random variables. *Kybernetika*, 43:87–96, 2007.
- E. Hosseinzadeh and H. Hassanpour . Estimating the parameters of fuzzy linear regression model with crisp inputs and Gaussian fuzzy outputs: A goal programming approach. *Soft Computing*, 25:2719–2728, 2021.
- V. Hrissanthou and M. Spiliotis . Conventional and Fuzzy Regression: Theory and Engineering Applications. Nova Science Pub Inc, New York, 2018.
- O. Hryniwicz . Goodman-Kruskal measure of dependence for fuzzy ordered categorical data. *Computational Statistics and Data Analysis*, 51:323–334, 2006.
- O. Hryniwicz . Possibilistic decisions and fuzzy statistical tests. *Fuzzy Sets and Systems*, 157:2665–2673, 2006.
- Zuo- M.J. Huang, H.Z. and Z.Q. Sun . Bayesian reliability analysis for fuzzy lifetime data. *Fuzzy Sets and Systems*, 157:1674–1686, 2006.
- M. Hudec and D. Praenka . Collecting and managing fuzzy data in statistical relational databases. *Statistical Journal of the IAOS*, 32:245–255, 2016.
- M.A. Hussain and E.A. Amin . Fuzzy reliability estimation for exponential distribution using ranked set sampling. *International Journal of Contemporary Mathematical Sciences*, 12:31–42, 2017.
- H. Inoue . A strong law of large numbers for fuzzy random sets. *Fuzzy Sets Systems*, 41:285–291, 1991.
- C. Jershian and J.S. Yao . Fuzzy probability over fuzzy  $\sigma$ -field with fuzzy topological spaces. *Fuzzy Sets and Systems*, 116:201–223, 2000.
- Parchami- A. Jiryaei, A. and M. Mashinchi . One-Way ANOVA and least squares method based on fuzzy random variables. *Turkish Journal of Fuzzy Systems*, 4:18–33, 2013.
- S.Y. Joo . Weak laws of large numbers for fuzzy random variables. *Fuzzy Sets Systems*, 147:453–464, 2004.
- Mc Clelland GH Judd, CM and SR Carey . Data Analysis: A Model Comparison Approach To Regression, ANOVA, and Beyond ( 3rd ed.). Routledge, New York, 2017.
- Lee W.J. Jung H.Y. and Choi S.H. Hybrid fuzzy regression analysis using the F-Transform. *Applied Sciences*, 10:14–24, 2020.
- Bozdag C.F. Kahraman, C. and Ruan D. . Fuzzy sets approaches to statistical parametric and non-parametric tests. *International Journal of Intelligent Systems*, 19:1069–1078, 2004.
- D. Kalpanapriya and P. Pandian . Fuzzy hypothesis testing Of ANOVA model with fuzzy data. *International Journal of Modern Engineering Research*, 2:2951–2956, 2007.
- C. Kao and C.L. Chyu . Least-squares estimates in fuzzy regression analysis. *European Journal of Operational Research*, 148:426–435, 2003.
- Khan M.F. Aslam M. - Niaki S.T.A. Khan, M.Z. and Mughal A.R. . A fuzzy EWMA attribute control chart to monitor process mean. *Information*, 9:24–33, 2018.
- Z. Khan , Hashim R. Yaqoob N. Gulistan, M. , and Chamamm W. . Design of S-control chart for neutrosophic data: An application to manufacturing industry. *Journal of Intelligent and Fuzzy Systems*, 38:4743–4751, 2020.
- H. Kim and H.Y. Jung . Ridge fuzzy regression modelling for solving multicollinearity. *Mathematics*, 8:63–77, 2020.

- Y.K. Kim . A strong law of large numbers for fuzzy random variables. *Fuzzy Sets Systems*, 111:319–323, 2000.
- E.P. Klement . Some remarks on a paper of R.R. Yager. *Information Sciences*, 27:211–220, 1982.
- Puri M.L. Klement, E.P. and Ralescu D.A. . Limit theorems for fuzzy random variables. *Proceedings of the Royal Society A*, 407:171–182, 1986.
- GJ Klir and B Yuan . Fuzzy sets and fuzzy logic: theory and applications. Prentice Hall, 1995.
- JE Kolassa . An Introduction to Nonparametric Statistics. Chapman and Hall/CRC, 2020.
- Okuda T. Konishi, M. and Asai K. . Analysis of variance based on fuzzy interval data using moment correction method. *International Journal of Innovative Computing*, 2:83–99, 2006.
- V. Kratschmer . A unified approach to fuzzy random variables. *Fuzzy Sets Systems*, 123:1–9, 2001.
- R Kruse and KD Meyer . Statistics with vague data. Amsterdam, The Netherlands: Reidel, 1987.
- A. Kumar and P. Dhiman . Reliability estimation of a network structure using generalized trapezoidal fuzzy numbers. *Journal of Kabin*, 51:225–241, 2021.
- Ram M. Kumar, A. and Yadav O.P. . Advancements in Fuzzy Reliability Theory. Hershey, PA: IGI Global, 2021.
- H. Kwakernaak . Fuzzy random variables. Part I: definitions and theorems. *Information Science*, 19:1–15, 1987.
- H.T. Lee . Cpk index estimation using fuzzy numbers. *European Journal of Operational Research*, 129:683–688, 2001.
- KH Lee . First course on fuzzy theory and applications. Springer-Verlag: Berlin, 2005.
- Pedrycz W. Lee, S.H. and Sohn G. . Design of similarity and dissimilarity measures for fuzzy sets on the basis of distance measure. *International Journal of Fuzzy System*, 11:67–72, 2009.
- EL Lehmann and G Casella . Theory of Point Estimation ( 2nd ed.). Springer, 1998.
- R. Li . Water quality forecasting of Haihe River based on improved fuzzy time series model. *Desalination and Water Treatment*, 108:285291, 2018.
- Zeng W. Xie J. Li, J. and Yin Q. . A new fuzzy regression model based on least absolute deviation. *Engineering Applications of Artificial Intelligence*, 52:54–64, 2016.
- Wu B. Lin, P. and Watada J. . Kolmogorov-Smirnov two sample test with continuous fuzzy data. *Advances in Intelligent and Soft Computing*, 68:175–186, 2010.
- B Liu . Uncertainty Theory. Springer-Verlag: Berlin, 2004.
- Y.K. Liu and J. Gao . Convergence criteria and convergence relations for sequences of fuzzy random variables. *Fuzzy Knowledge Based Networks*, 3613:321–331, 2005.
- Y.K. Liu and B. Liu . Fuzzy random variables: A scalar expected value operator. *Fuzzy Optimization and Decision Making*, 2:143–160, 2003.
- M. Loeve . Probability Theory I. Springer-Verlag. New York, 1977.
- Magott J. Lower, M. and Skorupski J. . Analysis of air traffic incidents using event trees with fuzzy probabilities. *Fuzzy Sets and Systems*, 293:5079, 2016.
- M.A. Lubiano and W. Trutschnig . ANOVA for fuzzy random variables using the R-package SAFD; Combining soft Computing and statistical methods in data. *Analysis Advances in Intelligent and Soft Computing*, 77:449–456, 2010.
- A.S. Mendes , Rizol P. Machado, M.A.G. , and Rocha M.S. . Fuzzy control chart for monitoring mean and range of univariate processes. *Pesquisa Operacional*, 39:339–357, 2019.
- Jain D. Mishra, A.R. and Hooda D.S. . On fuzzy distance and induced fuzzy information measures. *Journal of Information and Optimization Sciences*, 37:193–211, 2016.
- I. Molchanov . On strong laws of large numbers for random upper semicontinuous. *Journal of Mathematical Analysis and Applications*, 235:349–355, 1999.
- Rodríguez G Gil Ma-Colubi A Montenegro, M and MR Casals . *Introduction to ANOVA with fuzzy random variables* , In Soft Methodology and Random Information Systems , eds. M. Lopez-Díaz , M.A. Gil , P. Grzegorzewski , O. Hryniewicz , and J. Lawry . Berlin, Springer, pp. 487–494, 2004.
- D.C. Montgomery . Introduction to statistical quality control. John Wiley and Sons, New York, 2007.
- J Munkres . Topology ( 2nd ed.). Prentice-Hall, Upper Saddle River, NJ, USA, 2000.
- N. Mylonas and B. Papadopoulos . Unbiased fuzzy estimators in fuzzy hypothesis testing. *Algorithms*, 14:185–192, 2021.

- Behzad M.H. Razzaghnia T. Naderkhani, R. and Farnoosh R. . Fuzzy regression analysis based on fuzzy neural networks using trapezoidal data. International Journal of Fuzzy Systems, In press <https://doi.org/10.1007/s40815-020-01033-2>, 2021.
- K. Nakamura . Preference relations on a set of fuzzy utilities as a basis for decision making. *Fuzzy Sets and Systems*, 20:147–162, 1986.
- Riaz M. Nasir, A. and Does R.J.M.M. . An EWMA-type control chart for monitoring the process mean using auxiliary information. *Communications in Statistics; Theory and Methods*, 43:3485–3498, 2014.
- Mashinchi M. Nourbakhsh, N. and Parchamie A. . Analysis of variance based on fuzzy observations. *International Journal of Systems Science*, 44:714–726, 2013.
- V. Novak . Detection of structural breaks in time series using fuzzy techniques. *International Journal of Fuzzy Logic and Intelligent Systems*, 18:1–12, 2018.
- TW O'Gorman . *Applied Adaptive Statistical Methods; Tests of Significance and Confidence Intervals*. SIAM, 2004.
- Parham G.A. Pak, A. and Saraj M. . Reliability estimation in Rayleigh distribution based on fuzzy lifetime data. *International Journal of System Assurance Engineering and Management*, 5, 2014.
- W Palma . *Time Series Analysis*. John Wiley and Sons, New York, 2016.
- D.C. Panda . A new method for evaluation of fuzzy reliability of multistage interconnection. *International Journal of Research in Applied Science*, 3:14–20, 2016.
- P. Pandian and D. Kalpanapriya . Two-factor ANOVA technique for fuzzy data having membership grades. *Research Journal of Pharmacy and Technology*, 9:2394–2402, 2016.
- A. Parchami . Fuzzy decision in testing hypotheses by fuzzy data: Two case studies. *Iranian Journal of Fuzzy Systems*, 17:127–136, 2020.
- A. Parchami and M. Mashinchi . Fuzzy estimation for process capability indices. *Information Sciences*, 177:1452–1462, 2007.
- Mashinchi M. Yavari A.R. Parchami, A. and Maleki H.R. . Fuzzy confidence intervals for fuzzy process capability index. *Journal of Intelligent and Fuzzy Systems*, 17:287–295, 2006.
- K. Patra . Fuzzy risk analysis using a new technique of ranking of generalized trapezoidal fuzzy numbers. *Granular Computing*, In press <https://doi.org/10.1007/s41066-021-00255-5>, 2021.
- W.L. Pearn and S. Kotz . *Encyclopedia and Handbook of Process Capability Indices*. Series on Quality. Reliability and Engineering Statistics (vol. 12). World Scientific publishing Co, Pte. Ltd, 2006.
- W Pedrycz . *An Introduction to Computing with Fuzzy Sets: Analysis, Design, and Applications*. Springer, 2021.
- N. Pekin Alakoc . Fuzzy X-bar and S-control charts based on confidence intervals. *Journal of Advanced Research in Natural and Applied Sciences*, 7:114–131, 2021.
- N. Pekin Alakoc and A. Apaydin . A fuzzy control chart approach for attributes and variables. *Engineering, Technology and Applied Science Research*, 8:3360–3365, 2018.
- Bigand A. Phan, T.T.H. and Caillault E.P. . A new fuzzy logic-based similarity measure applied to large gap imputation for uncorrelated multivariate time series. *Applied Computational Intelligence and Soft Computing*, 2018:1–15, 2018.
- U.M. Pirzada and D.C. Vakaskar . Existence of Hukuhara differentiability of fuzzy-valued functions. *Journal of the Indian Mathematical Society*, 87, 2017.
- K. Plasecki . Probability of fuzzy events defined as denumerable additivity measure. *Fuzzy Sets and Systems*, 17:271–287, 1985.
- O.I. Provtar and O.O. Provtar . Fuzzy probabilities of fuzzy events. *Cybernetics and Systems Analysis*, 56:171180, 2020.
- M.L. Puri and D. Ralescu . Strong law of large numbers for Banach space valued random variables. *Annals of Probability*, 11:222–224, 1983.
- M.L. Puri and D.A. Ralescu . Differentials of fuzzy functions. *Journal of Mathematical Analysis and Applications*, 91:552–558, 1983.
- Othman M. Sokkalingam R. Rahim, N.F. and Kadir E.A. . Forecasting crude palm oil prices using fuzzy rule-based time series method. *IEEE Access*, 2:32216–32224, 2018.
- S.W. Roberts . Control chart tests based on geometric moving averages. *Technometrics*, 1:239–250, 1959.
- Colubi A. Rodríguez, G. and Gil M.A. . Fuzzy data treated as functional data: a one-way ANOVA test approach. *Computational Statistics and Data Analysis*, 56:943–955, 2011.

- Montenegro M. Colubi A. Rodríguez, G. and Gil M.A. . Bootstrap techniques and fuzzy random variables: Synergy in hypothesis testing with fuzzy data. *Fuzzy Sets and Systems*, 157:2608–2613, 2006.
- R Rodriguez-Lopez . On boundary value problems for fuzzy differential equations, *Soft Methods for Handling Variability and Imprecision. Advances in Soft Computing*, V. 48, Springer, Berlin, Heidelberg, pp. 218225, 2008.
- Tahir M. Ubaid U.R. - Zeeshan A. Ronnason, C. and Aiyared I. . Some novel cosine similarity measures based on complex hesitant fuzzy sets and their applications. *Journal of Mathematics*, 4:1–9, 2021.
- TP Ryan . Modern regression methods. John Wiley and Sons, New York, 2008.
- H. Sabahno , S.M. Mousavi , and A. Amiri . A new development of an adaptive X-R control chart under a fuzzy environment. *International Journal of Data Mining, Modelling and Management*, 11:19–44, 2019.
- H. Scheffe . The analysis of variance. John Wiley and Sons, New York, 1999.
- H Scott E, Maxwell and D Delaney . Designing Experiments and Analyzing Data: A Model Comparison Perspective ( 2nd ed.). Routledge, New York, Erlbaum.
- Patra K. Sen, S. and Mondal S.K. . Similarity Measure of Gaussian Fuzzy Numbers and Its Application. *International Journal of Applied and Computational Mathematics*, 96:<https://doi.org/10.1007/s40819-021-01040-3>, 2021.
- M. Shafi and R. Viertl . Empirical reliability functions based on fuzzy life time data. *Journal of Intelligent and Fuzzy Systems*, 28:707–711.
- Heydari A. Kazemi- M. Shahsavari-Pour, N. and M. Karami . A novel method for ranking fuzzy numbers based on the different areas fuzzy number. *International Journal of Mathematics in Operational Research*, 11:544–566, 2017.
- F.A. Shapiro . Fuzzy random variables. *Insurance: Journal of Mathematical Economics*, 44:307–314, 2009.
- AK Sharma . Text Book of Correlations and Regression. Discovery Publishing House, New Delhi, 2005.
- NZ Shi and J Tao . Statistical Hypothesis Testing: Theory and Methods. World Scientific Publishing Company, Singapore, 2008.
- Nguyen T.L. Shu, H.M. and Hsu B.M. . Fuzzy MaxGWMA chart for identifying abnormal variations of on-line manufacturing processes with imprecise information. *Expert System With Application*, 41:1342–1356, 2013.
- H Shumway and DS Stoffer . Time series analysis and its applications. Springer-Verlag Berlin, Heidelberg, 2011.
- P. Smets . Probability of a fuzzy event: An axiomatic approach. *Fuzzy Sets and Systems*, 7:153–164, 1982.
- M. Smithson and J. Verkuilen . Fuzzy set theory: Applications in the social sciences. Sage, USA, 2006.
- L. Stefanini and B. Bede . Generalized Hukuhara differentiability of interval-valued functions and interval differential equations. *Nonlinear Analysis: Theory Methods and Applications*, 71:1311–1328, 2009.
- A. Syropoulos and T. Grammenos . A Modern Introduction to Fuzzy Mathematics. John Wiley and Sons, New York, 2020.
- A Taff . Hypothesis Testing: The Ultimate Beginner's Guide to Statistical Significance. CreateSpace Independent Publishing Platform, 2018.
- J. Talašová and O. Pavlačka . Fuzzy probability spaces and their applications in decision making. *Austrian Journal of Statistics*, 35, 2006.
- Seymour L. Taylor, R.L. and Chen Y. . Weak laws of large numbers for fuzzy random sets. *Nonlinear Analysis*, 47:1245–1256, 2001.
- H. Toth . Probabilities and fuzzy events: An operational approach. *Fuzzy Sets and Systems*, 48:113–127, 1992.
- L. Tran and L. Duckstein . Comparison of fuzzy numbers using a fuzzy distance measure. *Fuzzy Sets and Systems*, 130:331–341, 2002.
- Ye J. Unver M. TÂYRKARSLAN, E. and Olgun M. . Consistency fuzzy sets and a cosine similarity measure in fuzzy multiset setting and application to medical diagnosis. *Mathematical Problems in Engineering*, pages 1–12, 2021.

- F.M. Tseng and G.H. Tzeng . A fuzzy seasonal ARIMA model for forecasting. *Fuzzy Sets and Systems*, 126:367–376, 2002.
- T. Uemura . A law of large numbers for random sets. *Fuzzy Sets and Systems*, 59:181–188, 1993.
- K. Vännman . A unified approach to the capability indices. *Statistica Sinica*, 5:805–820, 1995.
- R. Viertl . Univariate statistical analysis with fuzzy data. *Computational Statistics and Data Analysis*, 51:133–147, 2006.
- R. Viertl . Statistical methods for fuzzy data. John Wiley and Sons, New York, 2011.
- W. Voxman . Some remarks on distances between fuzzy numbers. *Fuzzy Sets and Systems*, 100:353–365, 1998.
- X. Wang and E.E. Kerre . Reasonable properties for the ordering of fuzzy quantities (I). *Fuzzy Sets and Systems*, 118:375–385, 2001.
- Xu Y. Wang, G. and Qin S. . Basic fuzzy event space and probability distribution of probability fuzzy space. *Mathematics*, 7:542–553, 2019.
- Y. Wang , G. Xu and S. Qin . Basic fuzzy event space and probability distribution of probability fuzzy space. *Mathematics*, 2019:1–15, 2019.
- Y.M. Wang and Y. Luo . Area ranking of fuzzy numbers based on positive and negative ideal points. *Computers and Mathematic Application*, 58:1769–1779, 2009.
- Zhang W.X. Wang, N. and Mei C.L. . Fuzzy non-parametric regression based on local linear smoothing technique. *Information Sciences*, 177:3882–3900, 2007.
- L Wasserman . All of non-parametric statistics. Springer, New York, 2007.
- C.W. Wu . Decision-making in testing process performance with fuzzy data. *European Journal of Operational Research*, 193:499–509, 2009.
- H.C. Wu . The central limit theorems for fuzzy random variables. *Information Science*, 120:239–256, 1999.
- H.C. Wu . The laws of large numbers for fuzzy random variables. *Fuzzy Sets and Systems*, 116:245–262, 2000.
- H.C. Wu . Statistical hypotheses testing for fuzzy data. *Fuzzy Sets and Systems*, 175:30–56, 2005.
- H.C. Wu . Analysis of variance for fuzzy data. *International Journal of Systems Science*, 38:235–246, 2007.
- R.R. Yager . A note on probabilities of fuzzy events. *Information Sciences*, 18:113–122, 1979.
- R.R. Yager . A representation of the probability of a fuzzy subset. *Fuzzy Sets and Systems*, 13:273–283, 1984.
- L. Yang and B. Liu . On inequalities and critical values of fuzzy random variables. *International Journal of Uncertainty, Fuzziness and Knowledge-Based Systems*, 13:163–175, 2005.
- J.S. Yao and K. Wu . Ranking fuzzy numbers based on decomposition principle and signed distance. *Fuzzy Sets and Systems*, 116:275–288, 2000.
- J. Yen and R. Langari . Fuzzy Logic: Intelligence, Control, and Information. Prentice Hall, 1999.
- Y. Yuan . Criteria for evaluating fuzzy ranking methods. *Fuzzy Sets and Systems*, 43:139–157, 1991.
- S Zacks . Introduction to reliability analysis: probability models and statistical methods. Springer Science and Business Media, 2012.
- L.A. Zadeh . Probability measures of fuzzy events. *Journal of Mathematical Analysis and Applications*, 23:421–427, 1968.
- L.A. Zadeh . Fuzzy sets as a basis for a theory of possibility. *Fuzzy Sets and Systems*, 1:3–28, 1978.
- M.Gh. Zarei , R. Akbari and J. Chachi . Modeling autoregressive fuzzy time series data based on semi-parametric methods. *Soft Computing*, 24:7295–7304, 2020.
- Mirzazadeha A. Salehana- S. Zavar Sabegha, M.H. and G.W. Weber . A literature review on the fuzzy control chart; classifications and analysis. *International Journal of Supply and Operations Management*, 1:167–189, 2014.
- X.R. Zhao and B.Q. Hu . Fuzzy and interval-valued fuzzy decision-theoretic rough set approaches based on fuzzy probability measure. *Information Science*, 298:534554, 2015.
- H.J. Zimmermann . Fuzzy set theory and its applications. (4th ed.) Kluwer, 2001.