1. The probability of a leap year selected at random contain 53									
Sunday is:									
(a) 53/366	(b) 1/7	(c) 2/7	(d) 53/365						
2. A bag contain	s 3 red and 2 b	olue marbles. A n	narble is drawn at						
random. The pro	bability of drav	wing a black ball	is:						
(a) 3/5	(b) 2/5	(c) 0/5	(d) 1/5						
3. The probabilit	y that it will ra	in tomorrow is 0	.85. What is the						
	probability that it will not rain tomorrow								
			(d) none of these						
-	-		ed from the numbers						
(1, 2, 3,,15	•								
	• •	(c) 2/15	• •						
5. What are the									
		(c) 8							
-	-	number selecte	d at random from the						
numbers (1,2,3, .		() 10 10 7	(1)						
• •		. , ,	(d) none of these						
7. The sum of th	•								
, ,		$0 \qquad (d) \text{ none}$							
_	-	are given; choos	e the correct answer						
for that which is			(1)						
			(d) none of these.						
		nultaneously, tha	n the probability of						
getting at least two heads, is: (a) 1/4 (b) 3/8 (c) ½ (d) 1/8									
10. A letter is ch									
**A55A55INATI	UN. The pro	papility that the	letter chosen has: (d) none of these.						
(a) 6/13	(D) // 13	(C) I	(a) none of these.						
11 A diag is thre	wn Eindthan	robobility of gott	ing on oven number						
4 4	_		ing an even number.						
(A) 2/3	(b) i	(0) 3/0	D) 1/2						
12. Two coins are thrown at the same time. Find the probability of getting both heads.									
(A) 3/4 (B) 1/4		(D) 0							
13. Two dice are thrown simultaneously. The probability of getting a									

1

sum of 9 is:

(A) 1/10	(B) 3/10	(C) 1/9	(D) 4	./9		
	ds are numbe ime number.	ered from 1 to 1	00. Find the	probability of		
(A) 3/4	(B) 27/50	(C) 1/4	(D)) 29/100		
_	a blue ball is o	double that of a		If the probability en the number of		
` ,		, ,	` '	0		
	t random fron	ontains 12 defe n this box. Ther				
(A) 143/150		7/150 (C)	1/25	(D) 1/50		
mixed thoro	oughly. One callity that the n	umbers 2 to 10 ard is drawn fro umber on card (C) 3/10	om this box is a perfect	randomly, then square.		
18. What is (A) 1/7	s the probabili (B) 53/366	ity of getting 53 (C) 2/7	•	n a leap year? 7/366		
19. A card is drawn from a well shuffled deck of 52 cards. Find the probability of getting a king of red suit. (A) $1/26$ (B) $3/26$ (C) $7/52$ (D) $1/13$						
equally like 1,2,312	ly to come to ,then the prob	onsists of spinr rest pointing to pability that it w (C) 7/12	one of the	number an odd number is:		
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game. (A) $3/4$ (B) $1/2$ (C) 1 (D) $1/4$						

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is:							
(A) 364/365	(B) 31/365	(C) 1/365	(D) 1/133225				
23. A number x is chosen at random from the numbers -2, -1, 0, 1, 2. Then the probability that $x^2 < 2$ is? (A) $1/5$ (B) $2/5$ (C) $3/5$ (D) $4/5$							
24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is 2/3, then the number of white marbles in the jar is: (A) 10 (B) 6 (C) 8 (D) 7							
Then the prob	25. A number is selected at random from first 50 natural numbers. Then the probability that it is a multiple of 3 and 4 is: (A) $7/50$ (B) $4/25$ (C) $1/25$ (D) $2/25$						
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?							
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	d) 4/21				
	-	•	nes are 50, 70, 82,				
	ne standard devia b) 25.49		d) 25.69				
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.							
a) 13, 15	b) 13, 18	c) 18, 15	d) 13, 16				
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is							
a) $\frac{1}{2}$ 30. X is a varia	b) $\frac{1}{3}$	d 3. The value of	$\frac{d}{1}$ 6 E(X²) is				
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?							

32.Out of to probability?	_	alues, which	one is not poss	ible in
	b) ∑ x d) P(P(x) = 3 x) = -0.5		
33.If E(x) =	2 and E(z) = 4 b) 6	1, then E(z - c) 0	•	sufficient data
34.The cov	ariance of two	independe	nt random variab	le is
a) 1	b) 0	c) - 1	d) Ur	ndefined
35.If Σ P(x) a) 0) = k² – 8 then , b) 1	, the value o c) 3		sufficient data
• •	0.5 and x = 4, b) 0.5	, ,	? d) 2	
37.In a disc is always?	rete probabili	ty distributio	on, the sum of al	l probabilities
a) 0	b) Infinite	c) 1	d) Und	defined
38.If the pr	obability of hi	tting the tar	get is 0.4, find m	nean and
	b) 0.6,	0.24	c) 0.4, 0.16	d) 0.6, 0.16
-	% and if 10 bo	mbs are dro	pped from a place pped, find mean 0.4, 0.16	
a) 2	e mean of toss b) 4 s the mean and	c) 8	d) 1 or standard norm	nal distribution?

c) 5

d) 7

a) 3

b) 4

				•		l variance nd varianc			
42. \a) E(e of a rand b) E(X			given by 2) - (E(X))		. d) (E(X))2		
	43.Mean of a random variable X is given by a) E(X)								
44.N a) 0	44.Mean of a constant 'a' is a) 0								
	45.Variance of a constant 'a' is . a) 0								
46.Find the mean and variance of X?									
F	Х	0	1	2	3	4			
	f(x)	1/9	2/9	3/9	2/9	1/9			
	, 4/3	b) 3	3, 4/3		c) 2, 2/3	.	d) 3, 2/3		

47. Find the expectation of a random variable X?

	Г				 		
	Х	0	1	2	3		
	f(x)	1/6	2/6	2/6	1/6		
a) (0.5		b) 1.5		c) 2	2.5	d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by



c) np2q

d) npq2

- 49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**
- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx
- 50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**

- a) \sqrt{np} b) \sqrt{pq} c) (np)2