Standard Precaution



Dr.Eman.A.Mahmoud

Clinical microbiology & immunology

Infection control consultant

NLI

- standard Precautions are the minimum infection prevention practices that apply to all patient care, regardless of suspected or confirmed infection status of the patient, in any setting where health care is delivered



components


- hand hygiene
- respiratory hygiene and cough etiquette
- patient placement
- Personal protective equipment
- aseptic technique
- safe injections and sharps injury prevention
- Environmental cleaning
- handling of laundry and linen
- waste management
- Decontamination and reprocessing of reusable patient care items and equipment

Hand hygiene

- **Hand Hygiene includes cleaning hands with *soap and water or alcohol-based hand rub* in order to remove germs,**
- transmission of several key healthcare-associated pathogens including MRSA, VR
  difficile. All these pathogens hav
  demonstrated to persist in the e
  (in some cases months)



When


1
BEFORE
TOUCHING
A PATIENT
2
BEFORE
CLEAN/ASEPTIC
PROCEDURE
3
AFTER BODY
FLUID EXPOSURE
RISK
4
AFTER
TOUCHING
A PATIENT
5
AFTER
TOUCHING PATIENT
SURROUNDINGS



- **Before contact with a patient or patient environment**

this will protect the patient from harmful organisms carried on your hands.

Before aseptic/clean procedure

This prevents the patient's own organisms from entering his or her body.

Handling dressings or touching open wounds

Performing invasive procedure





After body fluid *exposure*

This can help protect yourself and the health care environment from harmful patient organisms.

After contact with a patient/patient env

Help protect yourself and the health care environment from harmful patient organisms



Methods

1. Washing by soap and water.
2. Alcoholic hand rub after dirt removal.
3. Surgical hand scrub before surgery.



Respiratory hygiene:



Respiratory hygiene:




- Cover nose, mouth during cough, sneeze with tissue or upper sleeve
- Discard tissue with respiratory secretion
- Wear surgical mask in contact with patient with respiratory disease


Patient placment

Single Patient Rooms : Recommended for**:**

- Patients with known or suspected infections that are contagious (e.g., certain respiratory infections, draining wounds).
- Patients requiring additional privacy or if their condition poses a risk to others.



- **Cohorting Patients: can be done when:**
  - Patients have the same infection type (e.g., same pathogen) and are at similar risk levels.
  - This approach can be useful during outbreaks but should be done with caution to prevent cross-contamination.



Personal protective equipment



- Equipment used to protect person during deals with patient or its towels or secretions, or body fluids
- Choice of PPE depends on type of infection between patient and health care worker and type of disease transmission
- E.g. gloves, gown, surgical mask, goggles face shield, eye protector glasses.
- Disposable PPE must be discarded in biohozand bags

Aseptic technique

- Aseptic technique is a fundamental practice in healthcare that aims to prevent contamination and reduce the risk of infection during medical procedures.



Safe injection practice

- Use new needle and syringe in every medication vial.
- Skin at site of injection is rubbed by antiseptic.
- Preparation of medicine in a clean areas.
- Medication vials septum are disinfected before use.
- Wear disposal gloves during injection.
- New needle and syringe for every patient.






- Use a medication vials for one patient only if possible.
- Don't use the same syringes between patients even you changed the needle.
- Don't use the same syringes used to infuse medication into an IV tubing port for anther patients.
- Hand hygiene during injection and medication preparation.
- Needles, syringes are discarded into sharp containers as biohazard.


Needle stick and sharps injury prevention

- Used needle should be discarded immediately and not recapped, or cut, removed from syringe.
- Any needles or sharps should be places in a leak proof puncture resistant container.
- The container to be filed to 2/3 only not more than discarded as biohazard label.



Environmental cleaning

- Removal of dirt, organic mater.
- Use detergent, disinfection.
- Allow air to dry the area or item



handling of laundry and linen

- Effective handling of laundry and linen is vital in controlling infection in healthcare settings.
- reduce the risk of cross-contamination and protect both patients and healthcare workers

- By appropriate segregation, transportation, laundering, and storage

Waste disposal

Safety boxes resistant for puncture and leak are used then when filled to 2/3 they are transported after closing and labeling as biohazard material to be discarded.

Other disposable PPE, b body, body fluid polluted items are discarded in biohazard bags resistant to leaks and puncture and labeled to be discarded.

APPLICATION

MAKE THE BEST USE OF EVERYTHING

DIFFERENT COLOR,DIFFERNT USAGE



BLACK

HOUSEHOLD GARBAGE



RED

RADIOACTIVE, CHEMOTHERAPEUTIC MEDICAL WASTE



YELLOW

INFECTIOUS, PATHOLOGICAL WASTE



GREEN

RECYCLABLE WASTE



PURPLE

LABORATORY CYTOTOXIC WASTE



CLEAR

HIGH TEMPERATURE STERILIZATION OF LABORATORY SUPPLIES

Decontamination and reprocessing of reusable patient care items and equipment

- Reprocessing includes cleaning, disinfecting, and sterilizing reusable items to make them safe for patient use.
- **Sterilization**

This is the complete elimination of all microorganisms, including spores, usually performed on critical items (those that enter sterile tissue).