

# Homework 5

niceguy

February 17, 2023

1. Given a continuous uniform distribution, show that

(a)  $\mu = \frac{A+B}{2}$

(b)  $\sigma^2 = \frac{(B-A)^2}{12}$

**Solution:** The mean is

$$\begin{aligned}\mu &= \int_{-\infty}^{\infty} xf(x)dx \\ &= \int_A^B \frac{x}{B-A}dx \\ &= \frac{B^2 - A^2}{2(B-A)} \\ &= \frac{A+B}{2}\end{aligned}$$

The variance is

$$\begin{aligned}\sigma^2 &= \int_{-\infty}^{\infty} (x - \mu)^2 f(x)dx \\ &= \int_A^B \frac{x^2 - (A+B)x + \frac{(A+B)^2}{4}}{B-A}dx \\ &= \frac{B^3 - A^3}{3(B-A)} - \frac{(A+B)(B^2 - A^2)}{2(B-A)} + \frac{(A+B)^2(B-A)}{4(B-A)} \\ &= \frac{A^2 + AB + B^2}{3} - \frac{(A+B)^2}{2} + \frac{(A+B)^2}{4} \\ &= \frac{4A^2 + 4AB + 4B^2 - 6A^2 - 12AB - 6B^2 + 3A^2 + 6AB + 3B^2}{12} \\ &= \frac{A^2 - 2AB + B^2}{12} \\ &= \frac{(B-A)^2}{12}\end{aligned}$$

2. A bus arrives every 10 minutes at a bus stop. It is assumed that the waiting time for a particular individual is a random variable with a continuous uniform distribution.
- (a) What is the probability that the individual waits more than 7 minutes?
- (b) What is the probability that the individual waits between 2 and 7 minutes?

**Solution:**

$$P[X > 7] = \frac{10 - 7}{10} = 0.3$$
$$P[2 \leq X \leq 7] = \frac{7 - 2}{10} = 0.5$$

3. Given a standard normal distribution, find the value of  $k$  such that

- (a)  $P(Z > k) = 0.2946$
- (b)  $P(Z < k) = 0.0427$
- (c)  $P(-0.93 < Z < k) = 0.7235$

**Solution:**

$$0.54, -1.72, 1.28$$

4. The loaves of rye bread distributed to local stores by a certain bakery have an average length of 30 centimeters and a standard deviation of 2 centimeters. Assuming that the lengths are normally distributed, what percentage of the loaves are

- (a) longer than 31.7 centimeters?
- (b) between 29.3 and 33.5 centimeters in length?
- (c) shorter than 25.5 centimeters?

**Solution:**

$$P(X > 31.7) = P(Z > 0.85) = 19.8\%$$
$$P(29.3 < X < 33.5) = P(-0.35 < Z < 1.75) = 59.7\%$$
$$P(X < 25.5) = P(Z < -2.25) = 1.22\%$$

5. If a set of observations is normally distributed, what percent of these differ from the mean by

- (a) more than  $1.3\sigma$ ?
- (b) less than  $0.52\sigma$ ?

**Solution:**

$$P(|Z| > 1.3) = 19.4\%$$
$$P(|Z| < 0.52) = 39.7\%$$

6. A coin is tossed 400 times. Use the normal curve approximation to find the probability of obtaining

- (a) between 185 and 210 heads inclusive
- (b) exactly 205 heads
- (c) fewer than 176 or more than 227 heads

**Solution:**

$$Z = \frac{X - np}{\sqrt{np(1-p)}} = \frac{X - 200}{10}$$

Then

$$P(184.5 \leq X \leq 210.5) = P(-1.55 \leq Z \leq 1.05) = 0.793$$

$$P(204.5 < X < 205.5) = P(0.45 < Z < 0.55) = 0.0352$$

$$P(X < 175.5) + P(X > 227.5) = P(Z < -2.45) + P(Z > 2.75) = 0.0101$$

7. A pair of dice is rolled 180 times. What is the probability that a total of 7 occurs

- (a) at least 25 times?
- (b) between 33 and 41 times inclusive?
- (c) exactly 30 times?

**Solution:** The probability that a total of 7 occurs is  $\frac{1}{6}$ . Thus

$$Z = \frac{X - np}{\sqrt{np(1-p)}} = \frac{X - 30}{5}$$

Then

$$P(X > 24.5) = P(Z > -1.1) = 0.864$$

$$P(32.5 < X < 41.5) = P(0.5 < Z < 2.3) = 0.298$$

$$P(29.5 < X < 30.5) = P(-0.1 < Z < 0.1) = 0.0797$$

8. Use the gamma function with  $y = \sqrt{2x}$  to show that  $\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$ .

**Solution:**

$$\begin{aligned}\Gamma\left(\frac{1}{2}\right) &= \int_0^\infty x^{\frac{1}{2}-1} e^{-x} dx \\ &= \int_0^\infty \frac{e^{-x}}{\sqrt{x}} dx \\ &= \sqrt{2} \int_0^\infty e^{-\frac{y^2}{2}} dy \\ &= \sqrt{2} \sqrt{\frac{\pi}{2}} \\ &= \sqrt{\pi}\end{aligned}$$

9. Suppose that the service life, in years, of a hearing aid battery is a random variable having a Weibull distribution with  $\alpha = \frac{1}{2}$  and  $\beta = 2$ .

- (a) How long can such a battery be expected to last?
- (b) What is the probability that such a battery will be operating after 2 years?

**Solution:** The mean is

$$\mu = \alpha^{-\frac{1}{\beta}} \Gamma\left(1 + \frac{1}{\beta}\right) = \sqrt{2} \Gamma(1.5) = \sqrt{2} \int_0^{\infty} \sqrt{x} e^{-x} dx = \frac{1}{\sqrt{2}} \int_0^{\infty} \frac{e^{-x}}{\sqrt{x}} dx = \sqrt{\frac{\pi}{2}}$$

The probability it operates after 2 years is

$$\begin{aligned} P(X > 2) &= \int_2^{\infty} \alpha \beta x^{\beta-1} e^{-\alpha x^{\beta}} dx \\ &= \int_2^{\infty} x e^{-\frac{x^2}{2}} dx \\ &= e^{-2} \end{aligned}$$

10. Let  $X$  be a random variable with probability

$$f(x) = \begin{cases} \frac{1}{3} & x = 1, 2, 3 \\ 0 & \text{elsewhere} \end{cases}$$

Find the probability distribution of the random variable  $Y = 2X - 1$ .

**Solution:**

$$g(y) = \begin{cases} \frac{1}{3} & y = 1, 3, 5 \\ 0 & \text{elsewhere} \end{cases}$$

11. Let  $X_1$  and  $X_2$  be discrete random variables with joint probability distribution

$$f(x_1, x_2) = \begin{cases} \frac{x_1 x_2}{18} & x_1 = 1, 2; x_2 = 1, 2, 3 \\ 0 & \text{elsewhere} \end{cases}$$

Find the probability distribution of the random variable  $Y = X_1 X_2$ .

**Solution:**

$$g(y) = \begin{cases} \frac{1}{18} & y = 1 \\ \frac{2}{9} & y = 2, y = 4 \\ \frac{1}{6} & y = 3 \\ \frac{1}{3} & y = 6 \end{cases}$$

12. Let  $X$  be a random variable with probability distribution

$$f(x) = \begin{cases} \frac{1+x}{2} & -1 < x < 1 \\ 0 & \text{elsewhere} \end{cases}$$

Find the probability distribution of the random variable  $Y = X^2$ .

**Solution:**

$$\begin{aligned} G(y) &= P(Y \leq y) \\ &= P(-\sqrt{y} \leq X \leq \sqrt{y}) \\ &= \int_{-\sqrt{y}}^{\sqrt{y}} f(t) dt \end{aligned}$$

Then

$$\begin{aligned} g(y) &= \frac{d}{dy} G(y) \\ &= f(\sqrt{y}) \times \frac{1}{2\sqrt{y}} - f(-\sqrt{y}) \times \frac{-1}{2\sqrt{y}} \\ &= \frac{1 + \sqrt{y}}{2} \times \frac{1}{2\sqrt{y}} + \frac{1 - \sqrt{y}}{2} \times \frac{1}{2\sqrt{y}} \\ &= \frac{1}{2\sqrt{y}} \end{aligned}$$