1. The probability of a leap year selected at random contain 53						
Sunday is:						
(a) 53/366	(b) 1/7	(c) 2/7	(d) 53/365			
			marble is drawn at			
random. The prol	bability of draw	ring a black ball	is:			
(a) 3/5	(b) 2/5	(c) 0/5	(d) 1/5			
3. The probabilit	y that it will rai	n tomorrow is 0	0.85. What is the			
probability that it						
• •	, ,	• •	(d) none of these			
_	-		ed from the numbers			
(1, 2, 3,,15	•					
	• •	(c) 2/15	7 7			
5. What are the						
		(c) 8				
-	•	number selecte	ed at random from the			
numbers (1,2,3, .	35) is :					
* *	· · ·	* *	(d) none of these			
7. The sum of th						
• •	1 /	0 (d) none				
_	-	re given; choos	e the correct answer			
for that which is	not possible.					
			(d) none of these.			
		ultaneously, tha	n the probability of			
getting at least tw	wo heads, is:					
		(c) $\frac{1}{2}$				
10. A letter is ch						
ASSASSINATION	ON. The prob	ability that the	letter chosen has: (d) none of these.			
(a) 6/13	(b) 7/13	(c) 1	(d) none of these.			
	_		ting an even number.			
(A) 2/3	(B) 1	(C) 5/6 (D) 1/2			
40 Torressing and the control of the						
12. Two coins are thrown at the same time. Find the probability of						
getting both heads. (A) 3/4 (B) 1/4 (C) 1/2 (D) 0						
(A) 3/4 (B) 1/4	(6) 1/2	(D) 0				
13. Two dice are thrown simultaneously. The probability of getting a						

sum of 9 is:

(A) 1/10	(B) 3/10	(C) 1/9	(D) 4/	9			
14. 100 cards are numbered from 1 to 100. Find the probability of getting a prime number.							
(A) 3/4	(B) 27/50	(C) 1/4	(D)	29/100			
15. A bag contains 5 red balls and some blue balls .If the probability of drawing a blue ball is double that of a red ball, then the number of blue balls in a bag is:							
(A) 5	(B) 10	(C) 15	(D) 20				
taken out at non-defecti	random from	ontains 12 defeanthis box. Then 7/150 (C)					
mixed thoro	oughly. One ca lity that the nu	umbers 2 to 101 ard is drawn fro umber on card i (C) 3/10	m this box ras s a perfect s	andomly, then quare.			
18. What is the probability of getting 53 Mondays in a leap year? (A) 1/7 (B) 53/366 (C) 2/7 (D) 7/366							
19. A card is drawn from a well shuffled deck of 52 cards. Find the probability of getting a king of red suit. (A) 1/26 (B) 3/26 (C) 7/52 (D) 1/13							
equally likel 1,2,312 ,	y to come to the then the prob	ensists of spinning to eability that it wing (C) 7/12	one of the n	number n odd number is:			
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game. (A) $3/4$ (B) $1/2$ (C) 1 (D) $1/4$							

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is:							
_			(D) 1/133225				
23. A number x is chosen at random from the numbers -2, -1, 0, 1, 2. Then the probability that $x^2 < 2$ is? (A) $1/5$ (B) $2/5$ (C) $3/5$ (D) $4/5$							
24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is 2/3, then the number of white marbles in the jar is: (A) 10 (B) 6 (C) 8 (D) 7							
Then the prob	is selected at ran ability that it is a 4/25 (C) 1/25	mu <mark>ltiple of</mark> 3 and	0 natural numbers. 4 is:				
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?							
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	d) $\frac{4}{21}$				
	_	_	nes are 50, 70, 82,				
	e standard deviate b) 25.49		d) 25.69				
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.							
a) 13, 15	b) 13, 18	c) 18, 15	d) 13, 16				
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is							
a) $^{1}/_{2}$	b) $^1\!/_3$ ate between 0 and	d 3. The value of	d) $\frac{1}{6}$ E(X²) is				
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?							

probability?		·	n one is not poss	ible in				
a) $P(x) = 1$ c) $P(x) = 0.5$	b) ∑ x l d) P(x	P(x) = 3 0 = -0.5						
33.If E(x) = a) 2	2 and E(z) = 4 b) 6	, then E(z - c) 0	•	ufficient data				
34.The cov	ariance of two	independe	nt random variab	le is				
a) 1	b) 0	c) - 1	d) Un	defined				
35.If Σ P(x) a) 0	b) 1	the value o c) 3		sufficient data				
• •	36.If P(x) = 0.5 and x = 4, then E(x) = ? a) 1 b) 0.5 c) 4 d) 2							
37.In a disc is always?	rete probability	y distributio	on, the sum of all	probabilities				
•	b) Infinite	c) 1	d) Und	lefined				
•	obability of hit	ting the tar	get is 0.4, find m	ean and				
variance. a) 0.4, 0.24	b) 0.6, 0).24	c) 0.4, 0.16	d) 0.6, 0.16				
39.If the probability that a bomb dropped from a place will strike the target is 60% and if 10 bombs are dropped, find mean and variance? a) 0.6, 0.24 b) 6, 2.4 c) 0.4, 0.16 d) 4, 1.6								
 40. Find the mean of tossing 8 coins. a) 2 b) 4 c) 8 d) 1 41. What is the mean and variance for standard normal distribution? 								

c) 5

<mark>d) 7</mark>

a) 3

b) 4

<mark>a) Mean is</mark> c) Mean is			•			
42.Variano a) E(X)					(<mark>))2</mark>	d) (E(X))2
43.Mean o a) E(X)			_	•		d) (E(X))2
44.Mean of a) 0	f a consta b) a	ant 'a' is _	c) a/2	<u> </u>	d) 1	
45.Varianc a) 0			is		d) 1	
46.Find the	e mean ar	nd variand	ce of X?			
Х	0	1	2	3	4	
f(x)	1/9	2/9	3/9	2/9	1/9	
a) 2, 4/3 47.Find the	b) expecta		random	c) 2, 2/3 variable `		d) 3, 2/3

	Х	0	1	2	3	
	f(x)	1/6	2/6	2/6	1/6	
a) ().5		b) 1.5		c) 2.5	d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

b) npq

c) np2q

d) npq2

49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**

a) P(X = x) = nCx px qx

- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx
- 50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**

- a) \sqrt{np} b) \sqrt{pq} c) (np)2 d) \sqrt{npq}