

Lesson 5.3-5.4

Quiz, 8 questions

1
point

1.

We use the continuous version of Bayes' theorem if:

- ☐ θ is continuous
 - ☐ Y is continuous
 - ☐ $f(y | \theta)$ is continuous
 - ☐ All of the above
 - ☐ None of the above
-

1
point

2.

Consider the coin-flipping example from the lesson. Recall that the likelihood for this experiment was Bernoulli with unknown probability of heads, i.e., $f(y | \theta) = \theta^y (1 - \theta)^{1-y} I_{\{0 \leq \theta \leq 1\}}$, and we started with a uniform prior on the interval $[0, 1]$.

After the first flip resulted in heads ($Y_1 = 1$), the posterior for θ became $f(\theta | Y_1 = 1) = 2\theta I_{\{0 \leq \theta \leq 1\}}$.

Now use this posterior as your prior for θ before the next (second) flip. Which of the following represents the posterior PDF for θ after the second flip also results in heads ($Y_2 = 1$)?

- ☐ $f(\theta | Y_2 = 1) = \frac{(1-\theta) \cdot 2\theta}{\int_0^1 (1-\theta) \cdot 2\theta d\theta} I_{\{0 \leq \theta \leq 1\}}$
 - ☐ $f(\theta | Y_2 = 1) = \frac{\theta \cdot 2\theta}{\int_0^1 \theta \cdot 2\theta d\theta} I_{\{0 \leq \theta \leq 1\}}$
 - ☐ $f(\theta | Y_2 = 1) = \frac{\theta(1-\theta) \cdot 2\theta}{\int_0^1 \theta(1-\theta) \cdot 2\theta d\theta} I_{\{0 \leq \theta \leq 1\}}$
-

1
point

3.

Consider again the coin-flipping example from the lesson. Recall that we used a $\text{Uniform}(0,1)$ prior for θ . Which of the following is a correct interpretation of $P(0.3 < \theta < 0.9) = 0.6$?

- ☐ (0.3, 0.9) is a 60% credible interval for θ before observing any data.
 - ☐ (0.3, 0.9) is a 60% credible interval for θ after observing $Y = 1$.
 - ☐ (0.3, 0.9) is a 60% confidence interval for θ .
 - ☐ The posterior probability that $\theta \in (0.3, 0.9)$ is 0.6.
-

1
point

4.

Consider again the coin-flipping example from the lesson. Recall that the posterior PDF for θ , after observing $Y = 1$, was $f(\theta | Y = 1) = 2\theta I_{\{0 \leq \theta \leq 1\}}$. Which of the following is a correct interpretation of $P(0.3 < \theta < 0.9 | Y = 1) = \int_{0.3}^{0.9} 2\theta d\theta = 0.72$?

- ☐ (0.3, 0.9) is a 72% credible interval for θ before observing any data.
 - ☐ (0.3, 0.9) is a 72% credible interval for θ after observing $Y = 1$.
 - ☐ (0.3, 0.9) is a 72% confidence interval for θ .
 - ☐ The prior probability that $\theta \in (0.3, 0.9)$ is 0.72.
-

1
point

5.

Which two quantiles are required to capture the middle 90% of a distribution (thus producing a 90% equal-tailed interval)?

- ☐ .05 and .95
- ☐ .10 and .90
- ☐ .025 and .975

☐ 0 and .9

1
point

6.

Suppose you collect measurements to perform inference about a population mean θ . Your posterior distribution after observing data is $\theta \mid \mathbf{y} \sim N(0, 1)$.

Report the upper end of a 95% equal-tailed interval for θ . Round your answer to two decimal places.

Enter answer here

1
point

7.

What does "HPD interval" stand for?


- ☐ Highest partial density interval
 - ☐ Highest point distance interval
 - ☐ Highest posterior density interval
 - ☐ Highest precision density interval
-

1
point


8.

Each of the following graphs depicts a 50% credible interval from a posterior distribution. Which of the intervals represents the HPD interval?


- ☐ 50% interval: $\theta \in (0.196, 0.567)$




☐ 50% interval: $\theta \in (0.400, 0.756)$



☐ 50% interval: $\theta \in (0.500, 1.000)$



☐ 50% interval: $\theta \in (0.326, 0.674)$



☐ I, **Md Yousuf Ali**, understand that submitting work that isn't my own may result in permanent failure of this course or deactivation of my Coursera account.

[Learn more about Coursera's Honor Code](#)

Submit Quiz

