

✖ Try again once you are ready.

Required to pass: 80% or higher

You can retake this quiz up to 3 times every 8 hours.

[Back to Week 3](#)[Retake](#)

1. For which of the following situations would a quadratic loss function make the most sense?

1 / 1
points



A doctor's estimate for the life expectancy of a terminally ill patient.

Correct

Correct Answer. Large mistakes in estimating life expectancy would particularly painful for both the patient and their family.

This question refers to the following learning objective(s):

- Understand the concept of loss functions and how they relate to Bayesian decision making.



Your prediction for the number of bikes sold this year by a local bike shop.



Your prediction as to whether it will rain tomorrow.



Your answer choice on a Coursera multiple choice quiz.



1 / 1
points

2.

Fill in the blank: Under a **quadratic loss function**, the summary statistic that minimizes the posterior expected loss is the _____ of the posterior.

☐ Median

☐ Mode

☒ Mean

Correct

Correct Answer. The mean is the summary statistic that minimizes the posterior expected loss under the quadratic loss function.

This question refers to the following learning objective(s):

- Understand the concept of loss functions and how they relate to Bayesian decision making



0 / 1
points

3.

You are employed in the Human Resources department of a company and are asked to predict the number of employees that will quit their jobs in the coming year. Based on a Bayesian analysis of historical data, you determine the posterior predictive distribution of the number of quitters to follow a Poisson($\lambda = 10$) distribution. Given a **0/1** loss function, what is the prediction that minimizes posterior expected loss?

☐ a. 9

☒ b. 10

This should not be selected

10 is a mode of Poisson($\lambda = 10$), but not the only one. Since the loss function is 0/1, the mode of the posterior distribution minimizes posterior expected loss.

This question refers to the following learning objective(s):

- Make optimal decisions given a posterior distribution and a loss function.

- ☐ c. 11
- ☐ d. Either a or b



0 / 1
points

4. Suppose you are testing a new drug to determine whether it is more effective than an existing drug. Let H_0 denote the hypothesis that the new drug is no more effective than the existing drug and H_1 denote the hypothesis that the new drug is more effective than the existing drug. If you accept H_1 when in fact H_0 is true, the loss is 100. If you accept H_0 when in fact H_1 is true, the loss is 40. Assume that no loss is incurred if you accept a true hypothesis. At what posterior probability of H_0 will you be indifferent between the two hypotheses?



$\frac{1}{4}$

This should not be selected

If p is the posterior probability that H_0 is true, then the expected loss under H_0 is $40(1 - p)$ and the expected loss under H_1 is $100p$. Set the two equal and solve for p .

This question refers to the following learning objective(s):

- Decide between hypotheses given a loss function: Make optimal decisions given a

posterior distribution and a loss function.

- ☐ $\frac{2}{5}$
- ☐ $\frac{2}{7}$
- ☐ $\frac{1}{3}$



1 / 1
points

5. You are testing a hypothesis H_1 against an alternative hypothesis H_2 using Bayes Factors. You calculate $BF[H_1 : H_2]$ to be 0.427. According to guidelines first given by Jeffreys (and presented in the lecture), what conclusion can be drawn from the data?

- ☐ The data provides strong evidence against H_2 .
- ☐ The data provides strong evidence against H_1 .
- ☒ The data provides little to no evidence against H_1 .

Correct

Refer to lecture "Posterior probabilities of hypotheses and Bayes factors" to review interpretation of Bayes Factors. The greater $BF[H_1 : H_2]$, the stronger the evidence against H_2 .

This question refers to the following learning objective(s):

- Compare multiple hypotheses using Bayes Factors.

- ☐ The data provides significant evidence against H_1 .

Week 3 Quiz

Quiz, 10 questions

7/10 points (70.00%)



0 / 1
points

6. Suppose that you are trying to estimate the true proportion p of male births in the United States. Starting with a strong prior (Beta(500,500)) on the proportion, you randomly sample 10,000 birth certificates, observing 5029 males and 4971 females. What is the posterior probability that p is less than or equal to 0.5? (Hint: use function(s) in R to answer this question)

☐ 0.27

☒ 0.28



This should not be selected

Recall the conjugacy of the Beta-Binomial model. Use the pbeta function in R to calculate the posterior probability that p is less than 0.5.

This question refers to the following learning objective(s):

- Conceptualize Lindley's paradox and how the Bayes Factor depends on prior elicitation
- Make inferences about a proportion using a conjugate Beta prior.

☐ 0.29

☐ 0.30



1 / 1
points

7. Suppose that you are trying to estimate the true proportion p of male births in the United States. You want to evaluate two hypotheses against each other: first the null hypothesis (H_0) that $p = 0.5$ and an alternative hypothesis (H_1) that $p \neq 0.5$. To do this,

you assign a point-null prior to $p = 0.5$ under H_0 and a uniform $Beta(1, 1)$ prior to p under H_1 . Then, if we define k to be the number of male births out of a total sample of n birth certificates,

$$P(k|H_0) = \binom{n}{k} 0.5^k (1 - 0.5)^{n-k}$$

$$P(k|H_1) = \int_0^1 \binom{n}{k} p^k (1 - p)^{n-k} dp = \frac{1}{n+1}$$

Using the dbinom function in R calculate the Bayes Factor $BF[H_1 : H_0]$ if you observed 5029 male births out of 10,000 birth certificates.

☐ $BF[H_1 : H_0] = 67.44$

☐ $BF[H_1 : H_0] = 8.33$

☐ $BF[H_1 : H_0] = 0.12$

☒ $BF[H_1 : H_0] = 0.015$

Correct

This question refers to the following learning objective(s):

- Conceptualize Lindley's paradox and how the Bayes Factor depends on prior elicitation
- Compare multiple hypotheses using Bayes Factors.



1 / 1
points

8. Suppose you are trying to infer the difference in the proportion of yellow M&Ms between two bags of M&Ms. Each bag is estimated to contain roughly 30 M&Ms. To make inferences about the proportion, you draw 10 M&Ms from each bag **without** replacement and record the number of yellow M&Ms in each bag.

Is there a problem with the experimental design? If so, what is it?

- ☒ Yes, the probability of drawing a yellow M&M is not independent within groups.

Correct

This question refers to the following learning objective(s):

- Identify assumptions relating to a statistical inference.
- ☐ Yes, the probability of drawing a yellow M&M is not independent between groups.
- ☐ There is no problem with the experimental design.



1 / 1
points

9. Suppose that when testing $H_0 : p = 0.5$ versus $H_1 : p \neq 0.5$ using Bayes Factors, we get the posterior probability $P(H_0 \mid \text{data}) = 0.25$. Conditional on H_1 , the posterior mean of p is 0.6. Under **quadratic** loss, what is the point estimate for p that minimizes expected posterior loss?

- ☐ 0.5
- ☐ 0.55
- ☒ 0.575

Correct

This question refers to the following learning objective(s):

- Create point estimates and credible intervals by averaging over multiple hypotheses
- Make optimal decisions given a posterior

distribution and a loss function.

☐ 0.6



1 / 1
points

10. True or False: The use of the reference prior $Beta(1/2, 1/2)$ has little bearing on the posterior distribution of a proportion p , provided that the sample size is sufficiently large.

☒ True

Correct

This question refers to the following learning objective(s):

- Conceptualize Lindley's paradox and how the Bayes Factor depends on prior elicitation.

☐ False

