Chapter 5. Entire Functions

Jensen's formula, order of growth, infinite products, Weierstrass and Hadamard factorization

Contents

1 Jensen's formula **2**

2 Functions of finite order **3**

3 Infinite products **4**

3.1 Generalities 4

3.2 Example: the product formula for the sine function 4

4 Weierstrass infinite products **5**

5 Hadamard's factorization theorem **6**

Overview

Entire functions are holomorphic in the whole complex plane. The chapter is organized around three questions.

1. *Where can they vanish?* The obvious necessary condition is sufficient: for any sequence $\{z_n\}$ with no limit point in $\mathbb{C}$ there is an entire function vanishing exactly there. The construction is inspired by Euler's product for $\sin \pi z$, refined by the *Weierstrass canonical factors*.
2. *How do they grow at infinity?* The larger a function is, the more zeros it can have; for a polynomial of degree d, $\sup_{|z|=R}|P(z)| \approx R^d$ and P has d zeros. A precise version is *Jensen's formula*, linking the number of zeros in a disc to the logarithmic average of $|f|$ on the circle.
3. *To what extent are they determined by their zeros?* If f has finite order, it is determined by its zeros up to a simple factor: the *Hadamard factorization theorem*.

1 Jensen's formula

D_R and C_R denote the open disc and circle of radius R centered at 0. We exclude the function that vanishes identically.

Theorem 1.1 (Jensen's formula)

Let Ω be an open set that contains the closure of a disc D_R and suppose that f is holomorphic in Ω, $f(0) \neq 0$, and f vanishes nowhere on the circle C_R. If $z_1, \dots, z_N$ denote the zeros of f inside the disc (counted with multiplicities), then

$$\log|f(0)| = \sum_{k=1}^{N} \log\left(\frac{|z_k|}{R}\right) + \frac{1}{2\pi}\int_0^{2\pi} \log|f(Re^{i\theta})|\, d\theta. \tag{1}$$

Proof.

Step 1. If f_1, f_2 satisfy the hypotheses and (1), so does $f_1 f_2$, since $\log xy = \log x + \log y$ for $x, y > 0$ and the zeros of $f_1 f_2$ are the union of those of f_1 and f_2.

Step 2. $g(z) = f(z)/((z-z_1)\cdots(z-z_N))$, defined on $\Omega - \{z_1, \dots, z_N\}$, is bounded near each z_j, so each z_j is a removable singularity: $f(z) = (z-z_1)\cdots(z-z_N)g(z)$ with g holomorphic in Ω and nowhere vanishing on $\overline{D_R}$. By Step 1 it suffices to prove (1) for such g and for functions $z - z_j$.

Step 3. For g nowhere vanishing on $\overline{D_R}$ we must show $\log|g(0)| = \frac{1}{2\pi}\int_0^{2\pi} \log|g(Re^{i\theta})|\, d\theta$. In a slightly larger disc, $g = e^h$ with h holomorphic (discs are simply connected; Theorem 6.2). Then $|g| = e^{\mathrm{Re}(h)}$, so $\log|g| = \mathrm{Re}(h)$, and the mean value property (Corollary 7.3) gives the formula.

Step 4. For $f(z) = z - w$, $w \in D_R$, we must show $\log|w| = \log\frac{|w|}{R} + \frac{1}{2\pi}\int_0^{2\pi} \log|Re^{i\theta} - w|\, d\theta$. Since $\log(|w|/R) = \log|w| - \log R$ and $\log|Re^{i\theta} - w| = \log R + \log|e^{i\theta} - w/R|$, it suffices to show $\int_0^{2\pi} \log|e^{i\theta} - a|\, d\theta = 0$ for $|a| < 1$, equivalently (via $\theta \mapsto -\theta$, $|e^{i\theta} - a| = |1 - ae^{-i\theta}|$) $\int_0^{2\pi} \log|1 - ae^{i\theta}|\, d\theta = 0$. $F(z) = 1 - az$ vanishes nowhere on the closed unit disc, so $F = e^G$ in a disc of radius > 1 and $\log|F| = \mathrm{Re}(G)$ is harmonic. Since $\log|F(0)| = 0$, the mean value property gives the result. ∎

For f holomorphic on $\overline{D_R}$, let $n(r) = n_f(r)$ be the number of zeros of f (with multiplicities) in D_r, $0 < r < R$; $n(r)$ is non-decreasing. If $f(0) \neq 0$ and f does not vanish on C_R, then

$$\int_0^R n(r)\frac{dr}{r} = \frac{1}{2\pi}\int_0^{2\pi} \log|f(Re^{i\theta})|\, d\theta - \log|f(0)|, \tag{2}$$

which follows from Jensen's formula and:

Lemma 1.2

If $z_1, \dots, z_N$ are the zeros of f inside the disc D_R, then

$$\int_0^R n(r)\frac{dr}{r} = \sum_{k=1}^{N} \log\left|\frac{R}{z_k}\right|.$$

Proof. $\sum_{k=1}^N \log|\frac{R}{z_k}| = \sum_{k=1}^N \int_{|z_k|}^R \frac{dr}{r}$. With $\eta_k(r) = 1$ if $r > |z_k|$ and 0 otherwise, $\sum_k \eta_k(r) = n(r)$, so $\sum_k \int_{|z_k|}^R \frac{dr}{r} = \sum_k \int_0^R \eta_k(r)\frac{dr}{r} = \int_0^R n(r)\frac{dr}{r}$. ∎

2 Functions of finite order

Definition (order of growth)

An entire function f has an **order of growth** $\le \rho$ if $|f(z)| \le Ae^{B|z|^\rho}$ for all $z \in \mathbb{C}$, for some constants $A, B > 0$. The **order of growth** of f is $\rho_f = \inf \rho$ over all such $\rho > 0$.

For example, e^{z^2} has order of growth 2.

Theorem 2.1

If f is an entire function that has an order of growth $\le \rho$, then: (i) $n(r) \le Cr^\rho$ for some $C > 0$ and all sufficiently large r. (ii) If $z_1, z_2, \dots$ denote the zeros of f, with $z_k \neq 0$, then for all $s > \rho$ we have $\displaystyle\sum_{k=1}^{\infty} \frac{1}{|z_k|^s} < \infty$.

Proof. It suffices to treat $f(0) \neq 0$: otherwise use $F(z) = f(z)/z^\ell$, ℓ the order of the zero at 0; n_f and n_F differ by a constant and F also has order of growth $\le \rho$. By (2) with $R = 2r$,

$$\int_r^{2r} n(x)\frac{dx}{x} \le \int_0^{2r} n(x)\frac{dx}{x} = \frac{1}{2\pi}\int_0^{2\pi} \log|f(2re^{i\theta})|\,d\theta - \log|f(0)|.$$

Since n is increasing, the left side is $\ge n(r)\int_r^{2r}\frac{dx}{x} = n(r)\log 2$; and by the growth condition $\int_0^{2\pi}\log|f(2re^{i\theta})|\,d\theta \le \int_0^{2\pi}\log|Ae^{B(2r)^\rho}|\,d\theta \le C'r^\rho$ for large r. Hence $n(r) \le Cr^\rho$. For (ii),

$$\sum_{|z_k|\ge 1}|z_k|^{-s} = \sum_{j=0}^{\infty}\sum_{2^j\le|z_k|<2^{j+1}}|z_k|^{-s} \le \sum_{j=0}^{\infty} 2^{-js}n(2^{j+1}) \le c\sum_{j=0}^{\infty}2^{-js}2^{(j+1)\rho} \le c'\sum_{j=0}^{\infty}(2^{\rho-s})^j < \infty,$$

since $s > \rho$. ∎

The condition $s > \rho$ cannot be improved:

Example 1

$f(z) = \sin \pi z = \frac{e^{i\pi z} - e^{-i\pi z}}{2i}$ satisfies $|f(z)| \le e^{\pi|z|}$, and taking $z = ix$ shows the order of growth is exactly 1. f vanishes to order 1 at each $n \in \mathbb{Z}$, and $\sum_{n\neq 0} 1/|n|^s < \infty$ precisely when $s > 1$.

Example 2

$f(z) = \cos z^{1/2} = \sum_{n=0}^{\infty}(-1)^n\frac{z^n}{(2n)!}$ is entire with $|f(z)| \le e^{|z|^{1/2}}$, of order $1/2$. It vanishes at $z_n = ((n+1/2)\pi)^2$, and $\sum_n 1/|z_n|^s < \infty$ exactly when $s > 1/2$.

Given $z_1, z_2, \ldots$, is there an entire f with zeros exactly there? Necessarily $\lim_{k\to\infty}|z_k| = \infty$ (Theorem 4.8). Weierstrass showed this suffices, by inserting factors into $(z-z_1)(z-z_2)\cdots$ that guarantee convergence without introducing new zeros.

3 Infinite products

3.1 Generalities

For complex a_n, the product $\prod_{n=1}^{\infty}(1+a_n)$ **converges** if $\lim_{N\to\infty}\prod_{n=1}^{N}(1+a_n)$ exists.

Proposition 3.1

If $\sum|a_n| < \infty$, then the product $\prod_{n=1}^{\infty}(1+a_n)$ converges. Moreover, the product converges to 0 if and only if one of its factors is 0.

Proof. For large n, $|a_n| < 1/2$; discarding finitely many terms, assume this for all n. Define $\log(1+a_n)$ by the power series (6) of Chapter 3, so $1+z = e^{\log(1+z)}$ for $|z|<1$. Then $\prod_{n=1}^{N}(1+a_n) = \prod_{n=1}^{N} e^{\log(1+a_n)} = e^{B_N}$ with $B_N = \sum_{n=1}^{N} b_n$, $b_n = \log(1+a_n)$. From the power series, $|\log(1+z)| \le 2|z|$ for $|z| < 1/2$, so $|b_n| \le 2|a_n|$ and $B_N \to B$. By continuity of the exponential, $e^{B_N} \to e^B$. If $1+a_n \neq 0$ for all n, the limit is $e^B \neq 0$. ∎

Proposition 3.2

Suppose $\{F_n\}$ is a sequence of holomorphic functions on the open set Ω. If there exist constants $c_n > 0$ such that $\sum c_n < \infty$ and $|F_n(z) - 1| \le c_n$ for all $z \in \Omega$, then: (i) The product $\prod_{n=1}^{\infty} F_n(z)$ converges uniformly in Ω to a holomorphic function $F(z)$. (ii) If $F_n(z)$ does not vanish for any n, then

$$\frac{F'(z)}{F(z)} = \sum_{n=1}^{\infty} \frac{F_n'(z)}{F_n(z)}.$$

Proof. (i) Write $F_n(z) = 1 + a_n(z)$ with $|a_n(z)| \le c_n$ and argue as in Proposition 3.1; the estimates are uniform in z since the c_n are constants. The uniform limit is holomorphic (Theorem 5.2). (ii) Let $K \subset \Omega$ be compact and $G_N = \prod_{n=1}^{N} F_n$. Since $G_N \to F$ uniformly, Theorem 5.3 gives $G_N' \to F'$ uniformly on K. G_N is uniformly bounded below on K, so $G_N'/G_N \to F'/F$ uniformly on K; and $G_N'/G_N = \sum_{n=1}^{N} F_n'/F_n$ (Section 4). ∎

3.2 Example: the product formula for the sine function

$$\sin \pi z = \pi z \prod_{n=1}^{\infty}\left(1 - \frac{z^2}{n^2}\right) \tag{3}$$

for all $z \in \mathbb{C}$; it is derived from the sum formula for the cotangent

$$\pi \cot \pi z = \sum_{n=-\infty}^{\infty} \frac{1}{z+n} = \lim_{N\to\infty} \sum_{|n|\le N} \frac{1}{z+n} = \frac{1}{z} + \sum_{n=1}^{\infty} \frac{2z}{z^2 - n^2}, \tag{4}$$

valid for $z \notin \mathbb{Z}$. The halves of $\sum 1/(z+n)$ with $n > 0$ and $n < 0$ diverge separately; only the symmetric interpretation converges.

Proof of (4). $F(z) = \pi \cot \pi z$ has the properties: (i) $F(z+1) = F(z)$ for $z \notin \mathbb{Z}$; (ii) $F(z) = \frac{1}{z} + F_0(z)$ with F_0 analytic near 0; (iii) F has simple poles at the integers and no other singularities.

The function $\sum_{n=-\infty}^{\infty} \frac{1}{z+n} = \lim_N \sum_{|n|\le N} \frac{1}{z+n}$ has the same three properties: for (i),

$$\sum_{|n|\le N} \frac{1}{z+1+n} = \frac{1}{z+1+N} - \frac{1}{z-N} + \sum_{|n|\le N} \frac{1}{z+n},$$

and let $N \to \infty$; (ii), (iii) are evident from $\frac{1}{z} + \sum_{n\ge 1} \frac{2z}{z^2-n^2}$. Hence $\Delta(z) = F(z) - \sum \frac{1}{z+n}$ is periodic with period 1, has a removable singularity at 0 by (ii), hence at all integers by periodicity, and is entire. It suffices to show Δ is bounded, hence (by periodicity) bounded in the strip $|\mathrm{Re}(z)| \le 1/2$. It is bounded in the rectangle $|\mathrm{Im}(z)| \le 1$. For $z = x + iy$ with $y > 1$,

$$\pi \cot \pi z = \pi i \frac{e^{i\pi z} + e^{-i\pi z}}{e^{i\pi z} - e^{-i\pi z}} = \pi i \frac{e^{-2\pi y} + e^{-2\pi i x}}{e^{-2\pi y} - e^{-2\pi i x}}$$

is bounded, and

$$\Big|\frac{1}{z} + \sum_{n=1}^{\infty} \frac{2z}{z^2-n^2}\Big| = \Big|\frac{1}{x+iy} + \sum_{n=1}^{\infty} \frac{2(x+iy)}{x^2-y^2-n^2+2ixy}\Big| \le C + C\sum_{n=1}^{\infty} \frac{y}{y^2+n^2}.$$

The last sum is majorized by $\int_0^{\infty} \frac{y}{y^2+x^2}\,dx$ (the integrand is decreasing in x), which is independent of y (change variables $x \mapsto yx$). Similarly for $y < -1$. So Δ is bounded on $\mathbb{C}$, hence constant by Liouville; Δ is odd, so the constant is 0. ∎

Proof of (3). Let $G(z) = \sin \pi z$ and $P(z) = \pi z \prod_{n=1}^{\infty}(1 - z^2/n^2)$. Since $\sum 1/n^2 < \infty$, Proposition 3.2 shows P converges and, away from the integers, $\frac{P'(z)}{P(z)} = \frac{1}{z} + \sum_{n=1}^{\infty} \frac{2z}{z^2-n^2}$. Since $G'/G = \pi \cot \pi z$, (4) gives

$$\Big(\frac{P}{G}\Big)' = \frac{P}{G}\Big[\frac{P'}{P} - \frac{G'}{G}\Big] = 0,$$

so $P = cG$. Dividing by z and letting $z \to 0$, $c = 1$. ∎

4 Weierstrass infinite products

Theorem 4.1 (Weierstrass)

Given any sequence $\{a_n\}$ of complex numbers with $|a_n| \to \infty$ as $n \to \infty$, there exists an entire function f that vanishes at all $z = a_n$ and nowhere else. Any other such entire function is of the form $f(z)e^{g(z)}$, where g is entire.

Repetitions are allowed in $\{a_n\}$, so this gives entire functions with prescribed zeros *and multiplicities* (the multiplicity of a zero a is the m with $f(z) = (z-a)^m g(z)$, $g(a) \ne 0$).

Proof. If f_1, f_2 both vanish exactly at the a_n (with multiplicities), f_1/f_2 has removable singularities there, so it is entire and nowhere vanishing; hence $f_1/f_2 = e^g$ for an entire g (Theorem 6.2). This proves the last statement.

To construct f: the naive product $\prod(1 - z/a_n)$ converges only for suitable $\{a_n\}$, so we insert exponential factors. Define the **canonical factors** of degree k:

$$E_0(z) = 1 - z, \qquad E_k(z) = (1-z)e^{z+z^2/2+\cdots+z^k/k} \quad (k \ge 1).$$

Suppose f is to have a zero of order m at the origin and $a_1, a_2, \ldots$ are non-zero. The **Weierstrass product** is

$$f(z) = z^m \prod_{n=1}^{\infty} E_n(z/a_n).$$

Fix $R > 0$ and $|z| < R$. Factors with $|a_n| \le 2R$ are finitely many, and their product vanishes exactly at the a_n with $|a_n| < R$. If $|a_n| > 2R$ then $|z/a_n| \le 1/2$, and by Lemma 4.2 (with c independent of n)

$$|1 - E_n(z/a_n)| \le c\Big|\frac{z}{a_n}\Big|^{n+1} \le \frac{c}{2^{n+1}}.$$

By Section 3, $\prod_{|a_n|>2R} E_n(z/a_n)$ is holomorphic and non-vanishing in $|z| < R$. As R is arbitrary, f has the required properties. ■

Lemma 4.2

If $|z| \le 1/2$, then $|1 - E_k(z)| \le c|z|^{k+1}$ for some $c > 0$.

Proof. For $|z| \le 1/2$, $1 - z = e^{\log(1-z)}$ (power series), so $E_k(z) = e^{\log(1-z)+z+z^2/2+\cdots+z^k/k} = e^w$ with $w = -\sum_{n=k+1}^{\infty} z^n/n$. Since $|z| \le 1/2$,

$$|w| \le |z|^{k+1} \sum_{n=k+1}^{\infty} |z|^{n-k-1}/n \le |z|^{k+1} \sum_{j=0}^{\infty} 2^{-j} \le 2|z|^{k+1}.$$

In particular $|w| \le 1$, so $|1 - E_k(z)| = |1 - e^w| \le c'|w| \le c|z|^{k+1}$. ■

Remark

The constant c is independent of k: one may take $c' = e$, $c = 2e$.

5 Hadamard's factorization theorem

Weierstrass's theorem writes a function with zeros $a_1, a_2, \ldots$ as $e^{g(z)} z^m \prod E_n(z/a_n)$. Hadamard showed that for functions of finite order the degree of the canonical factors can be taken constant, and g is then a polynomial. Recall that if f has order of growth $\le \rho$, then $n(r) \le Cr^\rho$ for large r and $\sum |a_n|^{-s} < \infty$ for $s > \rho$ (Theorem 2.1).

Theorem 5.1 (Hadamard)

Suppose f is entire and has growth order ρ_0. Let k be the integer so that $k \le \rho_0 < k + 1$. If $a_1, a_2, \ldots$ denote the (non-zero) zeros of f, then

$$f(z) = e^{P(z)} z^m \prod_{n=1}^{\infty} E_k(z/a_n),$$

where P is a polynomial of degree $\le k$, and m is the order of the zero of f at $z = 0$.

Main lemmas.

Lemma 5.2

The canonical products satisfy

$$|E_k(z)| \ge e^{-c|z|^{k+1}} \text{ if } |z| \le 1/2 \qquad \text{and} \qquad |E_k(z)| \ge |1 - z|e^{-c'|z|^k} \text{ if } |z| \ge 1/2.$$

Proof. If $|z| \le 1/2$, $E_k(z) = e^{\log(1-z)+\sum_{n=1}^{k} z^n/n} = e^{-\sum_{n=k+1}^{\infty} z^n/n} = e^w$, and $|e^w| \ge e^{-|w|}$ with $|w| \le c|z|^{k+1}$. If $|z| \ge 1/2$, $|E_k(z)| = |1 - z||e^{z+z^2/2+\cdots+z^k/k}|$, and $|e^{z+\cdots+z^k/k}| \ge e^{-|z+\cdots+z^k/k|} \ge e^{-c'|z|^k}$ (as $|z| \ge 1/2$). ■

The key is a lower bound for the product of canonical factors away from the zeros.

Lemma 5.3

For any s with $\rho_0 < s < k+1$, we have

$$\Big|\prod_{n=1}^{\infty} E_k(z/a_n)\Big| \geq e^{-c|z|^s},$$

except possibly when z belongs to the union of the discs centered at a_n of radius $|a_n|^{-k-1}$, for $n = 1, 2, 3, \ldots$.

Proof. Write $\prod_n E_k(z/a_n) = \prod_{|a_n|\leq 2|z|} E_k(z/a_n) \prod_{|a_n|>2|z|} E_k(z/a_n)$. For the second product, with no restriction on z, Lemma 5.2 gives

$$\Big|\prod_{|a_n|>2|z|} E_k(z/a_n)\Big| \geq \prod_{|a_n|>2|z|} e^{-c|z/a_n|^{k+1}} = e^{-c|z|^{k+1}\sum_{|a_n|>2|z|}|a_n|^{-k-1}},$$

and since $|a_n| > 2|z|$ and $s < k+1$, $|a_n|^{-k-1} = |a_n|^{-s}|a_n|^{s-k-1} \leq C|a_n|^{-s}|z|^{s-k-1}$; as $\sum|a_n|^{-s} < \infty$, this product is $\geq e^{-c|z|^s}$. For the first product, by the second part of Lemma 5.2,

$$\Big|\prod_{|a_n|\leq 2|z|} E_k(z/a_n)\Big| \geq \prod_{|a_n|\leq 2|z|} \Big|1 - \frac{z}{a_n}\Big| \prod_{|a_n|\leq 2|z|} e^{-c'|z/a_n|^k}. \tag{5}$$

Here $\prod_{|a_n|\leq 2|z|} e^{-c'|z/a_n|^k} = e^{-c'|z|^k \sum_{|a_n|\leq 2|z|}|a_n|^{-k}}$ and $|a_n|^{-k} = |a_n|^{-s}|a_n|^{s-k} \leq C|a_n|^{-s}|z|^{s-k}$, so this is $\geq e^{-c|z|^s}$. For the first factor of (5) we need the restriction on z: if z is not in the disc of radius $|a_n|^{-k-1}$ about a_n, then $|a_n - z| \geq |a_n|^{-k-1}$, so

$$\prod_{|a_n|\leq 2|z|} \Big|1 - \frac{z}{a_n}\Big| = \prod_{|a_n|\leq 2|z|} \Big|\frac{a_n - z}{a_n}\Big| \geq \prod_{|a_n|\leq 2|z|} |a_n|^{-k-1}|a_n|^{-1} = \prod_{|a_n|\leq 2|z|} |a_n|^{-k-2}.$$

Finally $\log \prod_{|a_n|\leq 2|z|} |a_n|^{k+2} = (k+2)\sum_{|a_n|\leq 2|z|} \log|a_n| \leq (k+2)n(2|z|)\log 2|z| \leq c|z|^s \log 2|z| \leq c'|z|^{s'}$ for any $s' > s$, using $n(2|z|) \leq c|z|^s$ (Theorem 2.1). Since $s > \rho_0$ was arbitrary, starting with s close enough to ρ_0 gives the lemma (with s' in place of s). ∎

Corollary 5.4

There exists a sequence of radii $r_1, r_2, \ldots$ with $r_m \to \infty$, such that

$$\Big|\prod_{n=1}^{\infty} E_k(z/a_n)\Big| \geq e^{-c|z|^s} \qquad \text{for } |z| = r_m.$$

Proof. Since $\sum|a_n|^{-k-1} < \infty$, there is N with $\sum_{n\geq N}|a_n|^{-k-1} < 1/10$. Given consecutive large integers L, $L+1$, there is $r \in [L, L+1]$ such that the circle of radius r about 0 meets none of the forbidden discs of Lemma 5.3: otherwise the intervals $I_n = [|a_n| - |a_n|^{-k-1}, |a_n| + |a_n|^{-k-1}]$, of length $2|a_n|^{-k-1}$, would cover $[L, L+1]$, giving $2\sum_{n\geq N}|a_n|^{-k-1} \geq 1$, a contradiction. Apply Lemma 5.3 with $|z| = r$. ∎

Proof of Hadamard's theorem. Let $E(z) = z^m \prod_{n=1}^{\infty} E_k(z/a_n)$. E is entire, as in the proof of Theorem 4.1: by Lemma 4.2, $|1 - E_k(z/a_n)| \leq c|z/a_n|^{k+1}$ for large n, and $\sum|a_n|^{-k-1}$ converges ($\rho_0 < s < k+1$). E has the zeros of f, so f/E is entire and nowhere vanishing: $f(z)/E(z) = e^{g(z)}$ for an entire g. Since f has growth order ρ_0 and by Corollary 5.4,

$$e^{\mathrm{Re}(g(z))} = \Big|\frac{f(z)}{E(z)}\Big| \leq c'e^{c|z|^s} \qquad \text{for } |z| = r_m,$$

so $\mathrm{Re}(g(z)) \leq C|z|^s$ for $|z| = r_m$. Lemma 5.5 gives that g is a polynomial of degree $\leq s$, hence $\leq k$ (as $s < k+1$). ∎

Lemma 5.5

Suppose g is entire and $u = \operatorname{Re}(g)$ satisfies $u(z) \le Cr^s$ whenever $|z| = r$, for a sequence of positive real numbers r that tends to infinity. Then g is a polynomial of degree $\le s$.

Proof. Write $g(z) = \sum_{n=0}^{\infty} a_n z^n$. By Theorem 7.1,

$$\frac{1}{2\pi}\int_0^{2\pi} g(re^{i\theta})e^{-in\theta}\,d\theta = \begin{cases} a_n r^n & n \ge 0, \\ 0 & n < 0. \end{cases} \tag{6}$$

Taking complex conjugates, $\frac{1}{2\pi}\int_0^{2\pi} \overline{g(re^{i\theta})}e^{-in\theta}\,d\theta = 0$ for $n > 0$ (7). Since $2u = g + \overline{g}$, adding (6) and (7),

$$a_n r^n = \frac{1}{\pi}\int_0^{2\pi} u(re^{i\theta})e^{-in\theta}\,d\theta \quad (n > 0), \qquad 2\operatorname{Re}(a_0) = \frac{1}{\pi}\int_0^{2\pi} u(re^{i\theta})\,d\theta.$$

(Supplied completion.) Since $\int_0^{2\pi} e^{-in\theta}d\theta = 0$ for $n > 0$, $a_n r^n = -\frac{1}{\pi}\int_0^{2\pi}[Cr^s - u(re^{i\theta})]e^{-in\theta}\,d\theta$, and $Cr^s - u \ge 0$ on the circles in question, so

$$|a_n| r^n \le \frac{1}{\pi}\int_0^{2\pi}[Cr^s - u(re^{i\theta})]\,d\theta = 2Cr^s - 2\operatorname{Re}(a_0).$$

Letting $r \to \infty$ along the sequence, $a_n = 0$ for $n > s$. ∎