Chapter 7. Function Spaces

The space C(X), Arzelà–Ascoli, Weierstrass approximation, sequence spaces, improper integrals, spaces of integrable functions

Contents

7.1 The space $C(X)$ **2**

7.2 Sequence spaces **3**

7.3 Improper integrals **3**

7.4 Spaces of integrable functions **4**

7.1 The space $C(X)$

A **norm** on a real or complex vector space V is a function $\|\cdot\|$ with $\|v\| \geq 0$ (equality iff $v = 0$), $\|\lambda v\| = |\lambda|\|v\|$, and $\|u + v\| \leq \|u\| + \|v\|$; V is then a **normed space**, with distance $\|u - v\|$, and convergence, Cauchy sequences, open and closed sets are defined as in $\mathbb{R}^n$. A normed space in which every Cauchy sequence converges is **complete** (a **Banach space**). For compact $X \subset \mathbb{R}^n$, $C(X)$ is the space of continuous $f : X \to \mathbb{C}$ with $\|f\|_{\sup} = \max_X |f|$; convergence in this norm is uniform convergence.

Theorem 7.1 (7.1.1)

For compact X, every Cauchy sequence in $C(X)$ converges: $C(X)$ is complete.

Proof. A Cauchy sequence in $C(X)$ is uniformly Cauchy, so converges uniformly (Theorem 6.2) to a function that is continuous (Theorem 6.1). ■

The Heine–Borel theorem fails in $C(X)$: the functions $f_n(x) = x^n$ on $[0, 1]$ form a closed bounded set (no subsequence converges uniformly, since the pointwise limit is discontinuous). The missing ingredient is equicontinuity. A family $\mathcal{F} \subset C(X)$ is **equicontinuous** if for every $\epsilon > 0$ there is $\delta > 0$ with $|f(x) - f(y)| < \epsilon$ for all $f \in \mathcal{F}$ and $\|x - y\| < \delta$; it is **bounded** if $\sup_{f\in\mathcal{F}} \|f\|_{\sup} < \infty$.

Proposition 7.2 (7.1.2)

If X is compact and (f_n) converges uniformly in $C(X)$, then $\{f_n\}$ is equicontinuous.

Proof. Given ϵ, choose N with $\|f_n - f_N\|_{\sup} < \epsilon/3$ for $n \geq N$ (Theorem 6.2). The finitely many functions $f_1, \dots, f_N$ are uniformly continuous (Theorem 3.18 of Volume I), so some δ works for all of them with $\epsilon/3$. For $n > N$ and $\|x - y\| < \delta$, $|f_n(x) - f_n(y)| \leq |f_n(x) - f_N(x)| + |f_N(x) - f_N(y)| + |f_N(y) - f_n(y)| < \epsilon$. ■

Theorem 7.3 (7.1.3, Arzelà–Ascoli)

If X is compact and $\mathcal{F} \subset C(X)$ is bounded and equicontinuous, every sequence in $\mathcal{F}$ has a uniformly convergent subsequence.

Proof. Let (f_n) be in $\mathcal{F}$, $|f_n| \leq M$. X has a countable dense subset $E = \{p_1, p_2, \dots\}$ (for each k, finitely many balls of radius $1/k$ cover X; take their centres in X). The bounded sequence $(f_n(p_1))$ has a convergent subsequence $(f_{1,n})$; $(f_{1,n}(p_2))$ has a convergent subsequence $(f_{2,n})$; and so on. The diagonal sequence $g_n = f_{n,n}$ is, from the k-th term on, a subsequence of $(f_{k,n})$, so $(g_n(p_k))$ converges for every k. Given ϵ, take δ from equicontinuity for $\epsilon/3$, and finitely many $p_{k_1}, \dots, p_{k_r}$ with every $x \in X$ within δ of one of them. Choose N with $|g_m(p_{k_j}) - g_n(p_{k_j})| < \epsilon/3$ for $m, n \geq N$ and all j. For any x, pick $p = p_{k_j}$ with $\|x - p\| < \delta$; then $|g_m(x) - g_n(x)| \leq |g_m(x) - g_m(p)| + |g_m(p) - g_n(p)| + |g_n(p) - g_n(x)| < \epsilon$. So (g_n) is uniformly Cauchy and converges in $C(X)$ (Theorem 7.1). ■

Combined with Proposition 7.2: a subset of $C(X)$ has compact closure iff it is bounded and equicontinuous.

Theorem 7.4 (7.1.4, Weierstrass approximation)

For every continuous $f : [a, b] \to \mathbb{R}$ there are polynomials $P_n \to f$ uniformly on $[a, b]$.

Proof. By an affine change of variable take $[a, b] = [0, 1]$, and subtracting the linear function $f(0) + x(f(1) - f(0))$ assume $f(0) = f(1) = 0$. Extend f by 0 outside $[0, 1]$; it is uniformly continuous on $\mathbb{R}$. Let $Q_n(x) = c_n(1 - x^2)^n$ on $[-1, 1]$ with c_n chosen so that $\int_{-1}^{1} Q_n = 1$. Since

$(1-x^2)^n \geq 1-nx^2$,

$$\int_{-1}^{1}(1-x^2)^n dx \geq 2\int_0^{1/\sqrt{n}}(1-nx^2)dx = \frac{4}{3\sqrt{n}} > \frac{1}{\sqrt{n}},$$

so $c_n < \sqrt{n}$, and for $\delta \leq |x| \leq 1$, $Q_n(x) \leq \sqrt{n}(1-\delta^2)^n \to 0$ uniformly. Let

$$P_n(x) = \int_{-1}^{1} f(x+t)Q_n(t)\,dt = \int_0^1 f(t)Q_n(t-x)\,dt \qquad (0 \leq x \leq 1),$$

a polynomial in x. Given ϵ, choose δ with $|f(y)-f(x)| < \epsilon/2$ for $|y-x| < \delta$, and let $M = \max|f|$. Then

$$|P_n(x)-f(x)| \leq \int_{-1}^{1}|f(x+t)-f(x)|Q_n(t)\,dt \leq 2M\cdot 2\sqrt{n}(1-\delta^2)^n + \frac{\epsilon}{2} < \epsilon$$

for large n, uniformly in $x \in [0,1]$. ■

For complex-valued f, approximate real and imaginary parts. Consequently $C[a,b]$ has a countable dense set (polynomials with rational coefficients).

7.2 Sequence spaces

$\ell^1 = \ell^1(\mathbb{N})$ is the space of sequences $a = (a(n))$ with $\|a\|_1 = \sum|a(n)| < \infty$, and ℓ^2 those with $\|a\|_2 = (\sum|a(n)|^2)^{1/2} < \infty$. ℓ^2 is a vector space with inner product $\langle a,b\rangle = \sum a(n)\overline{b(n)}$, which converges absolutely since $|a(n)b(n)| \leq \frac{1}{2}(|a(n)|^2+|b(n)|^2)$; $\|\cdot\|_2$ is the associated norm (Theorem 6.17).

Theorem 7.5 (7.2.1)

ℓ^1 is complete.

Proof. Let (a_m) be Cauchy. Since $|a_m(n)-a_k(n)| \leq \|a_m-a_k\|_1$, each $(a_m(n))_m$ converges to some $a(n)$. Given ϵ, choose N with $\|a_m-a_k\|_1 < \epsilon$ for $m,k \geq N$. For each L, $\sum_{n\leq L}|a_m(n)-a_k(n)| < \epsilon$; let $k\to\infty$ and then $L\to\infty$: $\sum_n|a_m(n)-a(n)| \leq \epsilon$ for $m \geq N$. So $a = a_m-(a_m-a) \in \ell^1$ and $a_m \to a$. ■

Theorem 7.6 (7.2.2)

ℓ^2 is complete.

Proof. As in Theorem 7.5: the coordinates converge, and $\sum_{n\leq L}|a_m(n)-a_k(n)|^2 < \epsilon^2$ passes to the limit in k and then in L. ■

The same proof shows that ℓ^∞ (bounded sequences with the sup norm) is complete. The unit vectors e_n in ℓ^2 form a bounded sequence with $\|e_m-e_n\|_2 = \sqrt{2}$, so without convergent subsequence: closed bounded sets in infinite-dimensional spaces need not be compact.

7.3 Improper integrals

Let I be an interval and $f: I \to \mathbb{C}$ Riemann integrable on every compact subinterval. If $I = [a,b)$ ($b \leq \infty$), the **improper integral** $\int_I f = \lim_{c\to b^-}\int_a^c f$ when the limit exists; similarly at a left end point, and for open intervals both ends are treated separately. We write $\int_I|f| < \infty$ when this improper integral converges (**absolute convergence**).

Proposition 7.7 (7.3.1)

If $0 \le f \le g$ on I and $\int_I g$ converges, then $\int_I f$ converges.

Proof. $c \mapsto \int_a^c f$ is increasing and bounded by $\int_I g$, so it has a limit as $c \to b^-$ (as for monotone sequences). ■

Proposition 7.8 (7.3.2, comparison test)

If $|f| \le g$ and $\int_I g$ converges, then $\int_I |f|$ and $\int_I f$ converge and $|\int_I f| \le \int_I |f| \le \int_I g$.

Proof. $\int_I |f|$ converges by Proposition 7.7. Cauchy criterion: for $a < c < c' < b$, $|\int_c^{c'} f| \le \int_c^{c'} |f| \to 0$ as $c, c' \to b$, so $\lim \int_a^c f$ exists (as in Theorem 2.14 of Volume I). The inequality passes to the limit from finite intervals (Theorem 5.5 of Volume I). ■

Example 7.1

$\int_1^\infty x^{-p} dx < \infty$ iff $p > 1$, and $\int_0^1 x^{-p} dx < \infty$ iff $p < 1$. $\int_0^\infty \frac{\sin x}{x}\, dx$ converges (integrate by parts: $\int_1^c \frac{\sin x}{x} dx = \frac{\cos 1}{1} - \frac{\cos c}{c} - \int_1^c \frac{\cos x}{x^2} dx$) but not absolutely (on $[k\pi, (k+1)\pi]$, $\int |\sin x|/x \ge \frac{2}{(k+1)\pi}$).

Proposition 7.9 (7.3.3, integral test)

If $f : [1, \infty) \to (0, \infty)$ is decreasing, then $\sum f(n)$ and $\int_1^\infty f$ converge or diverge together; indeed $\sum_{n=2}^N f(n) \le \int_1^N f \le \sum_{n=1}^{N-1} f(n)$, and if $f \to 0$, then $\sum_{n \le N} f(n) - \int_1^N f$ converges.

Proof. On $[n, n+1]$, $f(n+1) \le f \le f(n)$; integrate and sum. The differences $d_N = \sum_{n \le N} f(n) - \int_1^N f$ decrease ($d_N - d_{N+1} = \int_N^{N+1} f - f(N+1) \ge 0$) and are $\ge f(N) > 0$, so converge. ■

For $f(x) = 1/x$ this gives Euler's constant $\gamma = \lim(1 + \frac{1}{2} + \cdots + \frac{1}{N} - \log N)$.

Theorem 7.10 (7.3.4)

If $\int_I |f| < \infty$, then for every $\epsilon > 0$ there is $\delta > 0$ such that $\int_x^y |f| < \epsilon$ whenever $x < y$ in I and $y - x < \delta$.

Proof. Choose a compact $[c, d] \subset I$ with $\int_{I - [c,d]} |f| < \epsilon/2$; f is bounded on $[c, d]$, say by K. With $\delta = \epsilon/(2K+1)$, $\int_x^y |f| \le \int_{[x,y] \cap [c,d]} |f| + \int_{I-[c,d]} |f| < K\delta + \epsilon/2 < \epsilon$. ■

7.4 Spaces of integrable functions

Proposition 7.11 (7.4.1)

For a normed space X the following are equivalent:
(a) every Cauchy sequence in X converges;
(b) whenever $\sum \|x_n\| < \infty$, the series $\sum x_n$ converges in X.

Proof. (a)⇒(b): the partial sums are Cauchy, since $\|\sum_{m < k \le n} x_k\| \le \sum_{m < k \le n} \|x_k\|$.

(b)⇒(a): given a Cauchy (y_n), choose $n_1 < n_2 < \cdots$ with $\|y_m - y_n\| < 2^{-k}$ for $m, n \ge n_k$. With $x_1 = y_{n_1}$, $x_{k+1} = y_{n_{k+1}} - y_{n_k}$, $\sum \|x_k\| < \infty$, so $y_{n_k} = \sum_{j \le k} x_j$ converges; a Cauchy sequence with a convergent subsequence converges (Lemma 2.13 of Volume I, same proof). ■

Let $\mathcal{R}^1(I)$ and $\mathcal{R}^2(I)$ be the locally Riemann integrable $f : I \to \mathbb{C}$ with $\int_I |f| < \infty$, resp. $\int_I |f|^2 < \infty$, with $\|f\|_1 = \int_I |f|$ and $\|f\|_2 = (\int_I |f|^2)^{1/2}$. These are seminorms ($\|f\| = 0$ does not force $f = 0$), and $\mathcal{R}^2$ carries the inner product $\langle f, g\rangle = \int_I f\bar{g}$. Neither space is complete; this is the reason for introducing the Lebesgue integral (Chapter 10). A **step function** is a finite linear combination of characteristic functions of bounded intervals.

Lemma 7.12 (7.4.2)

For $f \in \mathcal{R}^2(I)$ and $\epsilon > 0$ there is a step function s with $\|f - s\|_2 < \epsilon$.

Proof. Treating real and imaginary parts separately, let f be real. Choose a compact $[a, b] \subset I$ with $\int_{I-[a,b]} |f|^2 < \epsilon^2/4$. On $[a, b]$, f is Riemann integrable, so bounded by some K; choose a partition with $U(f, P) - L(f, P) < \epsilon^2/(8K)$ and let $s = m_i$ on the i-th interval (and 0 off $[a, b]$). Then $0 \le f - s \le 2K$ on $[a, b]$, so $\int_a^b |f - s|^2 \le 2K \int_a^b (f - s) \le 2K(U - L) < \epsilon^2/4$. Altogether $\|f - s\|_2^2 < \epsilon^2/2$. ∎

Lemma 7.13 (7.4.3)

For a step function s on I and $\epsilon > 0$ there is a continuous g with $\|s - g\|_2 < \epsilon$ (and $g = 0$ outside a compact subinterval).

Proof. It suffices to treat $s = \chi_{[a,b]}$ (with any type of end points). Let g be 1 on $[a + \eta, b - \eta]$, 0 outside (a, b), and linear in between; then $|s - g| \le 1$ and differs from 0 only on two intervals of length η, so $\|s - g\|_2^2 \le 2\eta$. ∎

Theorem 7.14 (7.4.4)

For $f \in \mathcal{R}^2(I)$ and $\epsilon > 0$ there is a continuous g with $\|f - g\|_2 < \epsilon$.

Proof. Combine Lemmas 7.12 and 7.13 with the triangle inequality for $\|\cdot\|_2$. ∎

Corollary 7.15 (7.4.5)

For $f \in \mathcal{R}^2[a, b]$ there are polynomials P_n with $\|f - P_n\|_2 \to 0$.

Proof. Choose continuous g with $\|f - g\|_2 < \epsilon$ and a polynomial P with $\|g - P\|_{\sup} < \epsilon$ (Theorem 7.4); then $\|g - P\|_2 \le \epsilon\sqrt{b - a}$. ∎

Theorem 7.16 (7.4.6)

For $f \in \mathcal{R}^1(I)$ and $\epsilon > 0$ there is a step function s with $\|f - s\|_1 < \epsilon$, and a continuous g vanishing outside a compact subinterval with $\|f - g\|_1 < \epsilon$. For $f \in \mathcal{R}^1[a, b]$ there are polynomials P with $\|f - P\|_1 < \epsilon$.

Proof. As in Lemma 7.12, with $\int_a^b |f - s| = \int_a^b (f - s) \le U - L$ for the lower step function. For the continuous approximation of $\chi_{[a,b]}$ the error in $\|\cdot\|_1$ is at most η. Polynomials: $\|g - P\|_1 \le (b - a)\|g - P\|_{\sup}$. ∎