Chapter 4. Inference in Simple Regression

Sampling distributions, confidence intervals, prediction, inference on correlation, residuals

Contents

4.1 Sampling distributions of the estimators **2**

4.2 Estimating the mean response and predicting **3**

4.3 Inference on the correlation **3**

4.4 Residuals **4**

Throughout, $y_i = \beta_0 + \beta_1 x_i + \epsilon_i$ with $\epsilon_i \overset{\text{iid}}{\sim} N(0, \sigma^2)$ and fixed x_i with $S_{xx} > 0$.

4.1 Sampling distributions of the estimators

Both estimators are linear in y:

$$\hat\beta_1 = \frac{S_{xy}}{S_{xx}} = \sum_i k_i y_i, \qquad k_i = \frac{x_i - \bar x}{S_{xx}}, \qquad \sum k_i = 0, \quad \sum k_i x_i = 1, \quad \sum k_i^2 = \frac{1}{S_{xx}},$$

(using $S_{xy} = \sum (x_i - \bar x) y_i$), and $\hat\beta_0 = \bar y - \hat\beta_1 \bar x$.

Theorem 4.1

(a) $\hat\beta_1 \sim N(\beta_1, \sigma^2 / S_{xx})$.

(b) $\operatorname{Cov}(\bar y, \hat\beta_1) = 0$.

(c) $\hat\beta_0 \sim N\Big(\beta_0,\ \sigma^2\big(\frac{1}{n} + \frac{\bar x^2}{S_{xx}}\big)\Big)$ and $\operatorname{Cov}(\hat\beta_0, \hat\beta_1) = -\dfrac{\bar x \sigma^2}{S_{xx}}$.

Proof. (a) $\hat\beta_1$ is a linear combination of independent normals, with mean $\sum k_i(\beta_0 + \beta_1 x_i) = \beta_1$ and variance $\sigma^2 \sum k_i^2 = \sigma^2 / S_{xx}$.

(b) $\operatorname{Cov}(\bar y, \hat\beta_1) = \sum_i \frac{1}{n} k_i \sigma^2 = 0$.

(c) $E\hat\beta_0 = (\beta_0 + \beta_1 \bar x) - \bar x \beta_1 = \beta_0$ and, by (b),

$$\operatorname{Var} \hat\beta_0 = \operatorname{Var} \bar y + \bar x^2 \operatorname{Var} \hat\beta_1 = \sigma^2\Big(\frac{1}{n} + \frac{\bar x^2}{S_{xx}}\Big), \qquad \operatorname{Cov}(\hat\beta_0, \hat\beta_1) = -\bar x \operatorname{Var} \hat\beta_1.$$

■

Theorem 4.2

Let $SSE = \sum e_i^2$ and $MSE = SSE/(n-2)$. Then $SSE/\sigma^2 \sim \chi^2(n-2)$, independent of $(\hat\beta_0, \hat\beta_1)$. In particular $E(MSE) = \sigma^2$.

This is the case $p = 1$ of Theorem 6.4; a direct proof uses an orthogonal transformation whose first two rows span $\mathbf{1}$ and $\mathbf{x} - \bar x \mathbf{1}$, exactly as for the normal sample variance. Consequently

$$\frac{\hat\beta_1 - \beta_1}{\sqrt{MSE / S_{xx}}} \sim t(n-2), \qquad \frac{\hat\beta_0 - \beta_0}{\sqrt{MSE(1/n + \bar x^2 / S_{xx})}} \sim t(n-2).$$

Corollary 4.3 (Intervals and tests for the coefficients)

$100(1-\alpha)\%$ confidence intervals are

$$\beta_1 :\ \hat\beta_1 \pm t_{\alpha/2}(n-2)\sqrt{\frac{MSE}{S_{xx}}}, \qquad \beta_0 :\ \hat\beta_0 \pm t_{\alpha/2}(n-2)\sqrt{MSE\Big(\frac{1}{n} + \frac{\bar x^2}{S_{xx}}\Big)}.$$

The test of $H_0 : \beta_1 = 0$ rejects when $|t_0| > t_{\alpha/2}(n-2)$, $t_0 = \hat\beta_1 / \sqrt{MSE/S_{xx}}$. If σ^2 is known, replace MSE by σ^2 and t by z.

Theorem 4.4 (ANOVA F test)

$E(SSR) = \sigma^2 + \beta_1^2 S_{xx}$ and $E(SSE) = (n-2)\sigma^2$. The statistic $F_0 = MSR/MSE$, $MSR = SSR/1$, satisfies $F_0 = t_0^2$, so the F test with $F_\alpha(1, n-2) = t_{\alpha/2}(n-2)^2$ is the two-sided t test of $\beta_1 = 0$.

Proof. $E(SSR) = S_{xx} E\hat\beta_1^2 = S_{xx}(\sigma^2/S_{xx} + \beta_1^2)$. Also $SSR/MSE = \hat\beta_1^2 S_{xx}/MSE = t_0^2$. ■

So $E(MSR)/E(MSE) = 1 + \beta_1^2 S_{xx}/\sigma^2$, which is 1 exactly when $\beta_1 = 0$; large F_0 is evidence against H_0.

4.2 Estimating the mean response and predicting

At a point x, the mean response $\mu_{y\cdot x} = \beta_0 + \beta_1 x$ is estimated by $\hat y = \hat\beta_0 + \hat\beta_1 x = \bar y + \hat\beta_1(x - \bar x)$.

Theorem 4.5

(a) $\hat y \sim N\Big(\mu_{y\cdot x},\ \sigma^2\Big(\dfrac{1}{n} + \dfrac{(x-\bar x)^2}{S_{xx}}\Big)\Big)$, and a $100(1-\alpha)\%$ confidence interval for $\mu_{y\cdot x}$ is

$$\hat y \pm t_{\alpha/2}(n-2)\sqrt{MSE\Big(\frac{1}{n} + \frac{(x-\bar x)^2}{S_{xx}}\Big)}.$$

(b) For a new observation $y_s = \beta_0 + \beta_1 x + \epsilon_s$, independent of the sample, $y_s - \hat y \sim N\big(0, \sigma^2(1 + \frac{1}{n} + \frac{(x-\bar x)^2}{S_{xx}})\big)$, and a $100(1-\alpha)\%$ prediction interval is

$$\hat y \pm t_{\alpha/2}(n-2)\sqrt{MSE\Big(1 + \frac{1}{n} + \frac{(x-\bar x)^2}{S_{xx}}\Big)}.$$

Proof. (a) By Theorem 4.1(b), $\operatorname{Var}\hat y = \operatorname{Var}\bar y + (x-\bar x)^2 \operatorname{Var}\hat\beta_1$. Studentize with MSE, which is independent of $\hat y$.

(b) y_s is independent of $\hat y$, so the variances add. ■

Both intervals are narrowest at $x = \bar x$ and widen quadratically away from it; extrapolation is costly. The prediction interval does not shrink to zero as $n \to \infty$, because of the irreducible variance σ^2 of a new observation.

Example 4.1

For the advertising data of Example 3.1 ($n = 6$, $\bar x = 5.667$, $S_{xx} = 45.33$, $MSE = 1.92/4 = 0.48$, $\hat y = 2.37 + 1.346x$), at $x = 7$: $\hat y = 11.79$, $\frac{1}{6} + \frac{(1.333)^2}{45.33} = 0.206$, and $t_{0.025}(4) = 2.776$. The 95% confidence interval for the mean is $11.79 \pm 2.776\sqrt{0.48 \cdot 0.206} = 11.79 \pm 0.87$, and the prediction interval is $11.79 \pm 2.776\sqrt{0.48 \cdot 1.206} = 11.79 \pm 2.11$.

4.3 Inference on the correlation

When (X, Y) is bivariate normal with correlation ρ, the conditional mean is linear with slope $\beta_1 = \rho\sigma_Y/\sigma_X$, so $\rho = 0 \iff \beta_1 = 0$.

Theorem 4.6

(a) The t statistic for $\beta_1 = 0$ equals $t_0 = \dfrac{r\sqrt{n-2}}{\sqrt{1-r^2}}$, which is $t(n-2)$ under $\rho = 0$.

(b) (Fisher) $Z = \frac{1}{2}\log\frac{1+r}{1-r}$ is approximately $N\big(\frac{1}{2}\log\frac{1+\rho}{1-\rho}, \frac{1}{n-3}\big)$ for moderate n (say $n \geq 25$).

Proof. (a) $SSE = SST(1-r^2)$ and $\hat\beta_1^2 S_{xx} = SSR = r^2 SST$, so

$$t_0^2 = \frac{\hat\beta_1^2 S_{xx}}{SSE/(n-2)} = \frac{(n-2)r^2}{1-r^2}.$$

The null distribution is conditionally $t(n-2)$ given the x_i, hence unconditionally as well.

(b) is the variance-stabilizing transformation of the asymptotic law $\sqrt{n}(r-\rho) \to N(0, (1-\rho^2)^2)$; the correction $n-3$ improves the approximation. ■

Part (b) gives tests of $H_0 : \rho = \rho_0 \neq 0$ and confidence intervals: $\tanh\big(Z \pm z_{\alpha/2}/\sqrt{n-3}\big)$ covers ρ with probability about $1-\alpha$.

4.4 Residuals

Residuals are our window on the errors. In vector form $\mathbf{e} = (I - H)\mathbf{y}$ with H the hat matrix (Chapter 6), and:

Theorem 4.7

$Ee_i = 0$ and

$$\operatorname{Var}(e_i) = \sigma^2\Big(1 - \frac{1}{n} - \frac{(x_i-\bar x)^2}{S_{xx}}\Big), \qquad \operatorname{Cov}(e_i, e_j) = -\sigma^2\Big(\frac{1}{n} + \frac{(x_i-\bar x)(x_j-\bar x)}{S_{xx}}\Big) \ (i \neq j).$$

Proof. $e_i = y_i - \bar y - \hat\beta_1(x_i - \bar x)$. Using $\operatorname{Var}\bar y = \sigma^2/n$, $\operatorname{Var}\hat\beta_1 = \sigma^2/S_{xx}$, $\operatorname{Cov}(y_i, \bar y) = \sigma^2/n$, $\operatorname{Cov}(y_i, \hat\beta_1) = \sigma^2 k_i$ and $\operatorname{Cov}(\bar y, \hat\beta_1) = 0$,

$$\operatorname{Var} e_i = \sigma^2\Big(1 + \frac{1}{n} + \frac{(x_i-\bar x)^2}{S_{xx}} - \frac{2}{n} - 2\frac{(x_i-\bar x)^2}{S_{xx}}\Big),$$

which simplifies to the claim; the covariance is computed in the same way. ■

Thus residuals are neither independent nor equally variable; points far from $\bar x$ have smaller residual variance (Chapter 10). Since $\sum e_i = 0$ and $\sum x_i e_i = 0$, residual plots against x_i or $\hat y_i$ have no linear trend by construction; what we look for is *non-random* structure:

- a curved band suggests a missing quadratic term or a transformation;
- a funnel suggests nonconstant variance (weighted least squares or a variance-stabilizing transformation of y);
- a trend in time order suggests correlated errors (Chapter 12).