Chapter 12. Planar Graphs

Do Not Cross

Contents

12.1 Euler's theorem for planar graphs **2**

12.2 Polyhedra **3**

12.3 Colouring maps **5**

12.1 Euler's theorem for planar graphs

Motivation. Can three houses be joined to three wells by non-crossing roads? Drawing a graph without crossings creates a new parameter, the number of faces, and Euler's formula $V+F=E+2$ ties it to V and E. Counting edges around faces then rules out $K_{3,3}$ and K_5.

Definition 12.1 (Planar graph)

A graph is **planar** if it can be drawn in the plane with no two edges crossing. The edges of such a drawing cut the plane into regions, the **faces**; one of them, the outer face, is unbounded.

Theorem 12.2 (Euler's theorem on planar graphs)

A connected planar graph with V vertices, E edges and F faces satisfies $V+F=E+2$.

Proof. Induction on E. For $E=1$: either one edge ($V=2$, $F=1$) or one loop ($V=1$, $F=2$). Let $E \geq 2$.

Step 1 (an edge on a cycle). If removing some edge e leaves G' connected, then e lies on a cycle, so the two sides of e belong to different faces, which merge in G'. G' has V vertices, $E-1$ edges and $F-1$ faces, so $V+F-1=E-1+2$ by induction.

Step 2 (a tree). Otherwise G is a tree, so $V=E+1$ (Theorem 10.4) and $F=1$. ■


Example 12.3

$K_{3,3}$ is not planar, so the three houses, three wells problem has no solution.
Solution. With $V=6$, $E=9$, a planar drawing would have $F=5$ faces. $K_{3,3}$ is bipartite and simple, so it has no odd cycles and no 2-cycles, and every face is bounded by at least 4 edges. Each edge borders at most two faces, so $4F \leq 2E$, i.e. $20 \leq 18$, a contradiction.

$K_{3,3} \subseteq K_6$, so K_6 is not planar either. K_4 is planar (draw a square with one diagonal inside and the other outside).

Example 12.4

K_5 is not planar.
Solution. With $V=5$, $E=10$, a drawing would have $F=7$ faces, each bounded by at least 3 edges (the graph is simple). Then $3F \leq 2E$, i.e. $21 \leq 20$, a contradiction.



K_5 $K_{3,3}$ K_4, drawn without crossings

Figure. *The two minimal non-planar graphs and a crossing-free drawing of K_4.*

Deleting a vertex of degree 2 and merging its two edges, or subdividing an edge by a new vertex, does not change planarity. Graphs related by a sequence of these operations are **edge-equivalent** (homeomorphic).

Theorem 12.5 (Kuratowski)

A graph is not planar iff it contains a subgraph that is edge-equivalent to K_5 or $K_{3,3}$.

The "if" direction follows from Examples 12.3 and 12.4. The converse is cited without proof, as in the book.

12.2 Polyhedra

Motivation. Regular polygons exist with any number of sides, but there are only five regular polyhedra. The reason is purely combinatorial: Euler's formula, together with "every face has ≥ 3 edges and every vertex has degree ≥ 3", leaves room for only five pairs (d, ℓ).

For a convex polyhedron (identified with its 1-skeleton) every face has at least 3 edges and every vertex lies on at least 3 edges. A polyhedron is **regular** (combinatorially) if every face is an ℓ-cycle and every vertex has degree d.

Theorem 12.6

A convex polyhedron with V vertices, F faces and E edges satisfies $V + F = E + 2$.

Proof (angle counting). Project P onto a plane p perpendicular to no face. Every face projects to a convex polygon, and the outer boundary B of the image is a convex v_1-gon; let v_2 vertices project inside it, so $v_1 + v_2 = V$. Each point inside B is covered by exactly one "front" and one "back" face. Let S be the sum of all angles of all projected faces.

Step 1 (by vertices). At a vertex projecting inside B, the image faces meeting there fill a full neighbourhood, so their angles sum to $360°$; this gives $360v_2$. At a vertex on B, the front faces and the back faces each fill the interior angle of B, so the boundary vertices contribute twice the angle sum $(v_1 - 2)180°$ of B. Hence

$$S = (v_1 - 2)360 + 360v_2 = (V - 2)360. \tag{12.1}$$

Step 2 (by faces). Let $f_1, \ldots, f_F$ be the numbers of edges of the faces. Each edge lies on two faces, so

$$\sum_{i=1}^{F} f_i = 2E, \tag{12.2}$$

and a convex k-gon has angle sum $(k - 2)180°$, so

$$S = \sum_{i=1}^{F} (f_i - 2)180 = 180 \sum_{i=1}^{F} f_i - 360F = 360(E - F). \tag{12.3}$$

Comparing (12.1) and (12.3) gives $V - 2 = E - F$. ∎

Corollary 12.7

In a convex polyhedron, $3F \leq 2E$.

Proof. $f_i \geq 3$ in (12.2). ∎

Proposition 12.8

In a convex polyhedron, $3V \leq 2E$.

Proof. Let $c_1, \ldots, c_V$ be the vertex degrees. Each edge has two ends, so

$$\sum_{i=1}^{V} c_i = 2E, \tag{12.4}$$

and $c_i \geq 3$. ■

Lemma 12.9

In a convex polyhedron, $E \leq 3V - 6$ and $E \leq 3F - 6$.

Proof. By Corollary 12.7, $E + 2 = V + F \leq V + \frac{2E}{3}$, so $\frac{E}{3} \leq V - 2$. By Proposition 12.8, $E + 2 \leq F + \frac{2E}{3}$, so $\frac{E}{3} \leq F - 2$. ■

Remark

Remark[$E \leq 3V - 6$ for every simple planar graph [Supplement]] Let G be simple and planar with $V \geq 3$; we may assume it connected (adding edges between components keeps it planar). If G is a tree, $E = V - 1 \leq 3V - 6$. Otherwise every face is bounded by a closed walk of length at least 3 (no loops or multiple edges) and each edge borders at most two faces (counted twice if it borders one face on both sides), so $3F \leq 2E$. Euler's formula then gives $E \leq 3V - 6$ as above. Consequently every simple planar graph has a vertex of degree at most 5: otherwise $2E \geq 6V$.

Lemma 12.10

Every convex polyhedron has a face with at most five edges.

Proof. By (12.2) and Lemma 12.9,

$$\sum_{i=1}^{F} f_i = 2E \leq 6F - 12 < 6F, \tag{12.5}$$

so not all $f_i \geq 6$. ■

Lemma 12.11

Every convex polyhedron has a vertex of degree at most five.

Proof. By (12.4) and Lemma 12.9,

$$\sum_{i=1}^{V} c_i = 2E \leq 6V - 12 < 6V. \tag{12.6}$$

■

The symmetry between V and F has a structural reason. Given a planar G, put a vertex inside each face and, for every edge e of G, join the vertices of the two faces on either side of e by an edge crossing e. The result G^* is planar with F vertices, E edges and V faces.

Definition 12.12 (Dual graph)

G^* is the **dual graph** of the plane graph G.

A statement about (V, E) of a polyhedron P is a statement about (F, E) of P^*. The dual of a regular polyhedron of type (d, ℓ) is regular of type (ℓ, d).

Classification. In a regular polyhedron, Lemmas 12.10 and 12.11 give $d, \ell \in \{3, 4, 5\}$, and by (12.2)

$$E = \frac{F\ell}{2}. \tag{12.7}$$

(A) $d = 3$: $3V = 2E$, so Euler gives $3F = E + 6$, and with (12.7), $(6 - \ell)F = 12$. Then $\ell = 3, 4, 5$ give $(V, E, F) = (4, 6, 4)$ (tetrahedron), $(8, 12, 6)$ (cube), $(20, 30, 12)$ (dodecahedron).

(B) $d = 4$: $4V = 2E$, so $E = 2F - 4$ and $(4 - \ell)F = 8$; only $\ell = 3$ works: $(6, 12, 8)$, the octahedron (dual of the cube).

(C) $d = 5$: $5V = 2E$, so $3E = 5F - 10$ and $(10 - 3\ell)F = 20$; only $\ell = 3$ works: $(12, 30, 20)$, the icosahedron (dual of the dodecahedron).

Theorem 12.13

There are exactly five regular polyhedra: the tetrahedron, the cube, the dodecahedron, the octahedron and the icosahedron.

Proof. The case analysis (A)–(C) leaves exactly these five parameter sets, and each is realised by the named solid. ■

	d	ℓ	V	E	F
Tetrahedron	3	3	4	6	4
Cube	3	4	8	12	6
Octahedron	4	3	6	12	8
Dodecahedron	3	5	20	30	12
Icosahedron	5	3	12	30	20

12.3 Colouring maps

Motivation. Colouring the countries of a map so that neighbours differ is, by duality, properly colouring the vertices of a planar graph. A vertex of degree ≤ 5 gives six colours at once; Kempe's chain-swapping trick brings this down to five. Four colours suffice too, but every known proof uses a computer.

Faces sharing an edge must get different colours; this is the same as properly colouring the vertices of G^*. K_4 is planar, so at least four colours are needed.

Proposition 12.14

Every planar graph can be properly vertex-coloured with six colours.

Proof. Induction on V; loops are excluded and multiple edges do not matter, so let G be simple. By the Remark after Lemma 12.9, G has a vertex A of degree at most 5. Colour $G - A$ with six colours by induction, and give A a colour not used by its at most five neighbours. ■

Theorem 12.15 (Five colour theorem)

Every planar graph can be properly vertex-coloured with five colours.

Proof. Induction on V for simple G. Take A of degree ≤ 5 and colour $G' = G - A$ properly with five colours. If the neighbours of A use at most four colours, colour A with the fifth. Otherwise A has neighbours $y_1, \dots, y_5$ in clockwise order, coloured $1, \dots, 5$.

Step 1 (Kempe chains). Let H_{13} be the subgraph of G' induced by the vertices coloured 1 or 3, and K the component of H_{13} containing y_1. Swapping colours 1 and 3 on K keeps the colouring proper. If $y_3 \notin K$, after the swap y_1 and y_3 both have colour 3, colour 1 is free at A, and we are done.

Step 2 (both chains cannot exist). So assume $y_3 \in K$: there is a path $y_1 \to y_3$ coloured alternately $1, 3$. Together with A it forms a cycle C through A that separates y_2 from y_4 in the plane (the neighbours occur around A in the order y_1, y_2, y_3, y_4). By the same argument with colours $2, 4$, we are done unless there is a path $y_2 \to y_4$ coloured $2, 4$. This path would have to cross C at a vertex, but its vertices are coloured $2, 4$ and those of $C - A$ are coloured $1, 3$. So this case cannot occur. ■




Figure. *Kempe chains in the five colour theorem: a 1–3 path from y_1 to y_3 and a 2–4 path from y_2 to y_4 would have to cross at a vertex.*

Remark

Remark[The four colour theorem] Four colours always suffice for maps with contiguous countries. Appel and Haken proved this in 1976 by a computer check of 1936 configurations, later reduced to 633 by Robertson, Sanders, Seymour and Thomas (1996). A machine-verified proof was given by Gonthier and Werner (2005).