Chapter 5. The Tychonoff Theorem

Compactness of products, Stone–Čech compactification

Contents

5.1 The Tychonoff theorem **2**

5.2 The Stone–Čech compactification **2**

5.1 The Tychonoff theorem

We prove compactness via the finite intersection property (Theorem 3.23): a space is compact iff every collection $\mathcal{A}$ of subsets with the finite intersection property (FIP) satisfies $\bigcap_{A\in\mathcal{A}} \overline{A} \neq \emptyset$.

Lemma 5.1 (37.1)

Let $\mathcal{A}$ be a collection of subsets of X with the FIP. Then there is a collection $\mathcal{D} \supset \mathcal{A}$ with the FIP such that no collection properly containing $\mathcal{D}$ has the FIP ($\mathcal{D}$ is *maximal* with respect to the FIP).

Proof. Let $\mathbb{A}$ be the set of all collections $\mathcal{B} \supset \mathcal{A}$ of subsets of X with the FIP, strictly partially ordered by proper inclusion. If $\mathbb{B} \subset \mathbb{A}$ is a chain, $\mathcal{C} = \bigcup_{\mathcal{B}\in\mathbb{B}} \mathcal{B}$ has the FIP: finitely many $C_1, \ldots, C_n \in \mathcal{C}$ lie in $\mathcal{B}_1, \ldots, \mathcal{B}_n$, all contained in the largest $\mathcal{B}_k$ (chain), which has the FIP. So $\mathcal{C}$ is an upper bound, and Zorn's lemma (Theorem 1.29) gives a maximal element. ■

Lemma 5.2 (37.2)

Let $\mathcal{D}$ be maximal with respect to the FIP. (a) Every finite intersection of elements of $\mathcal{D}$ is in $\mathcal{D}$. (b) If $A \subset X$ intersects every element of $\mathcal{D}$, then $A \in \mathcal{D}$.

Proof. (a) Let B be a finite intersection of elements of $\mathcal{D}$; $\mathcal{E} = \mathcal{D} \cup \{B\}$ has the FIP (a finite intersection of members of $\mathcal{E}$ is a finite intersection of members of $\mathcal{D}$), so $\mathcal{E} = \mathcal{D}$ by maximality. (b) $\mathcal{E} = \mathcal{D} \cup \{A\}$: a finite intersection $A \cap D_1 \cap \cdots \cap D_n$ is nonempty since $D_1 \cap \cdots \cap D_n \in \mathcal{D}$ by (a) and A meets it. So $A \in \mathcal{D}$. ■

Theorem 5.3 (37.3, Tychonoff theorem)

An arbitrary product of compact spaces is compact in the product topology.

Proof. Let $X = \prod_{\alpha\in J} X_\alpha$ and $\mathcal{A}$ a collection of subsets of X with the FIP; extend it to a maximal $\mathcal{D}$ (Lemma 5.1). It suffices to find $x \in \bigcap_{D\in\mathcal{D}} \overline{D}$. For each α, $\{\pi_\alpha(D) : D \in \mathcal{D}\}$ has the FIP in the compact X_α, so choose $x_\alpha \in \bigcap_D \overline{\pi_\alpha(D)}$, and let $x = (x_\alpha)$. Step 1: if U_β is a neighbourhood of x_β, then $\pi_\beta^{-1}(U_\beta)$ meets every $D \in \mathcal{D}$ (since U_β meets $\pi_\beta(D)$, as $x_\beta \in \overline{\pi_\beta(D)}$), so $\pi_\beta^{-1}(U_\beta) \in \mathcal{D}$ by Lemma 5.2(b). Step 2: every basis element containing x is a finite intersection of such subbasis elements, hence lies in $\mathcal{D}$ (Lemma 5.2(a)), hence meets every element of $\mathcal{D}$ (FIP). So $x \in \overline{D}$ for all $D \in \mathcal{D}$. ■

The theorem fails for the box topology ($[0,1]^\omega$ in the box topology is not compact), and it is equivalent to the axiom of choice.

Example 5.1

$[0,1]^\omega$ is a compact metrizable space; every regular second countable space imbeds in it (proof of Theorem 4.12). $[0,1]^J$ for uncountable J is compact Hausdorff but not metrizable: it is not even first countable (a countable family of basis neighbourhoods of a point restricts only countably many coordinates).

5.2 The Stone–Čech compactification

A **compactification** of X is a compact Hausdorff space Y containing X as a subspace with $\overline{X} = Y$; two are **equivalent** if there is a homeomorphism between them that is the identity on X. If X

has a compactification, it is completely regular (a subspace of a compact Hausdorff, hence normal, space).

Lemma 5.4 (38.1)

If $h : X \to Z$ is an imbedding in a compact Hausdorff space Z, there is a compactification Y of X and an imbedding $H : Y \to Z$ equal to h on X; Y is unique up to equivalence (Y is the compactification **induced** by h).

Proof. Let $X_0 = h(X)$ and $Y_0 = \overline{X_0}$, a compact Hausdorff space. Form $Y = X \sqcup (Y_0 - X_0)$ with the bijection $H : Y \to Y_0$ (h on X, identity elsewhere) and give Y the topology making H a homeomorphism. Then X is a subspace of Y, dense since X_0 is dense in Y_0. ■

Theorem 5.5 (38.2)

Every completely regular space X has a compactification Y such that every bounded continuous $f : X \to \mathbb{R}$ extends uniquely to a continuous $Y \to \mathbb{R}$.

Proof. Let $\{f_\alpha\}_{\alpha \in J}$ be all bounded continuous real functions on X, with $f_\alpha(X) \subset I_\alpha = [\inf f_\alpha, \sup f_\alpha]$. $h = (f_\alpha) : X \to Z = \prod I_\alpha$ is an imbedding (Theorem 4.13, using complete regularity), and Z is compact Hausdorff (Tychonoff). Let Y be the induced compactification with $H : Y \to Z$. For $f = f_\beta$, $\pi_\beta \circ H$ is a continuous extension. Uniqueness follows from the next lemma. ■

Lemma 5.6 (38.3)

If $A \subset X$ and $f : A \to Z$ is continuous with Z Hausdorff, there is at most one continuous extension $g : \overline{A} \to Z$.

Proof. If $g \neq g'$ at x, separate $g(x), g'(x)$ by disjoint U, U'; the neighbourhood $g^{-1}(U) \cap g'^{-1}(U')$ of x contains a point $y \in A$, but $g(y) = g'(y) = f(y)$ lies in $U \cap U'$. ■

Theorem 5.7 (38.4)

If Y is a compactification of the completely regular X with the extension property of Theorem 5.5, every continuous map f of X into a compact Hausdorff space C extends uniquely to a continuous $g : Y \to C$.

Proof. C is completely regular, so imbeds in some $[0,1]^J$ (Corollary 4.14); regard $C \subset [0,1]^J$. Each coordinate $\pi_\alpha \circ f$ extends to $g_\alpha : Y \to [0,1]$; $g = (g_\alpha) : Y \to [0,1]^J$ is continuous. Since $g(X) = f(X) \subset C$ and X is dense, $g(Y) \subset \overline{g(X)} \subset C$ (C is closed). Uniqueness: Lemma 5.6. ■

Theorem 5.8 (38.5)

Two compactifications Y_1, Y_2 of X with the extension property of Theorem 5.5 are equivalent.

Proof. Extend the inclusion $j_2 : X \to Y_2$ to $f_2 : Y_1 \to Y_2$ and j_1 to $f_1 : Y_2 \to Y_1$ (Theorem 5.7). $f_1 \circ f_2 : Y_1 \to Y_1$ extends the identity of X, as does id_{Y_1}; by Lemma 5.6, $f_1 \circ f_2 = \mathrm{id}$. Similarly $f_2 \circ f_1 = \mathrm{id}$. ■

This compactification is the **Stone–Čech compactification** $\beta(X)$. It is characterized by the universal property of Theorem 5.7.

Example 5.2

(a) The one-point compactification of $(0,1)$ is the circle and $[0,1]$ is a two-point compactification; neither is $\beta(0,1)$, since $\sin(1/x)$ extends to $\beta(0,1)$.
(b) $\beta(\mathbb{Z}_+)$ is not sequentially compact: $\mathbb{Z}_+$ has no convergent subsequence in $\beta\mathbb{Z}_+$ (separate even and odd subsequences by extending functions on $\mathbb{Z}_+$).
(c) $\beta(S_\Omega) = \overline{S}_\Omega$, because every continuous real function on S_Ω is eventually constant.