# 7. An Algorithmic Approach to Identification

A consolidated, mathematical lecture companion

English notes based on the lecture slides *An Algorithmic Approach to Identification* (Prof. Sanghack Lee, GSDS SNU)

The central question is how to decide identifiability of $P(\mathbf{y} \mid \mathrm{do}(\mathbf{x}))$ mechanically, without searching for a clever sequence of do-calculus steps.

**Roadmap.** (1) Define a factorisation of the distributions generated by an SCM based on its causal diagram (c-components and c-factors). (2) Establish operations that identify particular factors from a distribution. (3) Express the target effect in terms of factors and identify each one independently. These notes consolidate the slides, verify every example by explicit sums, and state the resulting algorithm with its completeness theorem. Physical PDF page numbers are listed in Section 5.

## Contents

**1 Factorising observational distributions** **2**

1.1 Confounded components . . . 2
1.2 Markovian case . . . 2
1.3 Semi-Markovian case . . . 2
1.4 C-factors . . . 3
1.5 C-factors are causal effects . . . 3

**2 C-factor algebra** **3**

2.1 Marginalising variables in c-factors . . . 3
2.2 C-component factorisation . . . 4
2.3 Visualisation . . . 5

**3 Causal queries in terms of c-factors: examples** **5**

3.1 Back-door . . . 5
3.2 Front-door . . . 5
3.3 Napkin . . . 5
3.4 Overview of the three derivations . . . 6

**4 A general identification algorithm** **6**

4.1 Reducing the query to c-factors . . . 6
4.2 The Identify procedure . . . 7

**5 Source coverage** **7**


# 1 Factorising observational distributions

**Purpose.** Group variables that share latent confounders and show that the joint distribution factorises over these groups.

## 1.1 Confounded components

**Definition 1.1** (C-component)**.** When $V_i$ and $V_j$ share a common unobservable parent $U$, their mechanisms are tied together through $U$, and $V_i$, $V_j$ are in the same *confounded component* (c-component). The relation is closed under chaining: two variables are in the same c-component iff they are connected by a path of bidirected edges.

Running example: $V_1 \to V_2 \to V_3 \to V_4 \to V_5$ with $U_1$: $V_1 \leftrightarrow V_3$, $U_2$: $V_3 \leftrightarrow V_5$, $U_3$: $V_2 \leftrightarrow V_4$.




$V_1$ is in the same c-component as $V_3$, $V_3$ as $V_5$, hence $V_1$ as $V_5$; $V_2$ is with $V_4$. To see this easily, look at the graph induced by the bidirected edges only. The c-component relation is reflexive, symmetric and transitive, so it *partitions* the observed variables: here $\{V_1, V_3, V_5\}$ and $\{V_2, V_4\}$.

## 1.2 Markovian case

$$P(\mathbf{v}) = \sum_{\mathbf{u}} P(\mathbf{u}) \prod_{V_i \in \mathbf{V}} P(v_i \mid v_1, \ldots, v_{i-1}, \mathbf{u}) = \sum_{\mathbf{u}} P(\mathbf{u}) \prod_{V_i \in \mathbf{V}} P(v_i \mid \mathbf{pa}_i, \mathbf{u}_i).$$

For $Z \to X$, $Z \to Y$, $X \to Y$:

$$\begin{aligned} P(z, x, y) &= \sum_{u_z} P(z \mid u_z) P(u_z) \sum_{u_x} P(x \mid z, u_x) P(u_x) \sum_{u_y} P(y \mid x, z, u_y) P(u_y) \\ &= P(z) P(x \mid z) P(y \mid x, z). \end{aligned}$$

The factors $P(v_i \mid \mathbf{pa}_i)$ are "canonical", and every one of them is computable from $P(\mathbf{v})$:

$$P(v_i \mid \mathbf{pa}_i) = \frac{\sum_{\mathbf{v} \setminus (\{v_i\} \cup \mathbf{pa}_i)} P(\mathbf{v})}{\sum_{\mathbf{v} \setminus \mathbf{pa}_i} P(\mathbf{v})},$$

$$\text{e.g. } P(z) = \sum_{x,y} P(\mathbf{v}),\ P(x \mid z) = \frac{\sum_y P(\mathbf{v})}{\sum_{x,y} P(\mathbf{v})},\ P(y \mid x, z) = \frac{P(\mathbf{v})}{\sum_y P(\mathbf{v})}.$$

## 1.3 Semi-Markovian case

With only $U_1$ ($V_1 \leftrightarrow V_3$) and $U_2$ ($V_3 \leftrightarrow V_5$):

$$P(\mathbf{v}) = P(v_2 \mid v_1) P(v_4 \mid v_3) \sum_{u_1, u_2} P(u_1, u_2) P(v_1 \mid u_1) P(v_3 \mid v_2, u_1, u_2) P(v_5 \mid v_4, u_2).$$

Adding $U_3$ ($V_2 \leftrightarrow V_4$):

$$\begin{aligned} P(\mathbf{v}) = &\Big[ \sum_{u_3} P(v_2 \mid v_1, u_3) P(v_4 \mid v_3, u_3) P(u_3) \Big] \\ &\cdot \Big[ \sum_{u_1, u_2} P(u_1, u_2) P(v_1 \mid u_1) P(v_3 \mid v_2, u_1, u_2) P(v_5 \mid v_4, u_2) \Big]. \end{aligned}$$

The sums separate along c-components because different c-components share no latent variable.

### 1.4 C-factors

These factors are long to write in terms of $P(\mathbf{v}, \mathbf{u})$, but their structure follows from the diagram. Define for $C \subseteq \mathbf{V}$

$$Q[C](\mathbf{c}, \mathbf{pa}_C) = \sum_{\mathbf{u}(C)} P(\mathbf{u}(C)) \prod_{V_i \in C} P(v_i \mid \mathbf{pa}_i, \mathbf{u}_i),$$
$$\mathbf{U}(C) = \bigcup_{V_i \in C} \mathbf{U}_i.$$

Then

$$P(\mathbf{v}) = Q[V_2, V_4](v_2, v_4, v_1, v_3) \cdot Q[V_1, V_3, V_5](v_1, v_3, v_5, v_2, v_4),$$

abbreviated $P(\mathbf{v}) = Q[V_1, V_3, V_5]\, Q[V_2, V_4]$, without naming $\mathbf{U}$ at all. Note $Q[\mathbf{V}] = P(\mathbf{v})$, and by convention $Q[\emptyset] = 1$.

### 1.5 C-factors are causal effects

For $C \subseteq \mathbf{V}$, consider the effect of all other variables on $C$. By the truncated factorisation,

$$P(\mathbf{c} \mid \mathrm{do}(\mathbf{v} \setminus \mathbf{c})) = \sum_{\mathbf{u}} P(\mathbf{u}) \prod_{V_i \in C} P(v_i \mid \mathbf{pa}_i, \mathbf{u}_i) = \sum_{\mathbf{u}(C)} \prod_{V_i \in C} P(v_i \mid \mathbf{pa}_i, \mathbf{u}_i) P(\mathbf{u}(C)) = Q[C],$$

since latent variables that are not parents of $C$ sum out. *C-factors are interventional distributions*; this is the key link between factorisation and identification. For example $Q[V_1, V_3, V_5](v_1, v_3, v_5, v_2, v_4) = \sum_{u_1, u_2} P(u_1, u_2) P(v_1 \mid u_1) P(v_3 \mid v_2, u_1, u_2) P(v_5 \mid v_4, u_2)$ is the distribution of $(V_1, V_3, V_5)$ when $V_2 = v_2$ and $V_4 = v_4$ are fixed by intervention.

## 2 C-factor algebra

**Purpose.** Develop the two operations, marginalising to ancestral sets and factorising into c-components, that drive identification.

### 2.1 Marginalising variables in c-factors

C-factors behave partly like probabilities. In $Q[V_1, V_3, V_5]$ each of $V_1, V_3, V_5$ appears in a single term (none is a parent of another member), so summing over $v_3$ gives

$$\sum_{v_3} Q[V_1, V_3, V_5] = \sum_{u_1, u_2} P(u_1, u_2) P(v_1 \mid u_1) \Big( \sum_{v_3} P(v_3 \mid v_2, u_1, u_2) \Big) P(v_5 \mid v_4, u_2) = Q[V_1, V_5].$$

In $Q[V_1, V_2, V_3]$, by contrast, $V_1$ is a parent of $V_2$. Summing over $v_1$:

$$\sum_{v_1} Q[V_1, V_2, V_3] = \sum_{\mathbf{u}} P(\mathbf{u}) P(v_3 \mid v_2, u_1, u_2) \sum_{v_1} P(v_1 \mid u_1) P(v_2 \mid v_1, u_3),$$

whereas

$$Q[V_2, V_3](v_2, v_3, v_1) = \sum_{\mathbf{u}} P(\mathbf{u}) P(v_3 \mid v_2, u_1, u_2) P(v_2 \mid v_1, u_3)$$

keeps $v_1$ as a fixed argument. The coupling of $v_1$ with $u_1$ (shared with $V_3$) inside the sum prevents the reduction, so $\sum_{v_1} Q[V_1, V_2, V_3] \neq Q[V_2, V_3]$ in general: one can only sum out variables that do not feed into the remaining ones.

**Lemma 2.1** (Ancestral reduction)**.** *Let $W \subseteq C \subseteq \mathbf{V}$. If $W$ is ancestral in $\mathcal{G}[C]$ (it contains all its ancestors within the subgraph induced by $C$), then*

$$Q[W] = \sum_{\mathbf{c}\setminus\mathbf{w}} Q[C].$$

Proof idea: sum out the variables of $C \setminus W$ in reverse topological order; each one, when summed, appears only in its own term (it has no children left in $C$) and that term sums to one. For $C = \{V_1, V_2, V_3\}$ in $\mathcal{G}[C]$ ($V_1 \to V_2 \to V_3$): $\{V_1, V_2\}$ and $\{V_1\}$ are ancestral, $\{V_2, V_3\}$ is not.

## 2.2 C-component factorisation

**Lemma 2.2** (Tian 2002)**.** *For any $H \subseteq \mathbf{V}$, let $H_1, \dots, H_k$ be the c-components of $\mathcal{G}[H]$. Then*

$$Q[H] = \prod_{j=1}^{k} Q[H_j].$$

*Moreover, let $V^{(1)} < \cdots < V^{(|H|)}$ be a topological order of $\mathcal{G}[H]$ and $H^{\le i}$ the variables of $H$ up to and including $V^{(i)}$. Then each factor is computable from $Q[H]$:*

$$Q[H_j] = \prod_{V^{(i)} \in H_j} \frac{Q[H^{\le i}]}{Q[H^{\le i-1}]}, \qquad Q[H^{\le i}] = \sum_{\mathbf{h}\setminus\mathbf{h}^{\le i}} Q[H].$$

($H^{\le i}$ is ancestral in $\mathcal{G}[H]$, so $Q[H^{\le i}]$ follows by ancestral reduction.)

**Example 2.3** (Factorising $P(\mathbf{v})$ by c-components)**.** Graph: $Z_2 \to X$, $Z_2 \to Z_1$, $Z_2 \to Z_3$, $X \to Z_1$, $Z_1 \to Y$, $Z_3 \to Y$, $Z_2 \leftrightarrow X$, $X \leftrightarrow Z_3$, $X \leftrightarrow Y$, $Z_2 \leftrightarrow Y$. The c-components are $\{Z_2, Z_3, X, Y\}$ and $\{Z_1\}$, so

$$Q[\mathbf{V}] = Q[Z_2, Z_3, X, Y]\, Q[Z_1],$$
$$P(\mathbf{v}) = P(z_2, z_3, x, y \mid \mathrm{do}(z_1))\, P(z_1 \mid \mathrm{do}(z_2, z_3, x, y)) = P(z_2, z_3, x, y \mid \mathrm{do}(z_1))\, P(z_1 \mid \mathrm{do}(z_2, x)),$$

the last step because only the parents $Z_2, X$ of $Z_1$ matter (Rule 3).

**Example 2.4** (Running example)**.** With $H = \mathbf{V}$ and the order $V_1 < \cdots < V_5$, $Q[\mathbf{V}] = Q[V_1, V_3, V_5]Q[V_2, V_4]$ and

$$\begin{aligned}
Q[V_1, V_3, V_5] &= \frac{Q[V_1]}{Q[\emptyset]} \cdot \frac{Q[V_1, V_2, V_3]}{Q[V_1, V_2]} \cdot \frac{Q[V_1, \dots, V_5]}{Q[V_1, \dots, V_4]} \\
&= \frac{\sum_{v_2,\dots,v_5} P(\mathbf{v})}{\sum_{\mathbf{v}} P(\mathbf{v})} \cdot \frac{\sum_{v_4,v_5} P(\mathbf{v})}{\sum_{v_3,v_4,v_5} P(\mathbf{v})} \cdot \frac{P(\mathbf{v})}{\sum_{v_5} P(\mathbf{v})} \\
&= \frac{P(v_1)}{1} \cdot \frac{P(v_1, v_2, v_3)}{P(v_1, v_2)} \cdot \frac{P(\mathbf{v})}{P(v_1, \dots, v_4)} = P(v_1)P(v_3 \mid v_1, v_2)P(v_5 \mid v_1, \dots, v_4), \\
Q[V_2, V_4] &= \frac{Q[V_1, V_2]}{Q[V_1]} \cdot \frac{Q[V_1, \dots, V_4]}{Q[V_1, V_2, V_3]} = P(v_2 \mid v_1)P(v_4 \mid v_1, v_2, v_3).
\end{aligned}$$

In $\mathcal{G}[\{V_2, V_4\}]$ there are no directed edges ($V_2 \leftrightarrow V_4$ only), so both $\{V_2\}$ and $\{V_4\}$ are ancestral there:

$$Q[V_2] = \sum_{v_4} Q[V_2, V_4] = P(v_2 \mid v_1) \sum_{v_4} P(v_4 \mid v_1, v_2, v_3) = P(v_2 \mid v_1),$$
$$Q[V_4] = \sum_{v_2} P(v_2 \mid v_1)P(v_4 \mid v_1, v_2, v_3).$$

### 2.3 Visualisation

Starting from $Q[\mathbf{V}]$ one can move down a lattice of c-factors: summing out a variable that is "last" (ancestral reduction, e.g. $Q[\mathbf{V}] \to Q[V_{1234}] \to Q[V_{123}] \to Q[V_{12}] \to Q[V_1]$) or splitting along c-components (e.g. $Q[\mathbf{V}] \to Q[V_{135}] \cdot Q[V_{24}]$, $Q[V_{1234}] \to Q[V_{13}] \cdot Q[V_{24}]$, $Q[V_{123}] \to Q[V_{13}] \cdot Q[V_2]$). From $Q[V_{135}]$ every subset is reachable by summation since $\mathcal{G}[\{V_1, V_3, V_5\}]$ has no directed edges, and $Q[V_{15}] = Q[V_1]Q[V_5]$ factorises because $V_1, V_5$ are not bidirected-connected inside $\{V_1, V_5\}$. Once a factor is fully factorised it cannot be factorised further; summation is commutative.

**Summary of the algebra.**

1. *Reduce to an ancestral set:* $Q[W] = \sum_{\mathbf{c}\setminus\mathbf{w}} Q[C]$ if $W$ is ancestral in $\mathcal{G}[C]$.
2. *Factorise into c-components:* $Q[H] = \prod_j Q[H_j]$, with $H_1, \ldots, H_k$ the c-components of $\mathcal{G}[H]$, each computable from $Q[H]$.

These two operations yield an if-and-only-if condition for identifiability.

## 3 Causal queries in terms of c-factors: examples

**Purpose.** Rederive the back-door, front-door and napkin formulas with c-factor algebra.

### 3.1 Back-door

$Z \to X$, $Z \to Y$, $X \to Y$ (Markovian): $P(\mathbf{v}) = Q[X]Q[Y]Q[Z]$. In $\mathcal{G}_{\overline{X}}$, $P(y \mid \mathrm{do}(x)) = \sum_z Q[Y]Q[Z]$ with

$$Q[Y] = \frac{Q[X,Y,Z]}{Q[X,Z]} = \frac{P(\mathbf{v})}{\sum_y P(\mathbf{v})} = P(y \mid x, z), \qquad Q[Z] = \sum_{x,y} Q[X,Y,Z] = P(z),$$

so $P(y \mid \mathrm{do}(x)) = \sum_z P(y \mid z, x)P(z)$.

### 3.2 Front-door

$X \to Z \to Y$, $X \leftrightarrow Y$: $P(\mathbf{v}) = Q[X,Y] \cdot Q[Z]$ with (order $X < Z < Y$)

$$Q[X,Y] = \frac{Q[X]}{Q[\emptyset]} \cdot \frac{Q[X,Z,Y]}{Q[X,Z]} = P(x)P(y \mid x, z), \qquad Q[Z] = \frac{Q[X,Z]}{Q[X]} = P(z \mid x).$$

Then

$$P(y \mid \mathrm{do}(x)) = \sum_z P(y, z \mid \mathrm{do}(x)) = \sum_z Q[Y,Z] = \sum_z Q[Y]Q[Z],$$

and since $\{Y\}$ is ancestral in $\mathcal{G}[\{X,Y\}]$ (no directed edge between them), $Q[Y] = \sum_{x'} Q[X,Y] = \sum_{x'} P(x')P(y \mid x', z)$. Hence $P(y \mid \mathrm{do}(x)) = \sum_z P(z \mid x) \sum_{x'} P(x')P(y \mid x', z)$.

### 3.3 Napkin

$W_1 \to W_2 \to X \to Y$, $W_1 \leftrightarrow X$, $W_1 \leftrightarrow Y$. Then $P(\mathbf{v}) = Q[W_1, X, Y]\, Q[W_2]$ and

$$P(y \mid \mathrm{do}(x)) = \sum_{w_1, w_2} Q[W_1, W_2, Y] = Q[Y],$$

because $\{Y\}$ is ancestral in $\mathcal{G}[\{W_1, W_2, Y\}]$ (in that subgraph $Y$ has no parents). Can we get $Q[Y]$ from $Q[W_1, X, Y]$, which is computable from $Q[\mathbf{V}]$? With the order $W_1 < W_2 < X < Y$,

$$Q[W_1, X, Y] = \frac{Q[W_1]}{Q[\emptyset]} \cdot \frac{Q[W_1, W_2, X]}{Q[W_1, W_2]} \cdot \frac{Q[W_1, W_2, X, Y]}{Q[W_1, W_2, X]} = P(w_1)P(x \mid w_1, w_2)P(y \mid w_1, w_2, x).$$

$\{X, Y\}$ is ancestral in $\mathcal{G}[\{W_1, X, Y\}]$, so $Q[X, Y] = \sum_{w_1} P(w_1)P(x \mid w_1, w_2)P(y \mid w_1, w_2, x)$. In $\mathcal{G}[\{X, Y\}]$, $X$ and $Y$ are no longer bidirected-connected ($W_1$ is gone), so $Q[X, Y] = Q[X]Q[Y]$, giving

$$P(y \mid \mathrm{do}(x)) = Q[Y] = \frac{Q[X, Y]}{Q[X]} = \frac{Q[X, Y]}{\sum_y Q[X, Y]} = \frac{\sum_{w_1} P(w_1)P(x \mid w_1, w_2)P(y \mid w_1, w_2, x)}{\sum_{w_1} P(w_1)P(x \mid w_1, w_2)}.$$

**The same result by do-calculus.**

$$\begin{aligned}
P(y \mid \mathrm{do}(x)) &= P(y \mid \mathrm{do}(x, w_1, w_2)) && \text{Rule 3} \\
&= P(y \mid x, \mathrm{do}(w_1, w_2)) && \text{Rule 2} \\
&= \frac{P(y, x \mid \mathrm{do}(w_1, w_2))}{P(x \mid \mathrm{do}(w_1, w_2))} = \frac{P(y, x \mid \mathrm{do}(w_2))}{P(x \mid \mathrm{do}(w_2))} && \text{Rule 3} \\
&= \frac{\sum_{w_1} P(y, x \mid \mathrm{do}(w_2), w_1)P(w_1 \mid \mathrm{do}(w_2))}{\sum_{w_1} P(x \mid \mathrm{do}(w_2), w_1)P(w_1 \mid \mathrm{do}(w_2))} && \\
&= \frac{\sum_{w_1} P(y, x \mid w_2, w_1)P(w_1)}{\sum_{w_1} P(x \mid w_2, w_1)P(w_1)} && \text{Rules 2 and 3.}
\end{aligned}$$

### 3.4 Overview of the three derivations

| Back-door | Front-door | Napkin |
|---|---|---|
| $Q[\mathbf{V}] \xrightarrow{\text{fact.}} Q[X]\,Q[Y]\,Q[Z]$ | $Q[\mathbf{V}] \xrightarrow{\text{fact.}} Q[X, Y]\,Q[Z]$ | $Q[\mathbf{V}] \xrightarrow{\text{fact.}} Q[W_1, X, Y]\,Q[W_2]$ |
| | $Q[X, Y] \xrightarrow{\text{sum}} Q[Y]$ | $Q[W_1, X, Y] \xrightarrow{\text{sum}} Q[X, Y] \xrightarrow{\text{fact.}} Q[X]\,Q[Y]$ |
| $P(y \mid \mathrm{do}(x)) = \sum_z Q[Y]Q[Z]$ | $P(y \mid \mathrm{do}(x)) = \sum_z Q[Y]Q[Z]$ | $P(y \mid \mathrm{do}(x)) = Q[Y]$ |

## 4 A general identification algorithm

**Purpose.** Combine the two operations into a sound and complete procedure.

### 4.1 Reducing the query to c-factors

Given $\mathcal{G}$ and $P(\mathbf{y} \mid \mathrm{do}(\mathbf{x}))$, first use Rule 3 to enlarge the intervention: every variable that is not an ancestor of $\mathbf{Y}$ in $\mathcal{G}_{\overline{\mathbf{X}}}$ (non-ancestors of $\mathbf{Y}$, and variables that affect $\mathbf{Y}$ only through $\mathbf{X}$) can be intervened on without changing the query. Let $C = An(\mathbf{Y})_{\mathcal{G}_{\overline{\mathbf{X}}}}$. Then

$$P(\mathbf{y} \mid \mathrm{do}(\mathbf{x})) = \sum_{\mathbf{c} \setminus \mathbf{y}} P(\mathbf{c} \mid \mathrm{do}(\mathbf{v} \setminus \mathbf{c})) = \sum_{\mathbf{v} \setminus (\mathbf{x} \cup \mathbf{y})} Q[\mathbf{V} \setminus \mathbf{X}] = \sum_{\mathbf{c} \setminus \mathbf{y}} Q[C].$$

If $\mathcal{G}[C]$ has c-components $C_1, \ldots, C_k$, then

$$P(\mathbf{y} \mid \mathrm{do}(\mathbf{x})) = \sum_{\mathbf{c} \setminus \mathbf{y}} \prod_{i=1}^{k} Q[C_i],$$

while the data give $P(\mathbf{v}) = \prod_j Q[T_j]$ with $T_1, \ldots, T_\ell$ the c-components of $\mathcal{G}$, each $Q[T_j]$ computable from $P(\mathbf{v})$. Each $C_i$ lies inside a single $T_j$. It remains to decide whether $Q[C_i]$ is computable from $Q[T_j]$.

### 4.2 The Identify procedure

**Algorithm 1** IDENTIFY$(C, T, Q[T], \mathcal{G})$ (decides whether $Q[C]$ is computable from $Q[T]$, $C \subseteq T$)

```
A ← An(C)_{G[T]}
if A = C then                                   ▷ C is ancestral in T
    return Q[C] = Σ_{t\c} Q[T]                  ▷ marginalise
else if A = T then                              ▷ nothing to marginalise, unable to factorise
    return FAIL
else
    A_i ← the c-component of G[A] containing C
    obtain Q[A] = Σ_{t\a} Q[T], then Q[A_i] by c-component factorisation   ▷ marginalise, then factorise
    return IDENTIFY(C, A_i, Q[A_i], G)
```

**The three cases.** (1) $A = C$: marginalise, $Q[C] = \sum_{\mathbf{t}\setminus\mathbf{c}} Q[T]$. (2) $A = T$: $C$ is strictly inside $T$, but $T$ is its own ancestral closure and a single c-component, so $T$ cannot be shrunk: fail. (3) $C \subsetneq A \subsetneq T$: marginalise to $A$, factorise $A$ into c-components $A_i, A_j, \ldots$, keep the one containing $C$ (it must be contained in a single component because $C$ is a c-component of $\mathcal{G}[C]$ and $C \subseteq A_i$), and recurse. Each step strictly shrinks $T$ (recursive shrinking: $T \supsetneq A_i \supsetneq \cdots$), so the procedure terminates in at most $|T|$ steps.

**Theorem 4.1** (Completeness; Tian 2002)**.** *$P(\mathbf{y} \mid \mathrm{do}(\mathbf{x}))$ is identifiable from $\mathcal{G}$ and $P(\mathbf{v})$ if and only if each c-factor $Q[C_i]$ is identified by* IDENTIFY$(C_i, T_i, Q[T_i], \mathcal{G})$*, where $T_i$ is the c-component of $\mathcal{G}$ containing $C_i$.*

See also Huang and Valtorta (2006) and Shpitser and Pearl (2006), who proved completeness of the equivalent ID algorithm and, through it, of do-calculus. The failure case $A = T$ corresponds to a graphical structure (a "hedge") from which two models agreeing on $P(\mathbf{v})$ but not on the query can be built.

**Example 4.2** (Bow graph fails)**.** $X \to Y$, $X \leftrightarrow Y$, query $P(y \mid \mathrm{do}(x))$: $C = An(Y)_{\mathcal{G}_{\overline{X}}} = \{Y\}$, $T = \{X, Y\}$. IDENTIFY$(\{Y\}, \{X, Y\}, \ldots)$: $A = An(Y)_{\mathcal{G}[T]} = \{X, Y\} = T$, so it fails, matching Lecture 4.

**Example 4.3** (Front-door succeeds)**.** $X \to Z \to Y$, $X \leftrightarrow Y$: $C = \{Z, Y\}$ with c-components $\{Z\}$, $\{Y\}$ in $\mathcal{G}[C]$. $Q[Z]$ is itself a c-component of $\mathcal{G}$. For $Q[Y]$ with $T = \{X, Y\}$: $A = An(Y)_{\mathcal{G}[T]} = \{Y\} = C$ (the edge $X \to Z \to Y$ is not inside $T$), so $Q[Y] = \sum_x Q[X, Y]$, as found above.

## 5 Source coverage

The source deck has 91 physical pages (41 numbered slides with build-ups).

| Topic | Physical pages |
|---|---|
| Roadmap, c-components, Markovian and semi-Markovian factorisation | 1–15 |
| C-factors, causal-effect interpretation, marginalisation | 16–27 |
| C-component factorisation and examples | 28–43 |
| Visualisation and summary of the algebra | 44–54 |
| Back-door, front-door, napkin via c-factors and do-calculus | 55–76 |
| General algorithm, Identify, completeness, visualisation, references | 77–91 |