Chapter 2. Matrix, Calculus and Probability Tools

Matrix identities, eigenvalue inequalities, matrix calculus, optimization, convergence

Contents

1 Basic Matrix Operations **2**

1.1 Block-matrix formulas 3

2 Eigenvalues and Positive Definiteness **3**

3 Eigenvalue Inequalities **4**

4 Vector Calculus **5**

5 Matrix Calculus **5**

6 Optimization **6**

7 Probability and Convergence **7**

This chapter collects the tools used in later proofs (MLE, Wishart, T^2, large-sample theory).

1 Basic Matrix Operations

Definition 2.1: basic facts

(i) $(\mathbf{AB})_{ik} = \sum_j a_{ij}b_{jk}$; $(\mathbf{A}^\top)^\top = \mathbf{A}$, $(\mathbf{AB})^\top = \mathbf{B}^\top\mathbf{A}^\top$; a 1×1 matrix equals its transpose.

(ii) $\operatorname{tr}(\mathbf{C}) = \sum_i c_{ii}$ is linear.

(iii) $|\mathbf{AB}| = |\mathbf{A}||\mathbf{B}|$, so $|\mathbf{A}^{-1}| = 1/|\mathbf{A}|$.

(iv) (Leibniz) $|\mathbf{A}| = \sum_\sigma \operatorname{sgn}(\sigma)\prod_i a_{i\sigma(i)}$; hence $|\mathbf{A}|$ is a polynomial in the entries, continuous and infinitely differentiable, and $\{|\mathbf{A}| \neq 0\}$ is open. $|\mathbf{I}_p| = 1$, $|\operatorname{diag}(d_i)| = \prod d_i$, $|c\mathbf{A}| = c^p|\mathbf{A}|$.

(v) (Adjugate) If $|\mathbf{A}| \neq 0$, $\mathbf{A}^{-1} = \operatorname{adj}(\mathbf{A})/|\mathbf{A}|$, so $\mathbf{A} \mapsto \mathbf{A}^{-1}$ is continuous and differentiable on $\{|\mathbf{A}| \neq 0\}$.

(vi) (Cauchy–Schwarz) $|\sum_k a_k b_k| \le (\sum a_k^2)^{1/2}(\sum b_k^2)^{1/2}$.

Definition 2.2: Frobenius norm

$\|\mathbf{A}\|_F = (\sum_{i,j} a_{ij}^2)^{1/2}$ and $\|\mathbf{x}\| = \|\mathbf{x}\|_F$ for vectors. Example: $\mathbf{M} = \begin{pmatrix} -0.1 & 0.3 \\ 0.2 & -0.2 \end{pmatrix}$ has $\|\mathbf{M}\|_F = \sqrt{0.18} \approx 0.424$. Estimation errors of the sample moments can be measured by $E_{\bar u} = \|\bar{\mathbf{u}} - \boldsymbol{\mu}\|_2$, $E_S = \|\mathbf{S} - \boldsymbol{\Sigma}\|_F$, $E_R = \|\mathbf{R} - \mathbf{R}_0\|_F$.

Theorem 2.3: basic identities

For $\mathbf{x} \in \mathbb{R}^p$, $\mathbf{y} \in \mathbb{R}^q$, $\mathbf{A} \in \mathbb{R}^{p\times q}$, $\mathbf{B} \in \mathbb{R}^{q\times p}$:

(i) $\mathbf{x}^\top\mathbf{A}\mathbf{y} = \sum_{i,j} a_{ij}x_i y_j$; if $\mathbf{A}$ is diagonal, $\mathbf{x}^\top\mathbf{A}\mathbf{x} = \sum_i a_{ii}x_i^2$.

(ii) $\operatorname{tr}(\mathbf{AB}) = \operatorname{tr}(\mathbf{BA})$.

(iii) $\|\mathbf{A}\|_F^2 = \operatorname{tr}(\mathbf{A}^\top\mathbf{A})$ and $\|\mathbf{x}\|^2 = \mathbf{x}^\top\mathbf{x}$.

Proof. (i) $(\mathbf{Ay})_i = \sum_j a_{ij}y_j$, so $\mathbf{x}^\top\mathbf{Ay} = \sum_i x_i(\mathbf{Ay})_i$.

(ii) $\operatorname{tr}(\mathbf{AB}) = \sum_i\sum_j a_{ij}b_{ji} = \sum_j\sum_i b_{ji}a_{ij} = \operatorname{tr}(\mathbf{BA})$.

(iii) $(\mathbf{A}^\top\mathbf{A})_{jj} = \sum_i a_{ij}^2$; sum over j. ■

Theorem 2.4: first-order expansion of the determinant

$|\mathbf{I}_p + t\mathbf{N}| = 1 + t\operatorname{tr}(\mathbf{N}) + O(t^2)$ as $t \to 0$.

Proof. In the Leibniz formula a non-identity σ moves at least two indices, so its term is $O(t^2)$. The identity contributes $\prod_i(1 + tn_{ii}) = 1 + t\sum_i n_{ii} + O(t^2)$. ■

Definition 2.5: open sets and continuity

With distance $\|\mathbf{B} - \mathbf{A}\|_F$: U is open iff every $\mathbf{A} \in U$ has a ball of radius $r > 0$ inside U; f is continuous at $\mathbf{A}$ iff $\forall\epsilon\,\exists\delta$ with $\|\mathbf{B} - \mathbf{A}\|_F < \delta \Rightarrow \|f(\mathbf{B}) - f(\mathbf{A})\|_F < \epsilon$. If every entry of a

map is continuous, so is the map ($\epsilon/\sqrt{m}$ argument).

1.1 Block-matrix formulas

Used for conditional distributions, the Wishart theorem and likelihood ratio tests.

Theorem 2.6: block determinant and inverse

For $\mathbf{A} = \begin{pmatrix} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{A}_{21} & \mathbf{A}_{22} \end{pmatrix}$ with $\mathbf{A}_{22}$ invertible,

$$|\mathbf{A}| = |\mathbf{A}_{22}|\,|\mathbf{A}_{11} - \mathbf{A}_{12}\mathbf{A}_{22}^{-1}\mathbf{A}_{21}| = |\mathbf{A}_{11}|\,|\mathbf{A}_{22} - \mathbf{A}_{21}\mathbf{A}_{11}^{-1}\mathbf{A}_{12}| \quad (\mathbf{A}_{11} \text{ invertible}).$$

With the Schur complement $\mathbf{A}_{11\cdot 2} := \mathbf{A}_{11} - \mathbf{A}_{12}\mathbf{A}_{22}^{-1}\mathbf{A}_{21}$,

$$\mathbf{A}^{-1} = \begin{pmatrix} \mathbf{A}_{11\cdot 2}^{-1} & -\mathbf{A}_{11\cdot 2}^{-1}\mathbf{A}_{12}\mathbf{A}_{22}^{-1} \\ -\mathbf{A}_{22}^{-1}\mathbf{A}_{21}\mathbf{A}_{11\cdot 2}^{-1} & \mathbf{A}_{22}^{-1} + \mathbf{A}_{22}^{-1}\mathbf{A}_{21}\mathbf{A}_{11\cdot 2}^{-1}\mathbf{A}_{12}\mathbf{A}_{22}^{-1} \end{pmatrix};$$

in particular $(\mathbf{A}^{-1})_{11} = \mathbf{A}_{11\cdot 2}^{-1}$.

Proof. Block Gaussian elimination:

$$\begin{pmatrix} \mathbf{I} & -\mathbf{A}_{12}\mathbf{A}_{22}^{-1} \\ \mathbf{0} & \mathbf{I} \end{pmatrix} \mathbf{A} \begin{pmatrix} \mathbf{I} & \mathbf{0} \\ -\mathbf{A}_{22}^{-1}\mathbf{A}_{21} & \mathbf{I} \end{pmatrix} = \begin{pmatrix} \mathbf{A}_{11\cdot 2} & \mathbf{0} \\ \mathbf{0} & \mathbf{A}_{22} \end{pmatrix}.$$

The outer factors have determinant 1, giving the determinant formula; inverting both sides gives $\mathbf{A}^{-1}$. ■

Theorem 2.7: matrix determinant lemma and Woodbury identity

$$|\mathbf{A} + \mathbf{u}\mathbf{v}^\top| = |\mathbf{A}|(1 + \mathbf{v}^\top\mathbf{A}^{-1}\mathbf{u}), \qquad |\mathbf{I}_p + \mathbf{u}\mathbf{v}^\top| = 1 + \mathbf{v}^\top\mathbf{u},$$
$$(\mathbf{A} + \mathbf{U}\mathbf{C}\mathbf{V})^{-1} = \mathbf{A}^{-1} - \mathbf{A}^{-1}\mathbf{U}(\mathbf{C}^{-1} + \mathbf{V}\mathbf{A}^{-1}\mathbf{U})^{-1}\mathbf{V}\mathbf{A}^{-1}.$$

Proof. Compute $\begin{vmatrix} \mathbf{A} & -\mathbf{u} \\ \mathbf{v}^\top & 1 \end{vmatrix}$ with both forms of the block formula: $|\mathbf{A}|(1+\mathbf{v}^\top\mathbf{A}^{-1}\mathbf{u}) = 1\cdot|\mathbf{A}+\mathbf{u}\mathbf{v}^\top|$. For Woodbury, multiply the right side by $\mathbf{A} + \mathbf{U}\mathbf{C}\mathbf{V}$ and check that the product is $\mathbf{I}$. ■

2 Eigenvalues and Positive Definiteness

Theorem 2.8: spectral decomposition and consequences

Symmetric $\mathbf{A} = \mathbf{U}\mathbf{\Lambda}(\mathbf{A})\mathbf{U}^\top = \sum_i \lambda_i(\mathbf{A})\mathbf{u}_i\mathbf{u}_i^\top$, $\lambda_1(\mathbf{A}) \geq \cdots \geq \lambda_p(\mathbf{A})$ ($\lambda_i(\mathbf{A})$ denotes the ith largest eigenvalue).

$$|\mathbf{A}| = \prod_i \lambda_i(\mathbf{A}), \qquad \operatorname{tr}(\mathbf{A}) = \sum_i \lambda_i(\mathbf{A}), \qquad \mathbf{A} \succ 0 \iff \text{all } \lambda_i(\mathbf{A}) > 0.$$

Proof. $|\mathbf{U}||\mathbf{U}^\top| = |\mathbf{U}\mathbf{U}^\top| = 1$ gives $|\mathbf{A}| = |\mathbf{\Lambda}(\mathbf{A})|$; $\operatorname{tr}(\mathbf{U}\mathbf{\Lambda}\mathbf{U}^\top) = \operatorname{tr}(\mathbf{\Lambda}\mathbf{U}^\top\mathbf{U}) = \operatorname{tr}\mathbf{\Lambda}$. Definiteness follows from $\mathbf{x}^\top\mathbf{A}\mathbf{x} = \sum_i \lambda_i(\mathbf{u}_i^\top\mathbf{x})^2$. ■

Theorem 2.9: smallest eigenvalue

For symmetric $\mathbf{A}, \mathbf{B}$:
(i) $\lambda_p(\mathbf{A}) = \min_{\|\mathbf{x}\|=1} \mathbf{x}^\top \mathbf{A}\mathbf{x}$;
(ii) $|\lambda_p(\mathbf{A}) - \lambda_p(\mathbf{B})| \le \|\mathbf{A} - \mathbf{B}\|_F$;
(iii) $\{\mathbf{A} \succ 0\}$ is convex and open in the space of symmetric matrices.

Proof. (i) With $\mathbf{z} = \mathbf{U}^\top \mathbf{x}$, $\|\mathbf{z}\| = 1$ and $\mathbf{x}^\top \mathbf{A}\mathbf{x} = \sum \lambda_i z_i^2 \ge \lambda_p$, equality at $\mathbf{x} = \mathbf{u}_p$.

(ii) For $\|\mathbf{x}\| = 1$, Cauchy–Schwarz gives $|\mathbf{x}^\top \mathbf{A}\mathbf{x} - \mathbf{x}^\top \mathbf{B}\mathbf{x}| = |\sum_{ij}(a_{ij} - b_{ij})x_i x_j| \le \|\mathbf{A} - \mathbf{B}\|_F (\sum_{ij} x_i^2 x_j^2)^{1/2} = \|\mathbf{A} - \mathbf{B}\|_F$; minimize both sides over $\mathbf{x}$.

(iii) Convex: $\mathbf{x}^\top\{(1-s)\mathbf{A}_0 + s\mathbf{A}_1\}\mathbf{x} > 0$. Open: if $\mathbf{A} \succ 0$ and $\|\mathbf{B} - \mathbf{A}\|_F < \lambda_p(\mathbf{A})$, (ii) gives $\lambda_p(\mathbf{B}) \ge \lambda_p(\mathbf{A}) - \|\mathbf{A} - \mathbf{B}\|_F > 0$. ■

Theorem 2.10: symmetric square root

For $\mathbf{A} \succ 0$, $\mathbf{A}^{\pm 1/2} = \mathbf{U}\mathbf{\Lambda}^{\pm 1/2}\mathbf{U}^\top$ are symmetric, $\mathbf{A}^{1/2}\mathbf{A}^{1/2} = \mathbf{A}$, $\mathbf{A}^{-1/2}\mathbf{A}^{-1/2} = \mathbf{A}^{-1}$, $\mathbf{A}^{-1/2}\mathbf{A}\mathbf{A}^{-1/2} = \mathbf{I}_p$. $\mathbf{A}^{-1} = \mathbf{U}\mathbf{\Lambda}^{-1}\mathbf{U}^\top \succ 0$ and $\lambda_i(\mathbf{A}^{-1}) = 1/\lambda_{p+1-i}(\mathbf{A})$ with eigenvector $\mathbf{u}_{p+1-i}$.

3 Eigenvalue Inequalities

Definition 2.11: Birkhoff's theorem and the rearrangement inequality

(i) A non-negative matrix with all row and column sums 1 (doubly stochastic) is a convex combination of permutation matrices.
(ii) If $a_1 \ge \cdots \ge a_p$ and $b_1 \ge \cdots \ge b_p$, then for every permutation σ, $\sum_i a_i b_{p+1-i} \le \sum_i a_i b_{\sigma(i)} \le \sum_i a_i b_i$.

Theorem 2.12: von Neumann's trace inequality

For symmetric $\mathbf{A} = \sum_i \lambda_i(\mathbf{A})\mathbf{u}_i\mathbf{u}_i^\top$, $\mathbf{B} = \sum_j \lambda_j(\mathbf{B})\mathbf{v}_j\mathbf{v}_j^\top$,

$$\sum_{i=1}^{p} \lambda_{p+1-i}(\mathbf{A})\lambda_i(\mathbf{B}) \le \operatorname{tr}(\mathbf{A}\mathbf{B}) \le \sum_{i=1}^{p} \lambda_i(\mathbf{A})\lambda_i(\mathbf{B}).$$

The upper bound is attained when $\mathbf{v}_i = \mathbf{u}_i$ and the lower bound when $\mathbf{v}_i = \mathbf{u}_{p+1-i}$.

Proof. Step 1. $\operatorname{tr}(\mathbf{A}\mathbf{B}) = \sum_{i,j} \lambda_i(\mathbf{A})\lambda_j(\mathbf{B}) w_{ij}$, $w_{ij} = (\mathbf{u}_i^\top \mathbf{v}_j)^2 \ge 0$. Since $\{\mathbf{u}_i\}, \{\mathbf{v}_j\}$ are orthonormal bases, $\sum_j w_{ij} = \|\mathbf{u}_i\|^2 = 1$ and $\sum_i w_{ij} = \|\mathbf{v}_j\|^2 = 1$: $\mathbf{W}$ is doubly stochastic. **Step 2.** By Birkhoff $\mathbf{W}$ is a convex combination of permutation matrices; $\operatorname{tr}(\mathbf{A}\mathbf{B})$ is linear in $\mathbf{W}$, so it lies between the minimum and maximum of $\sum_i \lambda_i(\mathbf{A})\lambda_{\sigma(i)}(\mathbf{B})$. **Step 3.** By rearrangement the maximum is at $\sigma(i) = i$, the minimum at $\sigma(i) = p+1-i$. **Step 4.** If $\mathbf{v}_i = \mathbf{u}_i$ (resp. $\mathbf{u}_{p+1-i}$), $\mathbf{W}$ is the identity (resp. reversal) permutation and the bound is attained. ■

Theorem 2.13: an elementary inequality

$x - \log x \ge 1$ for $x > 0$, with equality iff $x = 1$.

Proof. $\phi(x) = x - \log x$ has $\phi'(x) = 1 - 1/x$, negative on $(0,1)$, positive on $(1,\infty)$. For $x \ne 1$ the mean value theorem gives $\phi(x) - \phi(1) = \phi'(\xi)(x-1) > 0$ with ξ between 1 and x. ■

4 Vector Calculus

Definition 2.14: gradient, Jacobian, Hessian

For $f:\mathbb{R}^p\to\mathbb{R}$, $\partial f/\partial\mathbf{x}=(\partial f/\partial x_1,\dots,\partial f/\partial x_p)^\top$. The Jacobian of $\mathbf{g}:\mathbb{R}^p\to\mathbb{R}^q$ is the $q\times p$ matrix $(\partial g_i/\partial x_j)$. The Hessian $\partial^2 f/\partial\mathbf{x}\partial\mathbf{x}^\top$ is the Jacobian of the gradient; it is symmetric when second partials are continuous (Schwarz's theorem).

Theorem 2.15: chain rule

For $\mathbf{y}=\mathbf{g}(\mathbf{x})$ with $q\times p$ Jacobian $\mathbf{J}=\partial\mathbf{g}/\partial\mathbf{x}^\top$,

$$\frac{\partial f(\mathbf{y})}{\partial\mathbf{x}}=\mathbf{J}^\top\frac{\partial f}{\partial\mathbf{y}},\qquad \frac{\partial^2 f(\mathbf{y})}{\partial\mathbf{x}\partial\mathbf{x}^\top}=\mathbf{J}^\top\frac{\partial^2 f}{\partial\mathbf{y}\partial\mathbf{y}^\top}\mathbf{J}+\sum_{i=1}^{q}\frac{\partial f}{\partial y_i}\frac{\partial^2 g_i}{\partial\mathbf{x}\partial\mathbf{x}^\top}.$$

For an affine map $\mathbf{g}(\mathbf{x})=\mathbf{B}\mathbf{x}+\mathbf{b}$, $\mathbf{J}=\mathbf{B}$ and the last sum vanishes.

Theorem 2.16: derivatives of quadratic forms and basic maps

For symmetric $\mathbf{A}$, $\dfrac{\partial\,\mathbf{x}^\top\mathbf{A}\mathbf{x}}{\partial\mathbf{x}}=2\mathbf{A}\mathbf{x}$ and $\dfrac{\partial^2\,\mathbf{x}^\top\mathbf{A}\mathbf{x}}{\partial\mathbf{x}\partial\mathbf{x}^\top}=2\mathbf{A}$. Also $\partial\mathbf{x}/\partial\mathbf{x}^\top=\mathbf{I}_p$, $\partial(\mathbf{B}\mathbf{x}+\mathbf{b})/\partial\mathbf{x}^\top=\mathbf{B}$, and $\partial(\mathbf{x}-\boldsymbol{\mu})/\partial\boldsymbol{\mu}^\top=-\mathbf{I}_p$.

Proof. $\mathbf{x}^\top\mathbf{A}\mathbf{x}=\sum_{i,j}a_{ij}x_ix_j$, so $\partial/\partial x_k=\sum_j a_{kj}x_j+\sum_i a_{ik}x_i=2(\mathbf{A}\mathbf{x})_k$; differentiating in x_l gives $2a_{kl}$. ■

5 Matrix Calculus

To optimize over covariance matrices we differentiate with respect to a matrix: matrix gradients give first-order conditions and directional derivatives give second-order conditions along symmetric directions.

Definition 2.17: directional derivatives and matrix gradient

For f on an open set C, $\boldsymbol{\Theta}\in C$ and a direction $\mathbf{H}$,

$$Df(\boldsymbol{\Theta})[\mathbf{H}]=\frac{d}{dt}f(\boldsymbol{\Theta}+t\mathbf{H})\Big|_{t=0},\qquad D^2f(\boldsymbol{\Theta})[\mathbf{H}]=\frac{d^2}{dt^2}f(\boldsymbol{\Theta}+t\mathbf{H})\Big|_{t=0}.$$

The matrix gradient $\partial f/\partial\mathbf{X}$ has (i,j) entry $\partial f/\partial x_{ij}$.

Theorem 2.18: directional derivative and gradient

In $\mathbb{R}^p$: $Df(\boldsymbol{\mu})[\mathbf{h}]=\mathbf{h}^\top\frac{\partial f}{\partial\boldsymbol{\mu}}$, $D^2f(\boldsymbol{\mu})[\mathbf{h}]=\mathbf{h}^\top\frac{\partial^2 f}{\partial\boldsymbol{\mu}\partial\boldsymbol{\mu}^\top}\mathbf{h}$. For matrices: $Df(\mathbf{X})[\mathbf{H}]=\operatorname{tr}\Big\{\Big(\dfrac{\partial f}{\partial\mathbf{X}}\Big)^\top\mathbf{H}\Big\}=\sum_{i,j}\dfrac{\partial f}{\partial x_{ij}}h_{ij}$.

Theorem 2.19: derivatives of basic matrix functions

For $\boldsymbol{\Sigma}\succ 0$, symmetric $\mathbf{H}$ and $p\times p$ $\mathbf{B}$:

$$D\log|\boldsymbol{\Sigma}|[\mathbf{H}]=\operatorname{tr}(\boldsymbol{\Sigma}^{-1}\mathbf{H}),\qquad D^2\log|\boldsymbol{\Sigma}|[\mathbf{H}]=-\operatorname{tr}(\boldsymbol{\Sigma}^{-1}\mathbf{H}\boldsymbol{\Sigma}^{-1}\mathbf{H}),$$

$$D\,\mathrm{tr}(\boldsymbol{\Sigma}\mathbf{B})[\mathbf{H}] = \mathrm{tr}(\mathbf{H}\mathbf{B}), \qquad D^2\,\mathrm{tr}(\boldsymbol{\Sigma}\mathbf{B})[\mathbf{H}] = 0,$$
$$\frac{\partial \log|\boldsymbol{\Sigma}|}{\partial \boldsymbol{\Sigma}} = \boldsymbol{\Sigma}^{-1}, \qquad \frac{\partial\,\mathrm{tr}(\boldsymbol{\Sigma}\mathbf{B})}{\partial \boldsymbol{\Sigma}} = \mathbf{B}^\top.$$

D and D^2 are linear.

Proof. Let $\mathbf{M} = \boldsymbol{\Sigma}^{-1/2}\mathbf{H}\boldsymbol{\Sigma}^{-1/2} = \mathbf{V}\boldsymbol{\Lambda}(\mathbf{M})\mathbf{V}^\top$ (symmetric). Then $\boldsymbol{\Sigma} + t\mathbf{H} = \boldsymbol{\Sigma}^{1/2}(\mathbf{I} + t\mathbf{M})\boldsymbol{\Sigma}^{1/2}$ and $\mathbf{I} + t\mathbf{M}$ has eigenvalues $1 + t\lambda_k(\mathbf{M})$, so for small $|t|$

$$\log|\boldsymbol{\Sigma} + t\mathbf{H}| = \log|\boldsymbol{\Sigma}| + \sum_k \log\{1 + t\lambda_k(\mathbf{M})\}.$$

The first and second derivatives at $t = 0$ are $\sum_k \lambda_k(\mathbf{M}) = \mathrm{tr}\mathbf{M} = \mathrm{tr}(\boldsymbol{\Sigma}^{-1}\mathbf{H})$ and $-\sum_k \lambda_k(\mathbf{M})^2 = -\mathrm{tr}(\mathbf{M}^2) = -\mathrm{tr}(\boldsymbol{\Sigma}^{-1}\mathbf{H}\boldsymbol{\Sigma}^{-1}\mathbf{H})$. $\mathrm{tr}\{(\boldsymbol{\Sigma} + t\mathbf{H})\mathbf{B}\}$ is linear in t. Gradients: for any $\mathbf{H}$, $|\boldsymbol{\Sigma} + t\mathbf{H}| = |\boldsymbol{\Sigma}||\mathbf{I} + t\boldsymbol{\Sigma}^{-1}\mathbf{H}| = |\boldsymbol{\Sigma}|\{1 + t\mathrm{tr}(\boldsymbol{\Sigma}^{-1}\mathbf{H}) + O(t^2)\}$, so $D\log|\boldsymbol{\Sigma}|[\mathbf{H}] = \mathrm{tr}\{(\boldsymbol{\Sigma}^{-1})^\top\mathbf{H}\}$; compare with Theorem 2.18. Likewise $\mathrm{tr}(\mathbf{H}\mathbf{B}) = \mathrm{tr}\{(\mathbf{B}^\top)^\top\mathbf{H}\}$. ■

Theorem 2.20: symmetric matrix variables

If $\mathbf{X}$ is symmetric and $\mathbf{G} = \partial f/\partial \mathbf{X}$ is symmetric, then $Df(\mathbf{X})[\mathbf{H}] = 0$ for all symmetric $\mathbf{H}$ iff $\mathbf{G} = \mathbf{0}$.

Proof. $\Leftarrow$ is clear. $\Rightarrow$: take $\mathbf{H} = \mathbf{G}$: $0 = \mathrm{tr}(\mathbf{G}^\top\mathbf{G}) = \|\mathbf{G}\|_F^2$. ■

6 Optimization

Theorem 2.21: convexity and unique minimizer

Let C be open and convex and suppose $D^2 f(\boldsymbol{\Theta})[\mathbf{H}] > 0$ for all $\boldsymbol{\Theta} \in C$, $\mathbf{H} \neq \mathbf{0}$. Then (i) f is strictly convex on C;
(ii) a point $\hat{\boldsymbol{\Theta}}$ with $Df(\hat{\boldsymbol{\Theta}})[\mathbf{H}] = 0$ for all $\mathbf{H}$ is the unique global minimizer;
(iii) for $C = \mathbb{R}^p$ these read: Hessian $\succ 0$ and gradient $= \mathbf{0}$.

Proof. $g(t) = f(\boldsymbol{\Theta}_0 + t(\boldsymbol{\Theta}_1 - \boldsymbol{\Theta}_0))$ has $g''(t) = D^2 f(\cdot)[\mathbf{H}] > 0$, so g is strictly convex.

(ii) With $\boldsymbol{\Theta}_0 = \hat{\boldsymbol{\Theta}}$, $g'(0) = 0$ and g' is strictly increasing, so $g'(t) > 0$ on $(0, 1]$ and $f(\boldsymbol{\Theta}_1) = g(1) > g(0) = f(\hat{\boldsymbol{\Theta}})$ by the mean value theorem. ■

Theorem 2.22: transformations that keep the optimizer; profiling

For $a > 0$ and a constant c: $\arg\min\{af + c\} = \arg\min f$, $\arg\max f = \arg\min(-f)$, $\arg\max L = \arg\max \log L$ for $L > 0$, and for a bijection ϕ, $\arg\min_\theta f(\theta) = \phi^{-1}(\arg\min_\eta f \circ \phi^{-1}(\eta))$. **Profiling**: if $\hat{\theta}(\eta)$ minimizes $\ell(\cdot, \eta)$ for each η and $\hat{\eta}$ minimizes the profile $p(\eta) = \ell(\hat{\theta}(\eta), \eta)$, then $(\hat{\theta}(\hat{\eta}), \hat{\eta})$ minimizes ℓ jointly: $\ell(\theta, \eta) \geq \ell(\hat{\theta}(\eta), \eta) = p(\eta) \geq p(\hat{\eta})$.

The MLE (Ch. 4) first profiles out $\boldsymbol{\mu}$ and then minimizes over $\boldsymbol{\Sigma}$.

7 Probability and Convergence

Definition 2.23: standard results from mathematical statistics

(S0) $\mathbf{Y}_n \xrightarrow{p} \mathbf{c}$: $P(\|\mathbf{Y}_n - \mathbf{c}\| > \epsilon) \to 0$ for all $\epsilon > 0$. $\mathbf{Y}_n \xrightarrow{d} \mathbf{Y}$: $\mathrm{E}g(\mathbf{Y}_n) \to \mathrm{E}g(\mathbf{Y})$ for all bounded continuous g.

(S1) Markov: for $W \geq 0$, $P(W \geq a) \leq \mathrm{E}W/a$.

(S2) $\mathrm{E}(\mathbf{AY} + \mathbf{b}) = \mathbf{A}\mathrm{E}\mathbf{Y} + \mathbf{b}$, $\mathrm{Cov}(\mathbf{AY} + \mathbf{b}) = \mathbf{A}\mathrm{Cov}(\mathbf{Y})\mathbf{A}^\top$; for independent $\mathbf{Y}_j$, $\mathrm{Cov}(\sum \mathbf{Y}_j) = \sum \mathrm{Cov}(\mathbf{Y}_j)$; $\mathrm{E}\|\mathbf{Y} - \mathrm{E}\mathbf{Y}\|^2 = \mathrm{tr}\mathrm{Cov}(\mathbf{Y})$.

(S3) Weak law of large numbers; (S4) univariate CLT.

(S5) Continuous mapping theorem (CMT): $\mathbf{Y}_n \xrightarrow{d} \mathbf{Y}$, g continuous $\Rightarrow$ $g(\mathbf{Y}_n) \xrightarrow{d} g(\mathbf{Y})$; $\mathbf{Y}_n \xrightarrow{p} \mathbf{c}$, g continuous at $\mathbf{c} \Rightarrow g(\mathbf{Y}_n) \xrightarrow{p} g(\mathbf{c})$.

(S6) Slutsky: $\mathbf{Y}_n \xrightarrow{d} \mathbf{Y}$, $\mathbf{W}_n \xrightarrow{p} \mathbf{c} \Rightarrow (\mathbf{Y}_n, \mathbf{W}_n) \xrightarrow{d} (\mathbf{Y}, \mathbf{c})$.

(S7) Componentwise convergence in probability to constants implies joint convergence.

Theorem 2.24: vector Markov inequality and matrix convergence

$P(\|\mathbf{Y}\| > \epsilon) \leq \mathrm{E}\|\mathbf{Y}\|^2/\epsilon^2$. For random matrices, $\mathbf{A}_n \xrightarrow{p} \mathbf{A}$ (entrywise) $\iff$ $\|\mathbf{A}_n - \mathbf{A}\|_F \xrightarrow{p} 0$.

Proof. $\{\|\mathbf{Y}\| > \epsilon\} \subset \{\|\mathbf{Y}\|^2 \geq \epsilon^2\}$; apply Markov. Matrices: $\|\mathbf{A}_n - \mathbf{A}\|_F^2 \leq pq \max_{ij}(\mathbf{A}_n - \mathbf{A})_{ij}^2$, so $\{\|\mathbf{A}_n - \mathbf{A}\|_F > \epsilon\} \subset \bigcup_{ij}\{|(\mathbf{A}_n - \mathbf{A})_{ij}| > \epsilon/\sqrt{pq}\}$ and a union bound gives convergence; conversely $|(\mathbf{A}_n - \mathbf{A})_{ij}| \leq \|\mathbf{A}_n - \mathbf{A}\|_F$. ■

Theorem 2.25: characteristic functions and the Cramér–Wold device

$\varphi_{\mathbf{Y}}(\mathbf{t}) = \mathrm{E}\exp(i\mathbf{t}^\top\mathbf{Y})$.
(i) Uniqueness: $\varphi_{\mathbf{Y}} \equiv \varphi_{\mathbf{W}} \iff \mathbf{Y} \overset{d}{=} \mathbf{W}$.
(ii) Lévy continuity: $\mathbf{Y}_n \xrightarrow{d} \mathbf{Y} \iff \varphi_{\mathbf{Y}_n}(\mathbf{t}) \to \varphi_{\mathbf{Y}}(\mathbf{t})$ for all $\mathbf{t}$. **Cramér–Wold**:
(i) the distribution of $\mathbf{Y}$ is determined by those of all $\mathbf{t}^\top\mathbf{Y}$;
(ii) $\mathbf{Y}_n \xrightarrow{d} \mathbf{Y} \iff \mathbf{t}^\top\mathbf{Y}_n \xrightarrow{d} \mathbf{t}^\top\mathbf{Y}$ for every $\mathbf{t}$.

Proof. The key identity is $\varphi_{\mathbf{t}^\top\mathbf{Y}}(1) = \mathrm{E}e^{i\mathbf{t}^\top\mathbf{Y}} = \varphi_{\mathbf{Y}}(\mathbf{t})$.

(i) If $\mathbf{t}^\top\mathbf{Y} \overset{d}{=} \mathbf{t}^\top\mathbf{W}$, then $\varphi_{\mathbf{Y}}(\mathbf{t}) = \varphi_{\mathbf{t}^\top\mathbf{Y}}(1) = \varphi_{\mathbf{t}^\top\mathbf{W}}(1) = \varphi_{\mathbf{W}}(\mathbf{t})$.

(ii) $\Leftarrow$: Lévy for scalars gives $\varphi_{\mathbf{t}^\top\mathbf{Y}_n}(s) \to \varphi_{\mathbf{t}^\top\mathbf{Y}}(s)$; take $s = 1$. ■

"All one-dimensional linear projections determine the multivariate distribution" underlies the general definition of the multivariate normal (Ch. 3) and the multivariate CLT.