

FULLY DISCRETE NEEDLET APPROXIMATION ON THE SPHERE

Yu Guang Wang

School of Mathematics and Statistics
The University of New South Wales, Australia

AustMS 2015, Adelaide

Joint with Quoc T. Le Gia, Ian H. Sloan and




Robert S. Womersley



UNSW
AUSTRALIA

Applications


Applications



Cosmic Microwave Background

A needlet on the sphere \mathbb{S}^2

A needlet on the sphere \mathbb{S}^2



Needlets


Needlets are **filtered** polynomials
associated with **quadrature rules** on the sphere.

Needlets

Needlets are **filtered** polynomials
associated with **quadrature rules** on the sphere.

$$\psi_{jk}(\mathbf{x}) := \sqrt{w_{jk}} \sum_{\ell=0}^{\infty} h\left(\frac{\ell}{2^{j-1}}\right) (2\ell + 1) P_\ell(\mathbf{x} \cdot \mathbf{x}_{jk}), \quad j \geq 1.$$

C^5 -needlet filter h



- $0 \leq h(t) \leq 1$
- $\text{supp } h \subset [1/2, 2]$
- $[h(t)]^2 + [h(2t)]^2 = 1, \quad t \in [1/2, 1]$
- $h \in C^\kappa(\mathbb{R}_+), \quad \kappa \geq 2$

Quadrature rule for needlets

- Weights w_{jk} , Nodes $\mathbf{x}_{jk} \in \mathbb{S}^2$, $k = 1, \dots, N$.
- Exact for polynomials P of degree $\leq t$

$$\int_{\mathbb{S}^2} P(\mathbf{x}) \, d\sigma_2(\mathbf{x}) = \sum_{k=1}^N w_{jk} P(\mathbf{x}_{jk}).$$

Quadrature rule for needlets

- Weights w_{jk} , Nodes $\mathbf{x}_{jk} \in \mathbb{S}^2$, $k = 1, \dots, N$.
- Exact for polynomials P of degree $\leq t$

$$\int_{\mathbb{S}^2} P(\mathbf{x}) d\sigma_2(\mathbf{x}) = \sum_{k=1}^N w_{jk} P(\mathbf{x}_{jk}).$$


e.g. Symmetric spherical t -design:

- Equal weights $w_{jk} = 1/N$, $k = 1, \dots, N$.
- Exact for degree t .
- $N \sim \mathcal{O}(t^2/2)$.

Womersley., 2015.

Quadrature rule for needlets ψ_{jk} , $j = 4$

Fig. A symmetric spherical 31-design



- Equal weights
- Nodes $\mathbf{x}_{jk} \in \mathbb{S}^2$, $k = 1, \dots, 498$.
- Exact for polynomials of degree up to $2^{4+1} - 1 = 31$

Womersley, 2015.

What is a (semidiscrete) needlet approximation?

For a function f on \mathbb{S}^2 , $J = 0, 1, \dots$,

Definition (Semidiscrete needlet approximation)

$$V_J^{\text{need}}(f; \mathbf{x}) := \sum_{j=0}^J \sum_{k=1}^{N_j} (f, \psi_{jk})_{\mathbb{L}_2(\mathbb{S}^2)} \psi_{jk}(\mathbf{x}), \quad \mathbf{x} \in \mathbb{S}^2,$$

where

$$(f, \psi_{jk})_{\mathbb{L}_2(\mathbb{S}^2)} := \int_{\mathbb{S}^2} f(\mathbf{y}) \psi_{jk}(\mathbf{y}) d\sigma_2(\mathbf{y}).$$

The \mathbb{L}_p errors of needlet approximation

Narcowich, Petrushev & Ward., 2006.

Theorem

The \mathbb{L}_p error using the *needlet approximation* with sufficient smooth filter for $f \in \mathbb{W}_p^s(\mathbb{S}^2)$ and $1 \leq p \leq \infty$ and $s > 2/p$ has the optimal convergence order 2^{-Js} .

If $h \in C^\kappa(\mathbb{R}_+)$, $\kappa \geq 2$,

$$\|f - V_J^{\text{need}}(f)\|_{\mathbb{L}_p(\mathbb{S}^2)} \leq c E_{2^{J-1}}(f)_p \leq c 2^{-Js} \|f\|_{\mathbb{W}_p^s(\mathbb{S}^2)}.$$

The \mathbb{L}_p errors of needlet approximation

Narcowich, Petrushev & Ward., 2006.

Theorem

The \mathbb{L}_p error using the *needlet approximation* with sufficient smooth filter for $f \in \mathbb{W}_p^s(\mathbb{S}^2)$ and $1 \leq p \leq \infty$ and $s > 2/p$ has the optimal convergence order 2^{-Js} .

If $h \in C^\kappa(\mathbb{R}_+)$, $\kappa \geq 2$,

$$\|f - V_J^{\text{need}}(f)\|_{\mathbb{L}_p(\mathbb{S}^2)} \leq c E_{2^{J-1}}(f)_p \leq c 2^{-Js} \|f\|_{\mathbb{W}_p^s(\mathbb{S}^2)}.$$

$$\Rightarrow \sum_{j=0}^{\infty} \sum_{k=1}^{N_j} |(f, \psi_{jk})_{\mathbb{L}_2(\mathbb{S}^2)}|^2 = \|f\|_{\mathbb{L}_2(\mathbb{S}^2)}^2 \text{ (Tight frame).}$$

How to implement needlet approximations?

Given $J = 0, 1, \dots$

Discretisation quadrature $\mathcal{Q}_N := \{(W_i, \mathbf{y}_i) : i = 1, \dots, N\}$

Exact for polynomials of degree $3 \times 2^{J-1} - 1$

Need NOT be the same as needlet quadrature

Approximate needlet coefficient by quadrature \mathcal{Q}_N

$$\begin{aligned}(f, \psi_{jk})_{\mathbb{L}_2(\mathbb{S}^2)} &= \int_{\mathbb{S}^2} f(\mathbf{y}) \psi_{jk}(\mathbf{y}) \, d\sigma_2(\mathbf{y}) \\ &\approx \sum_{i=1}^N W_i f(\mathbf{y}_i) \psi_{jk}(\mathbf{y}_i) =: (f, \psi_{jk})_{\mathcal{Q}_N}\end{aligned}$$

How to implement needlet approximations?

Given $J = 0, 1, \dots$

Discretisation quadrature $\mathcal{Q}_N := \{(W_i, \mathbf{y}_i) : i = 1, \dots, N\}$

Exact for polynomials of degree $3 \times 2^{J-1} - 1$

Definition (Fully discrete needlet approximation)

$$V_{J,N}^{\text{need}}(f; \mathbf{x}) := \sum_{j=0}^J \sum_{k=1}^{N_j} (f, \psi_{jk})_{\mathcal{Q}_N} \psi_{jk}(\mathbf{x}), \quad \mathbf{x} \in \mathbb{S}^2.$$

$$V_J^{\text{need}}(f; \mathbf{x}) = \sum_{j=0}^J \sum_{k=1}^{N_j} (f, \psi_{jk})_{\mathbb{L}_2(\mathbb{S}^2)} \psi_{jk}(\mathbf{x}).$$

Discretisation does not lower the convergence rate!

Given $J = 0, 1, \dots$

Discretisation quadrature $\mathcal{Q}_N := \{(W_i, \mathbf{y}_i) : i = 1, \dots, N\}$

Exact for polynomials of degree $3 \times 2^{J-1} - 1$

Theorem


The **discrete needlet approximation** with smooth filter and \mathcal{Q}_N of a function f in $\mathbb{W}_p^s(\mathbb{S}^2)$, $1 \leq p \leq \infty$ and $s > 2/p$, converges at optimal rate 2^{-Js} in $\mathbb{L}_p(\mathbb{S}^2)$ -norm.

If $h \in C^\kappa(\mathbb{R}_+)$, $\kappa \geq 2$,

$$\|f - V_{J,\textcolor{red}{N}}^{\text{need}}(f)\|_{\mathbb{L}_p(\mathbb{S}^2)} \leq c 2^{-Js} \|f\|_{\mathbb{W}_p^s(\mathbb{S}^2)}, \quad f \in \mathbb{W}_p^s(\mathbb{S}^2).$$

Numerical examples: normalised Wendland functions

Fig. Test RBF function f_2



Chernih, Sloan & Womersley., 2014.

Le Gia, Sloan & Wendland., 2010.

Wendland., 2001.

$$f_k(\mathbf{x}) := \sum_{i=1}^6 \phi_k(|\mathbf{z}_i - \mathbf{x}|), \quad k \geq 0,$$


where $\mathbf{z}_i = (\pm 1, 0, 0), (0, \pm 1, 0), (0, 0, \pm 1)$
and

$$\phi_k(r) := \tilde{\phi}_k\left(\frac{r}{\delta_k}\right), \quad r \in \mathbb{R},$$

$$\phi_k(|\mathbf{z}_i - \mathbf{x}|) \in \mathbb{W}_2^{k+3/2}(\mathbb{S}^2).$$

Numerical examples: normalised Wendland functions

Fig. Test RBF function f_2



Chernih, Sloan & Womersley., 2014.

Le Gia, Sloan & Wendland., 2010.

Wendland., 2001.

$$f_k(\mathbf{x}) := \sum_{i=1}^6 \phi_k(|\mathbf{z}_i - \mathbf{x}|), \quad k \geq 0,$$


where $\mathbf{z}_i = (\pm 1, 0, 0), (0, \pm 1, 0), (0, 0, \pm 1)$
and

$$\phi_k(r) := \tilde{\phi}_k\left(\frac{r}{\delta_k}\right), \quad r \in \mathbb{R},$$


$$\phi_k(|\mathbf{z}_i - \mathbf{x}|) \in \mathbb{W}_2^{k+3/2}(\mathbb{S}^2).$$

e.g. $\tilde{\phi}_2(r) := (1 - r)_+^6 (35r^2 + 18r + 3)/3.$

Numerical examples support the theorems



(a) Semidiscrete



(b) Fully discrete

Key formula for proof

For $f \in C(\mathbb{S}^2)$ and $J \geq 0$,

$$V_{J,N}^{\text{need}}(f) = V_{2^J-1,H,N}(f),$$

where $H(t) = 1$, $0 \leq t < 1$ and $H(t) = h(t)^2$, $t \geq 1$.

Key formula for proof

For $f \in C(\mathbb{S}^2)$ and $J \geq 0$,

$$V_{J,N}^{\text{need}}(f) = V_{2^{J-1},H,N}(f),$$

where $H(t) = 1$, $0 \leq t < 1$ and $H(t) = h(t)^2$, $t \geq 1$.

Filtered hyperinterpolation $V_{2^{J-1},H,N}(f)$ has optimal error order 2^{-Js} for $f \in \mathbb{W}_p^s(\mathbb{S}^2)$, $1 \leq p \leq \infty$ and $s > 2/p$.

Sloan & H. Wang., 2015.

Key formula for proof

For $f \in C(\mathbb{S}^2)$ and $J \geq 0$,

$$V_{J,N}^{\text{need}}(f) = V_{2^{J-1},H,N}(f),$$

where $H(t) = 1$, $0 \leq t < 1$ and $H(t) = h(t)^2$, $t \geq 1$.

Filtered hyperinterpolation $V_{2^{J-1},H,N}(f)$ has optimal error order 2^{-Js} for $f \in \mathbb{W}_p^s(\mathbb{S}^2)$, $1 \leq p \leq \infty$ and $s > 2/p$.

Sloan & H. Wang., 2015.

Using ALL needlets, the error of discrete needlet approximation follows from the error of filtered hyperinterpolation!

Computational challenges in high orders


$$\# \text{ Needlets} \sim 2^{2j+1}$$

| Level j | # Needlets |
|-----------|------------|
| 5 | 2108 |
| 6 | 8130 |
| 7 | 32642 |

Tab. # Needlets using Sym. spherical design


Womersley., 2015.

Localised needlet approximation




Localisation of needlets, order $j = 5$

Fig. needlet ψ_{jk} with a C^5 -needlet filter




Localisation of needlets, order $j = 9$

Fig. needlet ψ_{jk} with a C^5 -needlet filter





Towards localised needlet approximations




- $j = 4$, # all needlets = 498
- $j = 6$, # all needlets = 8130
- In cap, # needlets = 544

Towards localised needlet approximations



Towards localised needlet approximations



More ...

- All results can be generalised to \mathbb{S}^d , $d \geq 3$.

References

-  Q. T. Le Gia, I. H. Sloan, Y. G. Wang, R. S. Womersley.
Needlet approximation for isotropic random fields on the sphere. *Preprint*.
-  F. J. Narcowich, P. Petrushev, J. D. Ward.
Decomposition of Besov and Triebel-Lizorkin spaces on the sphere. *J. Funct. Anal.*, 238(2):530–564, 2006.
-  F. J. Narcowich, P. Petrushev, J. D. Ward.
Localized tight frames on spheres. *SIAM J. Math. Anal.*, 38(2):574–594, 2006.
-  I. H. Sloan, H. Wang.
On filtered polynomial approximation on the sphere. *arXiv:1509.03792 [math.CA]*.
-  Y. G. Wang, Q. T. Le Gia, I. H. Sloan, R. S. Womersley.
Fully discrete needlet approximation on the sphere. *arXiv:1502.05806 [math.NA]*. *Submitted*.
-  R. S. Womersley.
Efficient spherical designs with good geometric properties.
<http://web.maths.unsw.edu.au/~rsw/Sphere/EffSphDes/>, 2015.

The End

Thank you!