

FULLY DISCRETE NEEDLET APPROXIMATION ON THE SPHERE

Yu Guang Wang

School of Mathematics and Statistics
University of New South Wales, Australia

ICIAM 2015, Beijing


Joint with Quoc T. Le Gia, Ian H. Sloan and
Robert S. Womersley



UNSW
AUSTRALIA

A needlet on the sphere

A needlet on the sphere



Needlets

Needlets are **filtered** spherical polynomials
associated with **quadrature rules**.


Needlets

Needlets are **filtered** spherical polynomials
associated with **quadrature rules**.

For $j \geq 1$,


$$\psi_{jk}(\mathbf{x}) := \sqrt{w_{jk}} \sum_{\ell=0}^{\infty} h\left(\frac{\ell}{2^{j-1}}\right) (2\ell + 1) P_\ell(\mathbf{x} \cdot \mathbf{x}_{jk}).$$

C^5 -needlet filter h



Quadrature rule for needlets for ψ_{jk} , $j = 5$

Fig. A symmetric spherical 31-design



Nodes $\mathbf{x}_{jk} \in \mathbb{S}^2$, $k = 1, \dots, 498$.

Exact for polynomials P of degree up to $2^{5+1} - 1 = 31$

$$\int_{\mathbb{S}^2} P(\mathbf{x}) \, d\sigma_2(\mathbf{x}) = \frac{1}{N} \sum_{k=1}^N P(\mathbf{x}_{jk}).$$

Womersley, 2015.

What is a (semidiscrete) needlet approximation?

For a function f on \mathbb{S}^2 , $J = 0, 1, \dots$,

Definition (Semidiscrete needlet approximation)

$$V_J^{\text{need}}(f; \mathbf{x}) := \sum_{j=0}^J \sum_{k=1}^{N_j} (f, \psi_{jk})_{\mathbb{L}_2(\mathbb{S}^2)} \psi_{jk}(\mathbf{x}), \quad \mathbf{x} \in \mathbb{S}^2,$$

where

$$(f, \psi_{jk})_{\mathbb{L}_2(\mathbb{S}^2)} := \int_{\mathbb{S}^2} f(\mathbf{y}) \psi_{jk}(\mathbf{y}) d\sigma_2(\mathbf{y}).$$

The \mathbb{L}_p errors of needlet approximation

Narcowich et al., 2006.

Theorem

The \mathbb{L}_p error using the *needlet approximation* with sufficient smooth filter for $f \in \mathbb{W}_p^s(\mathbb{S}^2)$ and $1 \leq p \leq \infty$ and $s > 2/p$ has the convergence order 2^{-Js} .

If $h \in C^\kappa(\mathbb{R}_+)$, $\kappa \geq 3$,

$$\|f - V_J^{\text{need}}(f)\|_{\mathbb{L}_p(\mathbb{S}^2)} \leq c E_{\lfloor 2^{J-1} \rfloor}(f)_p \leq c 2^{-Js} \|f\|_{\mathbb{W}_p^s(\mathbb{S}^2)}.$$

The \mathbb{L}_p errors of needlet approximation

Narcowich et al., 2006.

Theorem

The \mathbb{L}_p error using the *needlet approximation* with sufficient smooth filter for $f \in \mathbb{W}_p^s(\mathbb{S}^2)$ and $1 \leq p \leq \infty$ and $s > 2/p$ has the convergence order 2^{-Js} .

If $h \in C^\kappa(\mathbb{R}_+)$, $\kappa \geq 3$,

$$\|f - V_J^{\text{need}}(f)\|_{\mathbb{L}_p(\mathbb{S}^2)} \leq c E_{\lfloor 2^{J-1} \rfloor}(f)_p \leq c 2^{-Js} \|f\|_{\mathbb{W}_p^s(\mathbb{S}^2)}.$$

$$\Rightarrow \sum_{j=0}^{\infty} \sum_{k=1}^{N_j} |(f, \psi_{jk})_{\mathbb{L}_2(\mathbb{S}^2)}|^2 = \|f\|_{\mathbb{L}_2(\mathbb{S}^2)}^2 \text{ (Tight frame).}$$

How to implement needlet approximations?

Given $J = 0, 1, \dots$

Discretisation quadrature $\mathcal{Q}_N := \{(W_i, \mathbf{y}_i) : i = 1, \dots, N\}$

Exact for polynomials of degree $3 \times 2^{J-1} - 1$

Need NOT be the same as needlet quadrature

Approximate needlet coefficient by quadrature \mathcal{Q}_N

$$\begin{aligned}(f, \psi_{jk})_{\mathbb{L}_2(\mathbb{S}^2)} &= \int_{\mathbb{S}^d} f(\mathbf{y}) \psi_{jk}(\mathbf{y}) d\sigma_2(\mathbf{y}) \\ &\approx \sum_{i=1}^N W_i f(\mathbf{y}_i) \psi_{jk}(\mathbf{y}_i) =: (f, \psi_{jk})_{\mathcal{Q}_N}\end{aligned}$$

How to implement needlet approximations?

Given $J = 0, 1, \dots$

Discretisation quadrature $\mathcal{Q}_N := \{(W_i, \mathbf{y}_i) : i = 1, \dots, N\}$

Exact for polynomials of degree $3 \times 2^{J-1} - 1$

Definition (Fully discrete needlet approximation)

$$V_{J,N}^{\text{need}}(f; \mathbf{x}) := \sum_{j=0}^J \sum_{k=1}^{N_j} (f, \psi_{jk})_{\mathcal{Q}_N} \psi_{jk}(\mathbf{x}), \quad \mathbf{x} \in \mathbb{S}^2.$$

$$V_J^{\text{need}}(f; \mathbf{x}) = \sum_{j=0}^J \sum_{k=1}^{N_j} (f, \psi_{jk})_{\mathbb{L}_2(\mathbb{S}^2)} \psi_{jk}(\mathbf{x}).$$

Discretisation does not lower the convergence rate!

Given $J = 0, 1, \dots$

Discretisation quadrature $\mathcal{Q}_N := \{(W_i, \mathbf{y}_i) : i = 1, \dots, N\}$

Exact for polynomials of degree $3 \times 2^{J-1} - 1$

Theorem

The *discrete needlet approximation with smooth filter and \mathcal{Q}_N* of a function f in $\mathbb{W}_p^s(\mathbb{S}^2)$ converges at rate 2^{-Js} in $\mathbb{L}_p(\mathbb{S}^2)$ -norm.


If $h \in C^\kappa(\mathbb{R}_+)$, $\kappa \geq 2$,

$$\|f - V_{J,N}^{\text{need}}(f)\|_{\mathbb{L}_p(\mathbb{S}^2)} \leq c 2^{-Js} \|f\|_{\mathbb{W}_p^s(\mathbb{S}^2)}, \quad f \in \mathbb{W}_p^s(\mathbb{S}^2).$$

Numerical examples: normalised Wendland functions

Chernih et al., 2014 & Wendland., 2001.

Fig. Test RBF function f_2



$$f_k(\mathbf{x}) := \sum_{i=1}^6 \phi_k(|\mathbf{z}_i - \mathbf{x}|), \quad k \geq 0,$$

where $\mathbf{z}_i = (\pm 1, 0, 0), (0, \pm 1, 0), (0, 0, \pm 1)$ and

$$\phi_k(r) := \tilde{\phi}_k\left(\frac{r}{\delta_k}\right), \quad r \in \mathbb{R},$$


$$\phi_k(r) \in \mathbb{W}_2^{k+3/2}(\mathbb{S}^2).$$

Le Gia et al., 2010.

Numerical examples: normalised Wendland functions

Chernih et al., 2014 & Wendland., 2001.

Fig. Test RBF function f_2



$$f_k(\mathbf{x}) := \sum_{i=1}^6 \phi_k(|\mathbf{z}_i - \mathbf{x}|), \quad k \geq 0,$$

where $\mathbf{z}_i = (\pm 1, 0, 0), (0, \pm 1, 0), (0, 0, \pm 1)$ and

$$\phi_k(r) := \tilde{\phi}_k\left(\frac{r}{\delta_k}\right), \quad r \in \mathbb{R},$$


$$\phi_k(r) \in \mathbb{W}_2^{k+3/2}(\mathbb{S}^2).$$

Le Gia et al., 2010.


For $k = 2$,

$$\tilde{\phi}_2(r) := (1-r)_+^6 (35r^2 + 18r + 3)/3.$$

Numerical examples support the theorems



(a) Semidiscrete



(b) Fully discrete

Key formula for proof

For $f \in C(\mathbb{S}^2)$ and $J \geq 0$,

$$V_{J,N}^{\text{need}}(f) = V_{2^{J-1},H,N}(f).$$

Key formula for proof

For $f \in C(\mathbb{S}^2)$ and $J \geq 0$,

$$V_{J,N}^{\text{need}}(f) = V_{2^{J-1},H,N}(f).$$

Filtered hyperinterpolation $V_{2^{J-1},H,N}(f)$ has optimal error order 2^{-Js} for $f \in \mathbb{W}_p^s(\mathbb{S}^2)$, $1 \leq p \leq \infty$ and $s > 2/p$.

Sloan & Wang., 2015.

Key formula for proof

For $f \in C(\mathbb{S}^2)$ and $J \geq 0$,

$$V_{J,N}^{\text{need}}(f) = V_{2^{J-1},H,N}(f).$$

Filtered hyperinterpolation $V_{2^{J-1},H,N}(f)$ has optimal error order 2^{-Js} for $f \in \mathbb{W}_p^s(\mathbb{S}^2)$, $1 \leq p \leq \infty$ and $s > 2/p$.

Sloan & Wang., 2015.

Using **ALL** needlets, the error of **discrete needlet approximation** follows from the error of filtered hyperinterpolation!

Connection to wavelets

For $f \in C(\mathbb{S}^2)$ and $J \geq 0$,

$$V_{J,N}^{\text{need}}(f) = \sum_{j=0}^J \mathcal{U}_{jN}(f).$$

Connection to wavelets

For $f \in C(\mathbb{S}^2)$ and $J \geq 0$,

$$V_{J,N}^{\text{need}}(f) = \sum_{j=0}^J \mathcal{U}_{jN}(f).$$

The **contribution** \mathcal{U}_{jN} is a discrete wavelet on \mathbb{S}^2 satisfying

$$\|\mathcal{U}_{jN}(f)\|_{\mathbb{L}_p(\mathbb{S}^d)} \leq c 2^{-js} \|f\|_{\mathbb{W}_p^s(\mathbb{S}^d)}, \quad f \in \mathbb{W}_p^s(\mathbb{S}^2).$$

Computational challenges in high orders

$$\# \text{ Needlets} \sim 2^{2j+1}$$


Level j	# Needlets
5	2108
6	8130
7	32642

Tab. # Needlets using Sym. spherical design

Womersley., 2015.


Localisation of needlets, order $j = 5$

Fig. needlet ψ_{jk} with a C^5 -needlet filter



Localisation of needlets, order $j = 9$

Fig. needlet ψ_{jk} with a C^5 -needlet filter



Localisation of needlets


Narcowich *et al.*, 2006.

For $h \in C^\kappa(\mathbb{R}_+)$ with $\kappa \geq 3$,

$$\frac{|\psi_{jk}(\mathbf{x})|}{\sqrt{w_{jk}}} \leq \frac{c_{d,h} 2^{jd}}{(1 + 2^j \operatorname{dist}(\mathbf{x}, \mathbf{x}_{jk}))^\kappa}, \quad \mathbf{x} \in \mathbb{S}^2,$$



where $\operatorname{dist}(\mathbf{x}, \mathbf{x}_{jk}) := \cos^{-1}(\mathbf{x} \cdot \mathbf{x}_{jk})$.

Towards localised needlet approximations





- $j = 4$, # all needlets = 498
- $j = 6$, # all needlets = 8130
- In cap, # needlets = 544

Towards localised needlet approximations




Towards localised needlet approximations



More ...

- All results can be generalised to \mathbb{S}^d , $d \geq 3$.

- All results can be generalised to \mathbb{S}^d , $d \geq 3$.
- Applications to random fields on the sphere. *Le Gia et al, 2015.*



Cosmology

References

-  F. Narcowich, P. Petrushev, and J. Ward.
Decomposition of Besov and Triebel-Lizorkin spaces on the sphere. *J. Funct. Anal.*, 238(2):530–564, 2006.
-  F. J. Narcowich, P. Petrushev, and J. D. Ward.
Localized tight frames on spheres. *SIAM J. Math. Anal.*, 38(2):574–594, 2006.
-  Q. T. Le Gia, I. H. Sloan, Y. G. Wang and R. S. Womersley.
Needlet approximation for isotropic random fields on the sphere. *Preprint*.
-  I. H. Sloan and H. Wang.
On filtered polynomial approximation on the sphere. *Preprint*.
-  I. H. Sloan and R. S. Womersley.
Filtered hyperinterpolation: a constructive polynomial approximation on the sphere. *Int. J. Geomath.*, 3(1):95–117, 2012.
-  Y. G. Wang, Q. T. Le Gia, I. H. Sloan and R. S. Womersley.
Fully discrete needlet approximation on the sphere. *Submitted*.
-  R. S. Womersley.
Efficient spherical designs with good geometric properties.
<http://web.maths.unsw.edu.au/~rsw/Sphere/EffSphDes/>, 2015.

The End

Thank you!