

IRF & needlets
oooooooooo

Semidiscrete
oooo

Approx errors
ooo

Fully discrete
oooooooooooooooooooo

Computation of isotropic random fields on spheres via needlet decomposition

Quoc T. Le Gia Ian H. Sloan Yu Guang Wang
Robert S. Womersley

School of Mathematics and Statistics
University of New South Wales, Australia

“Minimal Energy Point Sets, Lattices, and Designs”
ESI, Vienna, Oct. 2014

IRF & needlets

Semidiscrete oooo

Approx errors

Fully discrete
○○○○○○○○○○○○○○○○

Microwave background — random field on sphere




Fig. Temperature fluctuations in the microwave background

NASA / WMAP Science Team

Random field on sphere

- (Ω, \mathcal{F}, P) probability measure space
 - $\mathbb{S}^d := \{\mathbf{x} \in \mathbb{R}^{d+1} : |\mathbf{x}| = 1\}$

Definition

An $\mathcal{F} \otimes \mathcal{B}(\mathbb{S}^d)$ -measurable function $T : \Omega \times \mathbb{S}^d \rightarrow \mathbb{R}$ is said to be a *real-valued random field* on the sphere \mathbb{S}^d .

Random field on sphere

- (Ω, \mathcal{F}, P) probability measure space
 - $\mathbb{S}^d := \{\mathbf{x} \in \mathbb{R}^{d+1} : |\mathbf{x}| = 1\}$

Definition

An $\mathcal{F} \otimes \mathcal{B}(\mathbb{S}^d)$ -measurable function $T : \Omega \times \mathbb{S}^d \rightarrow \mathbb{R}$ is said to be a *real-valued random field* on the sphere \mathbb{S}^d .

- **Centered**: $\mathbb{E}[T] = 0$
 - Notation: $T(\omega, \mathbf{x})$ or $T(\mathbf{x})$ or $T(\omega)$

IRF & needlets
○○●○○○○○

Semidiscrete

Approx errors

Needlets

Needlets are **filtered** spherical polynomials
associated with **quadrature rules**.


IRF & needlets
○○○●○○○○

Semidiscrete

Approx errors

Needlet $\psi_{jk}, j = 9, k = 10$

Fig. ψ_{jk} with a $C^5(\mathbb{R}_+)$ -filter, tensor product (Gauss-Legendre \times Equal)



Quadrature rule

- Quadrature rule $\{w_k, \mathbf{x}_k : k = 1, \dots, N\}$
 - Weights $w_k > 0$
 - Nodes $\mathbf{x}_k \in \mathbb{S}^d$
 - Exact for polynomials of degree up to ℓ

$$\int_{\mathbb{S}^d} P(\mathbf{x}) d\sigma_d(\mathbf{x}) = \sum_{k=1}^N w_k P(\mathbf{x}_k), \quad P \in \mathbb{P}_\ell(\mathbb{S}^d),$$

$\mathbb{P}_\ell(\mathbb{S}^d)$ is the space of all polynomials of degree $\leq \ell$ on \mathbb{S}^d .

Needlets

For order $j = 0, 1, \dots$,

- **Needlet Quadrature Rule** $\{w_{jk}, \mathbf{x}_{jk} : k = 1, \dots, N_j\}$ exact for polynomials of degree 2^{j+1}
 - **Needlets** ψ_{jk} , $k = 1, \dots, N_j$ and **needlet filter** h


$$\psi_{jk}(\mathbf{x}) := \begin{cases} \sqrt{w_{0k}}, & \text{if } j = 0, \\ \sqrt{w_{jk}} \sum_{\ell=0}^{\infty} h\left(\frac{\ell}{2^{j-1}}\right) Z(d, \ell) P_\ell^{(d)}(\mathbf{x} \cdot \mathbf{x}_{jk}), & \text{if } j \geq 1, \end{cases}$$

$P_\ell^{(d)}$ is the normalised ($P_\ell^{(d)}(1) = 1$) Gegenbauer polynomial.

[Narcowich, Petrushev, Ward. *SIAM J. Math. Anal.*, *J. Funct. Anal.* 2006]

C^5 -needlet filter

C^5 -needlet filter



Needlet filter h

$$0 \leq h(t) \leq 1$$

$$\text{supp } h \in [1/2, 2]$$

$$[h(t)]^2 + [h(2t)]^2 = 1,$$

$$t \in [1/2, 1]$$

$$h \in C^\kappa(\mathbb{R}_+), \kappa \geq d + 1$$

IRF & needlets


Semidiscrete

Approx errors

Localised zonal polynomial: needlet ψ_{jk} , $j = 9, k = 10$

Estimate

Fig. ψ_{jk} with a C^5 -needlet filter, tensor rule (Gauss-Legendre \times Equal)



IRF & needlets
○○○○○○○●

Semidiscrete
○○○○

Approx errors
○○○

Fully discrete
○○○○○○○○○○○○○○○○

Dubai Tower — A needlet with lower resolution



IRF & needlets
oooooooooooo

Semidiscrete

Approx errors

Fully discrete
○○○○○○○○○○○○○○○○

Karhunen-Loève expansion

Let $\{Y_{\ell,m} : m = 1, \dots, Z(d, \ell), \ell = 0, 1, \dots\}$ be an \mathbb{L}_2 -orthonormal basis for $\mathbb{L}_2(\mathbb{S}^d)$.

The Karhunen-Loève expansion of a random field T on \mathbb{S}^d is

$$T(\omega, \mathbf{x}) \approx \sum_{\ell=0}^{\infty} \sum_{m=1}^{Z(d,\ell)} (T(\omega), Y_{\ell,m})_{\mathbb{L}_2(\mathbb{S}^d)} Y_{\ell,m}(\mathbf{x}), \quad \mathbf{x} \in \mathbb{S}^d, \omega \in \Omega.$$

Karhunen-Loève expansion

Let $\{Y_{\ell,m} : m = 1, \dots, Z(d, \ell), \ell = 0, 1, \dots\}$ be an \mathbb{L}_2 -orthonormal basis for $\mathbb{L}_2(\mathbb{S}^d)$.

The Karhunen-Loëve expansion of a random field T on \mathbb{S}^d is

$$T(\omega, \mathbf{x}) \approx \sum_{\ell=0}^{\infty} \sum_{m=1}^{Z(d,\ell)} (T(\omega), Y_{\ell,m})_{\mathbb{L}_2(\mathbb{S}^d)} Y_{\ell,m}(\mathbf{x}), \quad \mathbf{x} \in \mathbb{S}^d, \omega \in \Omega.$$

For $d \geq 2$, the Karhunen-Loève expansion converges in $\mathbb{L}_2(\Omega \times \mathbb{S}^d, P \otimes \sigma_d)$ and also in $\mathbb{L}_2(\Omega, P)$.

E.g. [Marinucci, Peccati. Cambridge University Press, Cambridge, 2011]

IRF & needlets

Semidiscrete



Approx errors

Fully discrete
○○○○○○○○○○○○○○○○

Tight frame

Proposition (Almost orthogonal)

The needlets $\psi_{jk}, \psi_{j'k'}$ are \mathbb{L}_2 -orthogonal if $|j - j'| \geq 2$:

$$(\psi_{jk}, \psi_{j'k'})_{\mathbb{L}_2(\mathbb{S}^d)} = 0.$$

IRF & needlets
oooooooooooo

Semidiscrete



Approx errors

Fully discrete
○○○○○○○○○○○○○○○○

Tight frame

Proposition (Almost orthogonal)

The needlets ψ_{jk} , $\psi_{j'k'}$ are \mathbb{L}_2 -orthogonal if $|j - j'| \geq 2$:

$$(\psi_{jk}, \psi_{j'k'})_{\mathbb{L}_2(\mathbb{S}^d)} = 0.$$

Proposition (Parseval identity)

$$\sum_{j=0}^{\infty} \sum_{k=1}^{N_j} |(f, \psi_{jk})_{\mathbb{L}_2(\mathbb{S}^d)}|^2 = \|f\|_{\mathbb{L}_2(\mathbb{S}^d)}, \quad f \in \mathbb{L}_2(\mathbb{S}^d).$$

[Narcowich, Petrushev, Ward. *SIAM J. Math. Anal.*, *J. Funct. Anal.* 2006]

IRF & needlets
oooooooooooo

Semidiscrete



Approx errors

Fully discrete
○○○○○○○○○○○○○○○○

Semidiscrete needlet decomposition

For a random field T on \mathbb{S}^d ,

Definition (Needlet decomposition)

$$T(\omega, \mathbf{x}) \approx \sum_{j=0}^{\infty} \sum_{k=1}^{N_j} (T(\omega), \psi_{jk})_{\mathbb{L}_2(\mathbb{S}^d)} \psi_{jk}(\mathbf{x}), \quad \mathbf{x} \in \mathbb{S}^d, \quad \omega \in \Omega.$$

Mean convergence

Theorem

The needlet decomposition converges in $\mathbb{L}_p(\Omega \times \mathbb{S}^d, P \otimes \sigma_d)$,
 $1 \leq p \leq \infty$ for any $[p]$ -weakly isotropic random field.

$$\lim_{L \rightarrow +\infty} \mathbb{E} \left[\left\| T - \sum_{2^j \leq L} \sum_{k=1}^{N_j} (T, \psi_{jk})_{\mathbb{L}_2(\mathbb{S}^d)} \psi_{jk} \right\|_p^p \right] = 0, \quad 1 \leq p < \infty$$

and

$$\lim_{L \rightarrow +\infty} \mathbb{E} \left[\left\| T - \sum_{2^j \leq L} \sum_{k=1}^{N_j} (T, \psi_{jk})_{\mathbb{L}_2(\mathbb{S}^d)} \psi_{jk} \right\|_\infty \right] = 0, \quad p = \infty.$$

weak isotropy

IRF & needlets
oooooooooo

Semidiscrete
oooo

Approx errors
●○○

Fully discrete
oooooooooooooooooooo

Angular power spectrum

For T a centered two-weakly isotropic random field,
 $\mathbb{E}[T(\mathbf{x})T(\mathbf{y})]$ is a zonal function on \mathbb{S}^d .

Angular power spectrum $A_\ell = A_\ell^{(d)}$ is defined by:

$$\mathbb{E}[T(\mathbf{x})T(\mathbf{y})] = \sum_{\ell=0}^{\infty} A_\ell^{(d)} Z(d, \ell) P_\ell^{(d)}(\mathbf{x} \cdot \mathbf{y}).$$

IRF & needlets



Semidiscrete oooo

Approx errors

Fully discrete
○○○○○○○○○○○○○○○○

Smoothness vs Angular power spectrum

T two-weakly isotropic, satisfying

$$\tilde{c}_{d,s} := \sum_{\ell=0}^{\infty} A_\ell \ell^{2s+d-1} < +\infty.$$

Theorem (Kolmogorov-Chentsov continuation for sphere)

$T(\omega)$ is in Sobolev space $\mathbb{W}_2^s(\mathbb{S}^d)$ for almost every $\omega \in \Omega$.

For \mathbb{S}^2 , [Lang, Schwab. arXiv 2013].

IRF & needlets



Semidiscrete

Approx errors

Fully discrete
○○○○○○○○○○○○○○○○

Mean error of semidiscrete needlet approx.

Let $d \geq 2$, $s > d/2$, $\kappa \geq d + 1$ ($h \in C^\kappa(\mathbb{R}_+)$) and T be centered two-weakly isotropic satisfying

$$\tilde{c}_{d,s} = \sum_{\ell=0}^{\infty} A_\ell \ell^{2s+d-1} < +\infty.$$

Theorem

$$\mathbb{E} \left[\left\| T - \sum_{2^j \leq L} \sum_{k=1}^{N_j} (T, \psi_{jk})_{\mathbb{L}_2(\mathbb{S}^d)} \psi_{jk} \right\|_2^2 \right] \leq c \tilde{c}_{d,s} L^{-2s},$$

where the constant c depends only on d, s and needlet filter h .

Fully discrete needlet approximation

Given a positive integer L

- *Discretisation quadrature* $\mathcal{Q}_N := \{(W_i, \mathbf{y}_i) : i = 1, \dots, N\}$
 - Exact for polynomials of degree $3L$
 - Need NOT be the same as needlet quadrature
 - Approximate needlet coefficient by quadrature \mathcal{Q}_N

$$\begin{aligned} (T, \psi_{jk})_{\mathbb{L}_2(\mathbb{S}^d)} &= \int_{\mathbb{S}^d} T(\mathbf{y}) \psi_{jk}(\mathbf{y}) d\sigma_d(\mathbf{y}) \\ &\approx \sum_{i=1}^N W_i T(\mathbf{y}_i) \psi_{jk}(\mathbf{y}_i) =: (T, \psi_{jk})_{\mathcal{Q}_N} \end{aligned}$$

IRF & needlets
oooooooooooo

Semidiscrete

Approx errors

Fully discrete

Fully discrete needlet approximation (continued)

(Fully) discrete needlet approximation of degree L (or order η) for a random field T is

$$\sum_{2^j < L} \sum_{k=1}^{N_j} (T(\omega), \psi_{jk})_{\mathcal{Q}_N} \psi_{jk}(\mathbf{x}), \quad \mathbf{x} \in \mathbb{S}^d, \quad \omega \in \Omega$$

Discretisation does not lower the convergence rate

Let $d \geq 2$, $s > d/2$, $\kappa \geq d + 1$ and T be centered two-weakly isotropic satisfying

$$\tilde{c}_{d,s} = \sum_{\ell=0}^{\infty} A_\ell \ell^{2s+d-1} < +\infty.$$

Theorem (Mean error of discrete needlet approx.)

$$\mathbb{E} \left[\left\| T - \sum_{2^j \leq L} \sum_{k=1}^{N_j} (T, \psi_{jk})_{\mathcal{Q}_N} \psi_{jk} \right\|_2^2 \right] \leq c \tilde{c}_{d,s} L^{-2s},$$

where the constant c depends only on d , s and h .

Truncation error for a GRF, $A_\ell = 1/\ell^{2s+2}$, $s = 1, d = 2$

Mean error of dis. needlet approx., $L \leq 2^6$, $s=1$, C^5 filter




Fig. Order ≤ 6 , Sample = 100, Sym. Spherical design

IRF & needlets
○○○○○○○○



Semidiscrete
○○○○

Approx errors
○○○

Fully discrete
○○○●○○○○○○○○○○○○

GRF (left) by discrete needlet approx.(right), $A_\ell = 1/\ell^{2s+2}$, $s = 1$ [Skip](#)

Sample 1, $L = 64$



Evaluated at *generalized spiral points*
[Rakhmanov, Saff, Zhou. Math. Res. Lett. 1994]

IRF & needlets
○○○○○○○○



Semidiscrete
○○○○

Approx errors
○○○

Fully discrete
○○○○●○○○○○○○○○○○

GRF (left) by discrete needlet approx.(right), $A_\ell = 1/\ell^{2s+2}$, $s = 1$ [Skip](#)

Sample 2, $L = 64$



Evaluated at *generalized spiral points*
[Rakhmanov, Saff, Zhou. Math. Res. Lett. 1994]

IRF & needlets
○○○○○○○○



Semidiscrete
○○○○

Approx errors
○○○

Fully discrete
○○○○○●○○○○○○○○○○

GRF (left) by discrete needlet approx.(right), $A_\ell = 1/\ell^{2s+2}$, $s = 1$ [Skip](#)

Sample 3, $L = 64$



Evaluated at *generalized spiral points*

[Rakhmanov, Saff, Zhou. Math. Res. Lett. 1994]

IRF & needlets
○○○○○○○○



Semidiscrete
○○○○

Approx errors
○○○

Fully discrete
○○○○○○○●○○○○○○○○

GRF (left) by discrete needlet approx.(right), $A_\ell = 1/\ell^{2s+2}$, $s = 1$ [Skip](#)

Sample 4, $L = 64$



Evaluated at *generalized spiral points*
[Rakhmanov, Saff, Zhou. Math. Res. Lett. 1994]

IRF & needlets
○○○○○○○○

Semidiscrete
○○○○

Approx errors
○○○

Fully discrete
○○○○○○○○●○○○○○○

Examples of quadrature rules

Point set	Quadrature	
	Needlet	Discretisation
Gauss-Legendre Tensor	$j \geq 0$	$L \geq 0$
Sym. Spherical design	$2^{j+1} \leq 251^*$	$3L \leq 251$

* Private communication with Rob Womersley

IRF & needlets
○○○○○○○○Semi-discrete
○○○○Approx errors
○○○Fully discrete
○○○○○○○○●○○○○○

Challenge in computation

$$\text{Number of Needlets} \sim \frac{(2 * 2^j)^2}{2}$$

$$\text{Number of nodes for discretisation} \sim \frac{(3 * 2^j)^2}{2}$$

Level j	# Needlets	# QH
0	6	8
2	38	80
4	530	1178
6	8258	18530
8	131330	295298

Tab. Number of nodes required — Sym. spherical design

Filtered hyperinterpolation

With *hyperinterpolation filter* H and discretisation quadrature \mathcal{Q}_N

Definition (**Filtered hyperinterpolation**)

$$V_{L,H,N}(f; \mathbf{x}) := \sum_{i=1}^N W_i f(\mathbf{y}_i) v_{L,H}(\mathbf{y}_i \cdot \mathbf{x})$$

Filtered kernel

$$v_{L,H}(t) := \sum_{\ell=0}^{\infty} H\left(\frac{\ell}{L}\right) Z(d, \ell) P_\ell^{(d)}(t), \quad -1 \leq t \leq 1$$

[Le Gia, Mhaskar. *SIAM J. Numer. Anal.* 2008]

[Sloan, Womersley. *GEM Int. J. Geomath.* 2012]

[Ivanov, Petrushev. *Adv. Comput. Math.* 2014]

IRF & needlets
○○○○○○○○○○


Semidiscrete
○○○○

Approx errors
○○○

Fully discrete
○○○○○○○○○○○●○○○

An example: H in $C^5(\mathbb{R}_+)$

Degree 11 polynomial filter, in C^5



IRF & needlets
○○○○○○○○

Semidiscrete
○○○○

Approx errors
○○○

Fully discrete
○○○○○○○○○○○●○○

Needlets and filtered kernels

$$H(t) := \begin{cases} 1, & 0 \leq t \leq 1, \\ [h(t)]^2, & t > 1, \end{cases}$$

h : Needlet filter, $h(t) = 0$, $t \geq 2$.

Theorem (Filtered kernel decomposed via needlets)

$$\sum_{k=1}^{N_j} \psi_{jk}(\mathbf{x}) \psi_{jk}(\mathbf{y}) = v_{2^{j-1}, h^2}^{(d)}(\mathbf{x} \cdot \mathbf{y}), \quad j \geq 0,$$

$$\sum_{j=0}^{\eta} \sum_{k=1}^{N_j} \psi_{jk}(\mathbf{x}) \psi_{jk}(\mathbf{y}) = v_{2^{\eta-1}, H}^{(d)}(\mathbf{x} \cdot \mathbf{y}), \quad \eta \geq 0,$$

$$\eta = \lfloor \log_2(L) \rfloor.$$

IRF & needlets
○○○○○○○○

Semidiscrete
○○○○

Approx errors
○○○

Fully discrete
○○○○○○○○○○○○●○○

Discrete needlet approx. and filtered hyperinterpolation

$$\begin{aligned} V_{2^{\eta-1}, H, N}^{(d)}(f; \mathbf{x}) &= \sum_{i=1}^N W_i f(\mathbf{y}_i) v_{2^{\eta-1}, H}^{(d)}(\mathbf{y}_i \cdot \mathbf{x}) \\ &= \sum_{i=1}^N W_i f(\mathbf{y}_i) \sum_{j=0}^{\eta} \sum_{k=1}^{N_j} \psi_{jk}(\mathbf{y}_i) \psi_{jk}(\mathbf{x}) \\ &= \sum_{j=0}^{\eta} \sum_{k=1}^{N_j} (f, \psi_{jk})_{\mathcal{Q}_N} \psi_{jk}(\mathbf{x}). \end{aligned}$$

IRF & needlets
oooooooo

Semidiscrete
oooo

Approx errors
ooo

Fully discrete
oooooooooooo●○

Summary

- L_p convergence of needlet decomposition for $\lceil p \rceil$ -weakly isotropic random fields on \mathbb{S}^d
- Semidiscrete and fully discrete needlet approximations have same truncation errors
- Results proved for \mathbb{S}^d , $d \geq 2$

IRF & needlets
oooooooooo

Semidiscrete
oooo

Approx errors
ooo

Fully discrete
oooooooooooooooooooo●

End

Thanks!

Strong isotropy

We say T is *strongly isotropic* if for any $\mathbf{x} \in \mathbb{S}^d$ and for any rotation $\rho \in \mathrm{SO}(d+1)$, $T(\mathbf{x})$ and $T(\rho\mathbf{x})$ have the same law.

Weak isotropy

T is *two-weakly isotropic* if for all $\mathbf{x} \in \mathbb{S}^d$,

$$\mathbb{E} [|T(\mathbf{x})|^2] < +\infty$$


and if for $\mathbf{x}_1, \mathbf{x}_2 \in \mathbb{S}^d$ and for any rotation $\rho \in \mathrm{SO}(d+1)$,

$$\mathbb{E} [T(\mathbf{x}_1)] = \mathbb{E} [T(\rho \mathbf{x}_1)]$$

$$\mathbb{E} [T(\mathbf{x}_1)T(\mathbf{x}_2)] = \mathbb{E} [T(\rho \mathbf{x}_1)T(\rho \mathbf{x}_2)] .$$

Isotropy — Cosmological microwave background radiation

◀ Back



Gaussian random fields on spheres

We say T is a *Gaussian random field* (GRF) on \mathbb{S}^d if the vector $(T(\mathbf{x}_1), \dots, T(\mathbf{x}_k))$ follows the multivariate Gaussian distribution for all $k \geq 1$ and $\mathbf{x}_1, \dots, \mathbf{x}_k \in \mathbb{S}^d$.

Gaussian random fields on spheres

We say T is a *Gaussian random field* (GRF) on \mathbb{S}^d if the vector $(T(\mathbf{x}_1), \dots, T(\mathbf{x}_k))$ follows the multivariate Gaussian distribution for all $k \geq 1$ and $\mathbf{x}_1, \dots, \mathbf{x}_k \in \mathbb{S}^d$.

Remark

- (i) If the law of T is determined by its moments, then T is ∞ -weakly isotropic if and only if T is strongly isotropic and $\mathbb{E}[|T(\mathbf{x})|^\nu] < +\infty$ for every integer $\nu \geq 2$ and each $\mathbf{x} \in \mathbb{S}^d$;
- (ii) Let T be a GRF on \mathbb{S}^d . Then, T is strongly isotropic if and only if T is 2-weakly isotropic.

Needlet filter

Given $\kappa \geq 0$, a *needlet filter* h is a real function on \mathbb{R}_+ satisfying

- (i) (Compact support.) $h \in C^\kappa(\mathbb{R}_+)$, $\text{supp } h \subset [1/2, 2]$;
- (ii) (Partition of unity.) For all $t \geq 1$,

$$\sum_{j=0}^{\infty} \left[h\left(\frac{t}{2^j}\right) \right]^2 = 1.$$

Localisation of needlets

Let $d \geq 2$ and $h \in C^\kappa(\mathbb{R}_+)$ with $\kappa \geq 1$.

$$|\psi_{jk}(\mathbf{x})| \leq \frac{c_{d,h} 2^{jd}}{(1 + 2^j \operatorname{dist}(\mathbf{x}, \mathbf{x}_{jk}))^\kappa}, \quad \mathbf{x} \in \mathbb{S}^d.$$

[Mhaskar. *J. Approx. Theory* 2004]

[Petrushev, Xu. *J. Fourier Anal. Appl.* 2005]

[Narcowich, Petrushev, Ward. *SIAM J. Math. Anal.*, *J. Funct. Anal.* 2006]

◀ Back