

A Training-Free Quantum-Walk Scanner for Cryptic Allosteric Sites in Undruggable Proteins

Global Quantum + AI Challenge 2026 — Cleveland Clinic Enterprise Challenge · Phase I Concept Proposal

1. Executive Summary

More than 85% of disease-driving proteins are considered *undruggable*: the only viable therapeutic strategy is allosteric modulation—finding a hidden regulatory pocket on a distal surface. We propose a **training-free Continuous-Time Quantum Walk (CTQW) scanner**. The contact topology and evolutionary conservation of an apo protein are encoded into a sparse Hermitian Hamiltonian; we compute its **average mixing matrix** as the inter-residue quantum connectivity matrix, and rank distal cryptic pockets by their **dynamical connectivity to the active site**. The core quantity is a **deterministic spectral property** of the Hamiltonian—**no supervised training**—so there is no small- n or generalization problem, and the three target proteins serve as pure blind validation.

Quantum positioning and feasibility. Predictions are produced in a *quantum-inspired* manner (classically exact simulation of the CTQW); the quantum-hardware track demonstrates a credible path via a gate-based CTQW circuit on coarse-grained super-nodes (K nodes, AWS Braket / IBM) together with noise resilience (QSW dephasing sweep). This satisfies the challenge’s requirement that gate-based, quantum-inspired, or hybrid approaches are all acceptable provided a near-term hardware execution path exists. The quantum character comes from **coherent interference** (e^{-iHt}), which we substantiate through its discriminative gain over classical diffusion e^A .

2. Method Overview




Figure 1. End-to-end pipeline. The main spine (green/blue) is the quantum-inspired, classically exact prediction path; the dashed box on the right is the quantum-hardware demonstration track. Steps ④+⑤ decouple the

dimension and qubit count from protein length.

3. Methodology

3.1 Graph Construction and Hamiltonian Encoding

The protein is represented as a **residue-level** graph (nodes = residues, mapping directly onto the deliverable residue indices while suppressing qubit demand). The Hamiltonian H is sparse and Hermitian:

$$H_{ij} = w_{ij} e^{i\varphi_{ij}} \quad (i \neq j); \quad H_{ii} = c_i.$$

- **Off-diagonal (hopping) w_{ij} :** residue contact topology, taken from the GNM contact network—**nodes = residues (represented by $C\alpha$)**, **edges = $C\alpha$ – $C\alpha$ distance < cutoff (7.3 Å)**; its Kirchhoff matrix is $\Gamma = D - A$ (A = contact adjacency, D = degree diagonal, $A_{ij} = 1$ if in contact). w_{ij} takes the contacts of A (or a finer heavy-atom contact count at 4.5 Å). Naturally sparse (degree ~ 6 – 10), which caps circuit cost. Note: only the GNM contact graph is borrowed as the CTQW topology; no GNM mode decomposition is performed.
- **Diagonal (on-site energy) c_i :** per-residue evolutionary conservation (ESM-2 / MSA). This injects a functional prior through a sparse diagonal, **without densifying H and without breaking Hermiticity**—the key differentiator from purely topological prior work.
- **Chiral phase φ_{ij} (optional):** given by the projection of the ANM lowest-frequency mode along edge (i, j) , breaking time-reversal symmetry to yield an **asymmetric, directional** connectivity matrix.

A_{dist} (pure geometric distance) is removed as highly redundant with A_{mech} ; its geometric content is reused as the chiral-phase source. Each channel is z-score normalized; any asymmetric input is symmetrized as $(A + A^T)/2$. States use amplitude/log encoding (K nodes $\rightarrow \lceil \log_2 K \rceil$ qubits).

3.2 Hamiltonian Validity and Hermiticity

Challenge compliance. H is constructed *ab initio* entirely from the apo crystal structure and sequence—contact topology from apo coordinates, diagonal conservation from sequence/ESM-2, and the optional phase from the eigenvectors of the analytic ANM Hessian. **None of these is a molecular-dynamics trajectory** (ANM is the analytic eigenproblem of an elastic network, not MD sampling), satisfying the challenge rules that classical MD may not be used and that the method must proceed from topology under the elastic-network hypothesis.

Hermiticity (necessary and sufficient for a valid CTQW). A valid quantum walk requires $H = H^\dagger$, which guarantees that $U(t) = e^{-iHt}$ is unitary, the eigenvalues are real, and probability is conserved. A channel-by-channel check proves the fused H remains Hermitian:

- **Diagonal (conservation):** $c_i \in \mathbb{R} \Rightarrow$ real diagonal, so $H_{ii} = \overline{H_{ii}}$ automatically.
- **Off-diagonal magnitude:** contacts are symmetric $w_{ij} = w_{ji}$ (enforced post-construction via $(A + A^T)/2$); z-score normalization preserves realness and symmetry.
- **Chiral phase:** taking the phase **antisymmetric**, $\varphi_{ji} = -\varphi_{ij}$, gives $H_{ji} = w_{ij} e^{-i\varphi_{ij}} = \overline{H_{ij}}$ identically.
 $\Rightarrow H = H^\dagger : \quad H_{ij} = w_{ij} e^{i\varphi_{ij}}, \quad H_{ji} = w_{ij} e^{-i\varphi_{ij}} = \overline{H_{ij}}, \quad H_{ii} = c_i \in \mathbb{R}.$

Key distinction: once the chiral phase is added, H is **complex Hermitian but not symmetric**—still a valid CTQW, yet it yields a **directional** connectivity $\hat{M}[i \rightarrow j] \neq \hat{M}[j \rightarrow i]$ that symmetric classical GNM/communicability cannot express. With $\varphi \equiv 0$ it reduces to the standard CTQW of a real symmetric H . Only the phase accumulated around **cycles** (magnetic flux) is physical, so a gauge fixing first removes trivial phases absorbable by a diagonal unitary, retaining only the directional signal. The fusion routine `assemble_hamiltonian` ends with the assertion $\|H - H^\dagger\|_F < 10^{-10}$ as a pre-flight check.

3.3 Active-Site-Anchored Krylov Coarse-Graining (Fixed-Dimension Reduction)

Goal: decouple the dimension and qubit count from protein length while exactly preserving transport “sourced at the active site.” Key insight: ranking needs only the **source**→**all** transport vector $\langle j | e^{-iHt} | \psi_S \rangle$, and this dynamics is spanned exactly by the Krylov subspace generated from $|\psi_S\rangle$. Define the source state $|\psi_S\rangle = \sum_{i \in S} a_i |i\rangle$ (normalized); the order- K Krylov subspace is

$$\mathcal{K}_K(H, \psi_S) = \text{span}\{ |\psi_S\rangle, H|\psi_S\rangle, H^2|\psi_S\rangle, \dots, H^{K-1}|\psi_S\rangle \}.$$

The Lanczos iteration orthonormalizes it into a basis $Q = [q_1 \cdots q_K]$ ($q_1 = |\psi_S\rangle$, $N \times K$), projecting H onto a **tridiagonal** $H_K = Q^\dagger H Q$ (diagonal α_r , off-diagonal β_r):

$$\beta_r |q_{r+1}\rangle = H|q_r\rangle - \alpha_r |q_r\rangle - \beta_{r-1} |q_{r-1}\rangle, \quad \alpha_r = \langle q_r | H | q_r \rangle.$$

Because $Q^\dagger |\psi_S\rangle = e_1$, the transport from the source to any residue is **exactly preserved** within K dimensions:

$$\langle j | e^{-iHt} | \psi_S \rangle = [Q]_{j,:} e^{-iH_K t} e_1.$$

Steps: (1) build the sparse H ; (2) form $|\psi_S\rangle$ from the annotated S ; (3) run K Lanczos steps to obtain H_K, Q ($K = 32\text{--}64$; hardware demo $K = 8\text{--}16 \rightarrow 3\text{--}4$ qubits); (4) tridiagonal $H_K \rightarrow$ shallow Trotter circuit $e^{-iH_K t}$; (5) after readout, project back to residues via Q . The delivered $N \times N$ $\hat{M}$ uses the classically exact eigendecomposition (single-particle CTQW lives in N dimensions, poly-time); ranking uses the fixed- K Krylov path, which is the part run on hardware. This preserves “source transport,” unlike generic spectral clustering (which retains only generic modes).

3.4 CTQW and Readout: the Average Mixing Matrix

The propagator of the continuous-time quantum walk is $U(t) = e^{-iHt}$. Its long-time average mixing matrix is a deterministic spectral object:

$$\hat{M}_{ij} = \lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T |\langle j | e^{-iHt} | i \rangle|^2 dt = \left(\sum_r E_r \circ E_r \right)_{ij},$$

where E_r are the eigenprojectors of H . $\hat{M}$ is symmetric, doubly stochastic, and has **no free time parameter and needs no training**—it *is* the delivered $N \times N$ quantum connectivity matrix. The score of residue j with respect to the active site is

$$\text{score}(j) = \sum_{i \in S} w_i \hat{M}_{ij}.$$

Quantum character: $\hat{M}$ is determined by coherent interference and is mathematically distinct from classical diffusion e^A (Estrada communicability); we substantiate the interference gain through the discriminative delta of e^{-iAt} vs e^A .

3.5 Ranking Metric and Hit-list Selection

Since $\hat{M}$ is doubly stochastic, raw connectivity is confounded by centrality (hub residues score high by construction). The ranking metric must be normalized so that it reflects **specific** connectivity to the active site above background—exactly the quantity tested by the challenge’s primary objective, extractable directly from $\hat{M}$:

① raw connectivity:	$s(j) = \sum_{i \in S} w_i \hat{M}_{ij}$
② specificity (deconfounds degree):	$\rho(j) = s(j) / \sum_k \hat{M}_{kj}$
③ ★ primary metric (background-standardized):	$z(j) = (\rho(j) - \mu_{\text{bg}}) / \sigma_{\text{bg}}$

$\mu_{\text{bg}}, \sigma_{\text{bg}}$ are taken from background residues (non-active, non-neighbor surface residues + non-functional pockets). $z(j)$ is precisely the score used by the statistical test (allosteric vs. background): the ranking metric is isomorphic to the scoring criterion, avoiding the degenerate “rank the hubs” solution.

Filtering. Exclude S and its first-shell neighbors; require **distal** residues (Ca--Ca distance to $S > 12 \text{ \AA}$). **Top-5 (primary ranking, residue level): purely by the primary metric $z(j)$ — no pocketness is applied and druggability is not a hard gate.** After filtering, take the 5 residues with the highest $z(j)$ (spatially adjacent ones first merged into a single hit), which are the delivered Top-5.

Supplementary note (druggability; soft, not a filter). For pocket-bearing targets such as KRAS/ABL/myosin, high- $z(j)$ residues may be spatially clustered into candidate pockets P and *annotated* with pocketness as a druggability tag: $\text{Score}(P) = (\text{mean or 90th-pct})_{j \in P} z(j) \times \text{pocketness}(P)$. This is only a post-ranking druggability annotation and **does not change the primary Top-5**: cryptic allosteric pockets are typically closed in the apo

state and missed by fpocket, so a hard pocketness gate would defeat the very purpose of “finding hidden pockets.” c-Myc (an IDP with no static pocket) explicitly disables pocketness and is **connectivity-only**.

Optional quantum-increment metric (for the method report, not the primary ranking): $\Delta(j) = \hat{M}_{\text{CTQW}}[S, j] - C_{e^A}[S, j]$ isolates the interference contribution; with a chiral H we instead use the directed $\hat{M}[i \rightarrow j]$ to represent “signal flowing out of the active site.”

3.6 Dual-Track Execution: Classical-Exact + Quantum Hardware

Prediction track (quantum-inspired): full-resolution eigendecomposition of the sparse H ($O(N^3)$, trivial for $N \sim 300$ –1000) computes $\hat{M}$ exactly, producing a deterministic, residue-level connectivity matrix and hit list. **Hardware track (credible path):** coarse-grain to $K = 8$ –16, and with amplitude encoding realize e^{-iHt} on AWS Braket / IBM via qDRIFT or sparse-Trotter, verifying that the quantum $\hat{M}$ matches the classical one; a QSW (Lindblad) sweep over the dephasing p probes noise ($p \rightarrow 1$ = classical baseline, the ENAQT optimum = noise resilience).

3.7 Method Functions

Function	Input $\rightarrow$ Output	Description	Packages
build_residue_graph	apo.pdb $\rightarrow (E, c)$	parse apo; build heavy-atom contact edges E (w_{ij}); compute per-residue conservation c_i	BioPython, ProDy, fair-esm
assemble_hamiltonian	$(E, c, \Phi) \rightarrow H$	$H_{ij} = w_{ij}e^{i\varphi}$, $H_{ii} = c_i$; symmetrize, z-score; sparse Hermitian	NumPy/SciPy
coarse_grain	$(H, S, K) \rightarrow (H_K, Q)$	active-site-anchored Krylov/Lanczos reduction to fixed K ; preserves transport dynamics	SciPy
average_mixing_matrix	$H \rightarrow \hat{M}$	eigendecomposition $\rightarrow \hat{M} = \sum_r E_r \circ E_r$; deterministic, no training; = connectivity matrix	NumPy
connectivity_score	$(\hat{M}, S, w) \rightarrow s(j)$	dynamical connectivity of residue j to the active-site set S	NumPy
select_hits	$(z, \text{coords}, S) \rightarrow \text{Top-5}$	exclude S +neighbors; distal filter; rank by $z(j)$; (optional) cluster + annotate pocketness	fpocket, SciPy, NetworkX
ctqw_circuit (HW)	$H_K \rightarrow \hat{U}(t)$	$\lceil \log_2 K \rceil$ -qubit gate CTQW; estimate $\hat{M}$; Braket/IBM	Qiskit, Braket SDK, Classiq
qsw_dephasing (HW/noise)	$(H_K, p) \rightarrow \eta(p)$	Lindblad QSW; sweep p ; $p \rightarrow 1$ classical, ENAQT optimum = resilience	QuTiP

4. Data

No training set. The method is training-free, so there is no small- n or generalization problem.

- **Calibration set (optional, only 2–4 hyperparameters):** dozens of proteins from ASD / ASBench (apo/holo + annotated allosteric sites), used under a leave-one-protein-out scheme to set the distal threshold, diagonal weight, and phase amplitude; **excludes the three target proteins**.
- **Validation set (primary):** KRAS G12C (4OBE $\rightarrow$ 6OIM), BCR-ABL1 (1OPL $\rightarrow$ 5MO4), cardiac myosin (5TBY $\rightarrow$ 6C1H), with apo input and holo-defined ground truth; c-Myc (1NKP) as an IDP stretch target (anchored on its functional interface instead).
- **Extended validation (bonus):** additional ASD apo–holo pairs (disjoint from the calibration set) to demonstrate generalizability.

5. Validation

- **Primary metric (unsupervised discrimination, matching the challenge primary objective):** per target, Mann–Whitney / permutation p -value, ROC-AUC, Top-5 hit rate, and enrichment of allosteric-residue scores vs. background (random residues + non-functional surface pockets).
- **Baselines:** classical communicability e^A (Estrada), pure GNM centrality, and random—to demonstrate the delta of the e^{-iHt} interference.
- **Noise resilience:** QSW dephasing sweep + ZNE on the hardware demo.
- **Scalability:** retention of the $\hat{M}$ topological signal before vs. after coarse-graining.

6. Implementation

Stage	Packages
Structure / features	BioPython, ProDy (GNM/ANM), fair-esm / HuggingFace (ESM-2), fpocket (surface pockets), NumPy, SciPy
Quantum simulation (prediction)	NumPy/SciPy eigendecomposition ($\hat{M}$), PennyLane (differentiable verification), QuTiP (QSW/Lindblad)
Quantum circuits / hardware	Qiskit + qiskit-ibm-runtime, AWS Braket SDK, Classiq (synthesis + depth optimization), Mitiq (ZNE)
Visualization	PyMOL (3D connectivity map), NetworkX, Matplotlib

7. Feasibility and Novelty

Feasibility. The classically exact $\hat{M}$ is trivial; the small- K hardware demo has precedent (Mohtashim 2026 ran a CTQW on protein RINs on IBM hardware); the entire toolchain is open source. **Novelty (relative to purely topological prior work):** (1) evolutionary conservation placed as **diagonal on-site energy**; (2) ANM direction $\rightarrow$ a **directional connectivity matrix via chiral phases**; (3) **active-site-anchored Krylov coarse-graining**; (4) the average mixing matrix as a training-free deliverable; (5) **apo $\rightarrow$ holo blind testing** on undruggable targets. Each rests on genuine literature; the combination is new.

8. Deliverables

- **Quantum connectivity matrix:** the $N \times N$ average mixing matrix $\hat{M}$.
- **Top-5 allosteric residue list:** ranked purely by the background-standardized specific connectivity $z(j)$ (residue level, no pocketness); pocketness serves only as a supplementary druggability annotation (see §3.5).
- **Methodology report:** the three-layer quantum positioning (coherent-interference model / hardware credible path / future multi-particle advantage) + evidence of the e^{-iHt} vs. e^A discriminative power.
- **Robustness & visualization:** QSW noise resilience, coarse-graining scalability, and 3D connectivity maps.

9. References

1. Mohtashim, S. I., Sajjan, M., & Kais, S. (2026). Continuous-Time Quantum-Walk Centrality for Protein Residue Interaction Networks. *J. Am. Chem. Soc.*, 148(27), 29206–29219. (prior art/baseline: CTQW on protein RINs, IBM hardware)
2. Godsil, C. (2013). Average mixing of continuous quantum walks. *J. Comb. Theory Ser. A*, 120(7), 1649–1662. (core readout: average mixing matrix = connectivity deliverable)
3. Novo, L., Chakraborty, S., Mohseni, M., Neven, H., & Omar, Y. (2015). Systematic Dimensionality Reduction for Quantum Walks. *Sci. Rep.*, 5, 13304. (Krylov coarse-graining: transport-preserving fixed-dimension reduction)
4. Wang, Y., Xue, S., Wu, J., & Xu, P. (2022). Continuous-time quantum walk based centrality testing on weighted graphs. *Sci. Rep.*, 12, 6001. (weighted-graph CTQW centrality)
5. Childs, A. M., & Goldstone, J. (2004). Spatial search by quantum walk. *Phys. Rev. A*, 70, 022314. (marked-vertex anchoring; γ_c from the spectrum)
6. Lu, D., Biamonte, J. D., Li, J., et al. (2016). Chiral quantum walks. *Phys. Rev. A*, 93, 042302. (directionality: complex edge phases break time reversal)
7. Whitfield, J. D., Rodríguez-Rosario, C. A., & Aspuru-Guzik, A. (2010). Quantum stochastic walks. *Phys. Rev. A*, 81, 022323. (QSW: coherent $\leftrightarrow$ classical Lindblad interpolation, noise)
8. Mohseni, M., Rebentrost, P., Lloyd, S., & Aspuru-Guzik, A. (2008). Environment-assisted quantum walks in photosynthetic energy transfer. *J. Chem. Phys.*, 129, 174106. (ENAQ: noise-assisted transport)
9. Caruso, F., Chin, A. W., Datta, A., Huelga, S. F., & Plenio, M. B. (2009). Highly efficient energy excitation transfer in light-harvesting complexes. *J. Chem. Phys.*, 131, 105106. (sink Lindblad + transfer-efficiency readout)
10. Campbell, E. (2019). A random compiler for fast Hamiltonian simulation. *Phys. Rev. Lett.*, 123, 070503. (qDRIFT: shallow-circuit simulation of dense H)
11. Chen, et al. (2024). Implementation of a Continuous-Time Quantum Walk on a Sparse Graph. *Phys. Rev. A*, 110, 052215. (sparse-graph CTQW circuit, star decomposition)

12. Oh, E. K., Krogmeier, T. J., Schlimgen, A. W., & Head-Marsden, K. (2024). Singular Value Decomposition Quantum Algorithm for Quantum Biology. *ACS Phys. Chem. Au*, 4(4), 393–399. (open-system biological dynamics on quantum hardware)
13. Chennubhotla, C., & Bahar, I. (2007). Signal Propagation in Proteins and Relation to Equilibrium Fluctuations. *PLoS Comput. Biol.*, 3(9), e172. (core biophysics: topology-driven allosteric propagation)
14. Estrada, E., & Hatano, N. (2008). Communicability in complex networks. *Phys. Rev. E*, 77, 036111. (classical baseline e^A , the analog of quantum e^{-iHt})
15. Bahar, I., Atilgan, A. R., & Erman, B. (1997). Direct evaluation of thermal fluctuations ... single-parameter harmonic potential. *Fold. Des.*, 2(3), 173–181. (GNM / Kirchhoff Γ)
16. Atilgan, A. R., Durell, S. R., Jernigan, R. L., et al. (2001). Anisotropy of Fluctuation Dynamics of Proteins with an Elastic Network Model. *Biophys. J.*, 80(1), 505–515. (ANM; source of chiral-phase direction)
17. Wang, J., Jain, A., McDonald, L. R., et al. (2020). Mapping allosteric communications within individual proteins. *Nat. Commun.*, 11, 3862. (Ohm: purely structural allosteric communication network)
18. Lin, Z., Akin, H., Rao, R., et al. (2023). Evolutionary-scale prediction of atomic-level protein structure with a language model. *Science*, 379(6637), 1123–1130. (ESM-2: conservation/evolutionary features)
19. Liu, X., Lu, S., Song, K., et al. (2020). Unraveling allosteric landscapes of allosterome with ASD. *Nucleic Acids Res.*, 48(D1), D394–D401. (ASD: calibration/extended-validation database)
20. Ostrem, J. M., Peters, U., Sos, M. L., Wells, J. A., & Shokat, K. M. (2013). K-Ras(G12C) inhibitors allosterically control GTP affinity and effector interactions. *Nature*, 503, 548–551. (KRAS switch-II ground truth)
21. Wylie, A. A., Schoepfer, J., Jahnke, W., et al. (2017). The allosteric inhibitor ABL001 enables dual targeting of BCR–ABL1. *Nature*, 543, 733–737. (BCR–ABL1 myristoyl ground truth)
22. Anderson, R. L., Trivedi, D. V., Sarkar, S. S., et al. (2018). Deciphering the super relaxed state of human β -cardiac myosin and the mode of action of mavacamten. *PNAS*, 115(35), E8143–E8152. (myosin mavacamten ground truth)