

# Chapter 2

---

# Magnetism and Magnetic Materials

韩伟

量子材料科学中心

2018年9月28日

# 助教



马扬

mmm@pku.edu.cn

Homework:  
Paper or electronic  
by email

# 上节课总结

本课程介绍自旋电子学基础知识以及其最新进展。

- 1) 介绍自旋电子学的基础知识，包括铁磁、反铁磁性、磁阻等。
- 2) 着重介绍自旋电子学的最近进展，包括自旋阀、自旋转移力矩、热自旋电子学、拓扑自旋、反铁磁自旋电子学等。


# 上节课总结

## 本科模式 VS 博士模式

被动接受知识  
考前突击 (A+)  
作业检查  
独立学习 (死磕)  
已知知识  
不懂不问  
定律  
全部学会  
老师全都会  
课堂上讲答案  
Correct/Wrong

主动学习知识  
平时积累, 没有考试  
(谁懂谁知道)  
研讨seminar  
未知知识  
提问讨论  
尝试  
效率  
用什么学什么  
合作性  
老师告诉哪  
里找答案  
Critical, Creative

# 上节课总结



# 提纲

**1. Introduction to magnetism**

**2. How to induce magnetic moment**

**3. How to control magnetization**

# 提纲

## 1. Introduction to magnetism

# Introduction to Magnetism

- **Magnetism of Electrons**

- **Spin orbit Coupling**

- **Magnetism**

**Diamagnetism, Paramagnetism,  
FM, AFM, Ferrimagnet, Half metallic**


- **Magnetic resonance**

- **Magnetic domains**




# Magnetism of Electrons

**Orbital moment**



(a)


**Spin moment**



(b)

# Orbital moment

## Bohr Atom



Speed of the electron:  $v$

Period of rotation:

$$\tau = 2\pi r / v$$

The equivalent current:

$$I = -e / \tau$$

The magnetic moment:

$$m = IA = -\frac{1}{2} e \vec{r} \times \vec{v}$$

Angular momentum:


$$\vec{l} = m_e \vec{r} \times \vec{v}$$

The moment:

$$m = -\frac{e}{2m_e} \vec{l}$$

# Orbital moment

## Bohr Atom



The moment:


$$m = -\frac{e}{2m_e} \vec{l}$$

**gyromagnetic ratio**

$$m_z = -\frac{e}{2m_e} m_l \hbar, m_l = 0, \pm 1, \pm 2, \dots$$

# Orbital moment

Bohr Atom




Bohr Magneton:

$$\mu_B = \frac{e\hbar}{2m_e}$$

$$1 \mu_B = 9.274 \times 10^{-24} \text{ A m}^2$$

# Spin moment

Bohr Atom



(b)

The spin moment:

$$\mathbf{m} = -\frac{e}{m_e} \vec{s}$$

$$m_z = -\frac{e}{m_e} m_s \hbar, m_s = \pm \frac{1}{2}$$

# Magnetism of Electrons

**Table 3.1.** Properties of the electron

Mass	$m_e$	$9.109 \times 10^{-31} \text{ kg}$
Charge	$-e$	$-1.6022 \times 10^{-19} \text{ C}$
Spin quantum number	$s$	$1/2$
Spin angular momentum	$\frac{1}{2}\hbar$	$5.273 \times 10^{-34} \text{ J s}$
Spin g-factor	$g$	$2.0023$
Spin magnetic moment	$\mu$	$-9.285 \times 10^{-24} \text{ A m}^2$
Classical radius $\mu_0 e^2 / 4\pi m_e$	$r_e$	$2.818 \times 10^{-15} \text{ m}$

# Gyromagnetic ratio

$$\gamma = -\frac{q}{2m}$$

**For a free electron:**


$$\gamma = -\frac{e}{2m_e} g_e = 1.760 \times 10^{11} \frac{rad}{s * T}$$

# Gyromagnetic ratio

$$\gamma = -\frac{q}{2m}$$




For a nucleus:

$$\gamma_n = -\frac{e}{2m_p} g_n = g_n \mu_n / \hbar$$






# Gyromagnetic ratio



Chudo, et al, Applied Physics Express (2014)

# Gyromagnetic ratio




Chudo, et al, Applied Physics Express (2014)

# Gyromagnetic ratio

Nucleus	$\gamma$ ( $10^6 \text{ rad s}^{-1} \text{ T}^{-1}$ )
$^1\text{H}$	267.513
$^2\text{H}$	41.065
$^3\text{He}$	-203.789
$^{13}\text{C}$	67.262
$^{14}\text{N}$	19.331
$^{15}\text{N}$	-27.116
$^{17}\text{O}$	-36.264
$^{29}\text{Si}$	-53.190
$^{31}\text{P}$	108.291



# Gyromagnetic ratio




$H_2O$

# Hund's Rule

- (1) First maximize  $S$  for the configuration.
- (2) Then maximize  $L$  consistent with  $S$ .
- (3) Finally couple  $L$  and  $S$  to form  $J$ :  $J = L - S$  if the shell is less than half full, and  $J = L + S$  if the shell is more than half full. When the shell is exactly half full,  $L = 0$  and  $J = S$ .



# Hund's Rule



(a)

# Hund's Rule

**Table 4.7.** The 3d ions.  $m_{eff}$  is in units of  $\mu_B$

$3d^n$		$S$	$L$	$J$	$g$	$m_{eff} = \frac{m_{eff}}{g\sqrt{J(J+1)}}$	$m_{eff} = \frac{m_{eff}}{g\sqrt{S(S+1)}}$	$m_{eff}^{exp}$
1	Ti <sup>3+</sup> , V <sup>4+</sup>	$\frac{1}{2}$	2	$\frac{3}{2}$	$\frac{4}{5}$	1.55	1.73	1.7
2	Ti <sup>2+</sup> , V <sup>3+</sup>	1	3	2	$\frac{2}{3}$	1.63	2.83	2.8
3	V <sup>2+</sup> , Cr <sup>3+</sup>	$\frac{3}{2}$	3	$\frac{3}{2}$	$\frac{2}{5}$	0.78	3.87	3.8
4	Cr <sup>2+</sup> , Mn <sup>3+</sup>	2	2	0			4.90	4.9
5	Mn <sup>2+</sup> , Fe <sup>3+</sup>	$\frac{5}{2}$	0	$\frac{5}{2}$	2	5.92	5.92	5.9
6	Fe <sup>2+</sup> , Co <sup>3+</sup>	2	2	4	$\frac{3}{2}$	6.71	4.90	5.4
7	Co <sup>2+</sup> , Ni <sup>3+</sup>	$\frac{3}{2}$	3	$\frac{9}{2}$	$\frac{4}{3}$	6.63	3.87	4.8

3d metals: magnetism mainly due to spin moment

# Spin orbit coupling

4f metals: orbit moment is large

$$H_{SO} = \lambda \vec{l} * \vec{s}$$

$$J = L + S$$



# Spin orbit coupling


**Table 4.6.** The 4 *f* ions. The paramagnetic moment  $m_{eff}$  and the saturation moment  $m_0$  are in units of  $\mu_B$

$4f^n$		$S$	$L$	$J$	$g$	$m_0 = gJ$	$m_{eff} = g\sqrt{J(J+1)}$	$m_{eff}^{exp}$
1	Ce <sup>3+</sup>	$\frac{1}{2}$	3	$\frac{5}{2}$	$\frac{6}{7}$	2.14	2.54	2.5
2	Pr <sup>3+</sup>	1	5	4	$\frac{4}{5}$	3.20	3.58	3.5
3	Nd <sup>3+</sup>	$\frac{3}{2}$	6	$\frac{9}{2}$	$\frac{8}{11}$	3.27	3.52	3.4
4	Pm <sup>3+</sup>	2	6	4	$\frac{3}{5}$	2.40	2.68	
5	Sm <sup>3+</sup>	$\frac{5}{2}$	5	$\frac{5}{2}$	$\frac{2}{7}$	0.71	0.85	1.7
6	Eu <sup>3+</sup>	3	3	0	0	0	0	3.4
7	Gd <sup>3+</sup>	$\frac{7}{2}$	0	$\frac{7}{2}$	2	7.0	7.94	8.9
8	Tb <sup>3+</sup>	3	3	6	$\frac{3}{2}$	9.0	9.72	9.8
9	Dy <sup>3+</sup>	$\frac{5}{2}$	5	$\frac{15}{2}$	$\frac{4}{3}$	10.0	10.65	10.6
10	Ho <sup>3+</sup>	2	6	8	$\frac{5}{4}$	10.0	10.61	10.4
11	Er <sup>3+</sup>	$\frac{3}{2}$	6	$\frac{15}{2}$	$\frac{6}{5}$	9.0	9.58	9.5
12	Tm <sup>3+</sup>	1	5	6	$\frac{7}{6}$	7.0	7.56	7.6
13	Yb <sup>3+</sup>	$\frac{1}{2}$	3	$\frac{7}{2}$	$\frac{8}{7}$	4.0	4.53	4.5

4f metals: magnetism described better by “J”

# D i a m a g n e t i s m



## Curie's Law: localized electron



*Diamagnetism*

# Paramagnetism

## Curie's Law: localized electron



$$M = n\mu_B \tanh(\mu_B B/k_B T)$$


# Curie-law paramagnetism

**Curie's Law: localized electron**

$$M = n\mu_B \tanh(\mu_B B / k_B T)$$


$$\chi = \mu_0 M / B = C / nT$$

$$C = n\mu_0\mu_B^2 / k_B$$



# FM and AFM


## Curie-Weiss Law for FM



$$\chi = C / (T - \theta_p)$$

# FM and AFM

## Susceptibility




# FM

Typical FM

	$T_C$ (K)	$d$ (kg m <sup>-3</sup> )	$\sigma_s$ (A m <sup>2</sup> kg <sup>-1</sup> )	$M_s$ (kA m <sup>-1</sup> )
Fe	1044	7874	217	1710
Co	1360( $\epsilon$ )	8920	162( $\epsilon$ )	1440
Ni	628	8902	54.8	488

# FM


## The Slater - Pauling curve





# Characterization of FM

## Vibrating sample magnetometer



Sensitivity:  $10^{-6}$  emu




# Characterization of FM

## SQUID Magnetometry




Sensitivity:  $5 \times 10^{-8}$  emu



# Characterization of FM

MOKE: Surface sensitive



**Table 5.13.** Kerr rotation in ferromagnetic metals at 830 nm (1.5 eV)

$\theta_K$ (°)		$\theta_K$ (°)	
Fe	−0.53	CoPd	−0.17
FePt	−0.39	CoPt	−0.36
FeCo	−0.60	Ni	−0.09
Co	−0.36	PtMnSb	−1.3

# AFM

## Typical AFM


**Table 6.1.** Some common antiferromagnets

	Structure	$T_N$ (K)	$\theta_p$ (K)	$\mu_0 M_\alpha$ (T)
Cr	sdw	311		0.20
Mn	Complex	96	$\sim -2000$	0.20
NiO	Néel	524	$-1310$	0.54
$\alpha\text{Fe}_2\text{O}_3$	Canted	958	$-2000$	0.92
$\text{MnF}_2$	Néel	67	$-80$	0.78
FeMn	Néel	510		0.53
$\text{IrMn}_3$	Néel	690		0.50

sdw – spin density wave; Néel – two collinear sublattices.


# AFM

M vs. H loop



# AFM



SAF: Synthetic antiferromagnets



Antiferromagnetic exchange coupling between two FM layers

# AFM


SAF: Synthetic antiferromagnets






Baibich, et al, PRL (1988)  
Fert, Rev. Mod. Phys. (2007)

# Ferrimagnet

AFM



Ferrimagnet




Temperature,  $T$





# Ferrimagnet

FM-Gd alloys



# Ferrimagnet





white:  $\text{O}^{2-}$ , form a fcc sublattice  
(He,  $2s^2$ ,  $2p^4$ )

blue: A site, tetrahedral,  $\text{Fe}^{3+}$  only  
(Ar,  $4s^2$ ,  $3d \downarrow^5$ )

red: B site, octahedral, half  $\text{Fe}^{3+}$ ,  
half  $\text{Fe}^{2+}$   
(Ar,  $4s^2$ ,  $3d \uparrow^5$ ,  $3d \downarrow^{0.5}$ )

# Ferrimagnet



Blue:  $\text{Fe}^{3+}$

Red:  $\text{Fe}^{3+}$  and  $\text{Fe}^{2+}$


$\text{Fe}^{3+} : 5\mu_{\text{B}}$

$\text{Fe}^{2+} : 4\mu_{\text{B}}$

Total moment:  $4\mu_{\text{B}} / \text{Fe}_3\text{O}_4$


# Ferrimagnet

YIG:  $\text{Gd}_3\text{Fe}_5\text{O}_{12}$



# Half Metallic

Density of states




Coey & Venkatesan, JAP (2002)

# Half Metallic

$\text{CrO}_2$

Spin polarization: 100%



Coey & Venkatesan, JAP (2002)

# Introduction to Magnetism

- **Magnetism of Electrons**

- **Spin orbit Coupling**

- **Magnetism**

**Diamagnetism, Paramagnetism,  
FM, AFM, Ferrimagnet, Half metallic**

- **Magnetic resonance**

- **Magnetic domains**

**休息10分钟**





# 提纲

- 1. Introduction to magnetism**
- 2. How to induce magnetic moment**
- 3. How to control magnetization**

# Review of last class

➤ **Magnetism of Electrons**

➤ **Spin orbit Coupling**

➤ **Magnetism**

**Diamagnetism, Paramagnetism,  
FM, AFM, Ferrimagnet, Half metallic**

# Introduction to Magnetism

➤ **Magnetism of Electrons**

➤ **Spin orbit Coupling**

➤ **Magnetism**

**Diamagnetism, Paramagnetism,  
FM, AFM, Ferrimagnet, Half metallic**


➤ **Magnetic resonance**

➤ **Magnetic domains**

# Magnetic resonance

## Landau-Lifshitz-Gilbert equation

$H_{eff}$  (static)



$$\frac{d\hat{m}}{dt} = -\gamma \hat{m} \times \vec{H}_{eff} + \alpha \hat{m} \times \frac{d\hat{m}}{dt}$$


$$\gamma = \frac{g e}{2 m_e c} \text{ is gyromagnetic ratio}$$

$\alpha$  is the Gilbert damping

$H_x e^{i\omega t}$  (rf): small perturbation

# Magnetic resonance



## FMR



# Magnetic resonance

## FMR system


**PPMS**  
**T: 2 K - 300 K**  
**B: 9T**



**Coplanar Waveguide**

# Magnetic resonance

## Metallic FM– Py (NiFe)




Zhao, et al, Scientific Reports (2016)



# Magnetic resonance



## Semiconducting FM—Cr<sub>2</sub>Ge<sub>2</sub>Te<sub>6</sub>



Zhang, et al, JJAP (2016)


# Magnetic resonance

## Insulating FM-- YIG



# Antiferromagnetic resonance



## Precessing Modes



$$\omega_0 = \gamma \mu_0 [H_a(H_a + 2H_{ex})]^{\frac{1}{2}}.$$


# Magnetic domains

## FM domains



# Magnetic domains

## Racetrack memory




## Outline

# **2. How to induce magnetic moment**


# Mainly two methods

## 1) Impurity doping

### Mn impurity in GaMnAs



Low doping





High doping

MacDonald, et al, Nature Mater. (2005)

# Mainly two methods

## 1) Impurity doping




MacDonald, et al, Nature Mater. (2005)



# Mainly two methods

## 1) Impurity doping

### p-d Zener mode




Dietl, et al, Science (2010)

# Mainly two methods


## 2) Proximity effect

At the atomic level, when two atoms come into proximity, the highest energy, or valence, orbitals of the atoms change substantially and the electrons on the two atoms reorganize.



# Mainly two methods


## 2) Proximity effect



Klewe, et al, PRB (2016)


# Mainly two methods

## 2) Proximity effect



# Mainly two methods

## 2) Proximity effect





Composition	Pt thickness (nm)	FM thickness (nm)	Magnetic moment ( $\mu_B$ per atom)
Pt/Fe	1.8	9.8	$0.2 \pm 0.1^a$
	3.4	9.2	$0.6 \pm 0.1^a$
	5.9	9.8	$0.6 \pm 0.1^a$
	20.0	9.8	$0.6 \pm 0.1^a$
	3.4	9.2	$0.43 \pm 0.08^b$
Pt/Ni <sub>33</sub> Fe <sub>67</sub>	3.3	8.3	$0.44 \pm 0.10^b$
Pt/Ni <sub>81</sub> Fe <sub>19</sub>	3.3	9.8	$0.22 \pm 0.10^b$
Pt/Ni	3.2	9.8	$0.08 \pm 0.08^b$

# Induce M in two Quantum Materials

## Two Dirac Materials


### Graphene




# Induce M in two Quantum Materials

Two Dirac Materials

Topological Insulator



3D Topological insulator







# Induce M in graphene

## Why making graphene magnetic

PHYSICAL REVIEW B 83, 155447 (2011)

### Quantum anomalous Hall effect in single-layer and bilayer graphene

Wang-Kong Tse,<sup>1</sup> Zhenhua Qiao,<sup>1</sup> Yugui Yao,<sup>1,2</sup> A. H. MacDonald,<sup>1</sup> and Qian Niu<sup>1,3,\*</sup>



PHYSICAL REVIEW B 77, 115406 (2008)

### Spin transport in proximity-induced ferromagnetic graphene


Håvard Haugen,<sup>\*</sup> Daniel Huertas-Hernando, and Arne Brataas

*Department of Physics, Norwegian University of Science and Technology, N-7491 Trondheim, Norway*



# Induce M in graphene






How to make graphene magnetic



Yazyev and Helm, PRB (2007)  
Han, et al, Nature Nanotech (2014)

# Vacancies Defects

Vacancies Defects → FM




Cervenka, et al, Nature Phy. (2009)

# Vacancies Defects

Vacancies Defects → PM


$$M = NgJ\mu_B \left[ \frac{2J}{2J+1} \operatorname{ctnh} \left( \frac{(2J+1)z}{2J} \right) - \frac{1}{2J} \operatorname{ctnh} \left( \frac{z}{2J} \right) \right]$$



Nair, et al, Nature Phy. (2012)

# Vacancies Defects


Vacancies Defects  $\rightarrow$  PM



Nair, et al, Nature Phy. (2012)

# Vacancies Defects

**Question?**



**Ferromagnetic ??**

**Paramagnetic ??**

# 下一节课: Oct. 12th

---

**1. Introduction to magnetism**

**2. How to induce magnetic moment**

**3. How to control magnetization**

课件下载：

<http://www.phy.pku.edu.cn/~LabSpin/teaching.html>

**谢谢！**