

Edge conduction in monolayer WTe_2


Zaiyao Fei¹, Tauno Palomaki¹, Sanfeng Wu¹, Wenjin Zhao¹, Xinghan Cai¹, Bosong Sun¹, Paul Nguyen¹, Joseph Finney¹, Xiaodong Xu^{1,2*} and David H. Cobden^{1*}

Outline


- Background
- Experimental results & explanations
- Summary

◦ Background

Quantum spin Hall effect and 2DTI



Schematic of the spin-polarized edge channels in a quantum spin Hall insulator in a quantum well.



Band structure of 2DTI

◦ Background

Some puzzles in experimental works on 2DTI of quantum wells


- The conductance at low temperatures is not perfectly quantized, becoming small in long edges
- The edge conductance showing mesoscopic fluctuations as a function of gate voltage.
- The edges show signs of conducting even at high magnetic field.
- Non-helical edge conduction may also be present, due for instance to band bending when a gate voltage is applied in the quantum wells.



So, identification of a natural monolayer 2DTI would be helpful for elucidating and exploiting TI physics


◦ Experimental results & explanations

Temperature dependence of G_{diff} in different thickness WTe_2




- Experimental results & explanations

Distinguishing edge and bulk conduction




Traditional nonlocal measurement to detect edge conductance




Quantitative separation of bulk and edge contributions

- Experimental results & explanations


Distinguishing edge and bulk conduction




Temperature dependence of the conductance G_{edge} at zero magnetic field



Gate-averaged linear-response edge conductance for 19 adjacent-contact pairs



Effect of in-plane magnetic field B_{\parallel} at 1.6 K



Sweeps of B_{\parallel} at $V_g = 0$ for various temperatures

◦ Experimental results & explanations

Some explanations

- The monolayer edge conductance is independent of V_g . This is consistent with a single gapless mode, and not with carrier accumulation due to band bending.
- No edge conduction in bilayers. This can be explained by the fact that TR symmetry does not prohibit backscattering at the bilayer edge.
- The conductance is dramatically suppressed by $B_{//}$, consistent with the expectation that elastic backscattering is allowed once TR symmetry is broken.

◦ Summary

- At temperatures below about 100 K, monolayer WTe₂ does become insulating in its interior, while the edges still conduct.
- The edge conduction is strongly suppressed by an in-plane magnetic field and is independent of gate voltage.
- significant elastic scattering in the edge.