

Tracing the Unlabeled Storm: Cross-Variable Transfer in a Lagrangian Atmospheric JEPA Framework

K M Anirudh^{1*}, S Sandeep^{1,3}, Hariprasad Kodamana^{2,3}

¹Centre for Atmospheric Sciences, IIT Delhi, New Delhi, India

²Department of Chemical Engineering, IIT Delhi, New Delhi, India

³Yardi School of Artificial Intelligence, IIT Delhi, New Delhi, India
anirudh@cas.iitd.ac.in

Abstract

Deep atmospheric convection governs South Asian monsoon variability, yet attempting to learn its latent world model directly from zero-inflated, heavy-tailed precipitation yields suboptimal predictive representations. Continuous atmospheric proxies, such as outgoing longwave radiation (OLR), express this convective organization far more coherently. We address this mismatch with *cross-variable proxy learning*: M-JEPA, a multiscale Monsoon Joint-Embedding Predictive Architecture, is pretrained on five continuous proxy fields over Lagrangian patches tracking moving convective systems—without rainfall supervision at any point. The resulting frozen representation is transferred to daily precipitation forecasts through a shared decoder trunk featuring parallel probabilistic and deterministic branches. Because rainfall is strictly unobserved during pretraining, downstream skill directly measures the predictive information captured in the latent rollout. A frozen-backbone probing framework with two controls (an identical architecture trained on rainfall alone, and a randomly initialized backbone) attributes the transfer specifically to proxy pretraining: direct rainfall training exhibits 36% higher CRPS error (7.52 vs. 5.54 mm/day). Against the 51-member operational ECMWF ensemble, the transferred model attains a statistically resolved CRPS advantage (6.81 vs. 6.89 mm/day) and higher Brier skill (+0.05 vs. −0.04) using 15.4M parameters on a single consumer GPU, concentrated at heavy-rain thresholds and fine spatial scales, while the ensemble retains an advantage in neighborhood skill and deterministic references on point metrics. The result provides a competitive monsoon precipitation forecast grounded in intraseasonal dynamics and a diagnostic framework for evaluating transferred atmospheric representations.

1 Introduction

The South Asian monsoon (SAM) supplies the water on which nearly a quarter of the world’s population depends, and its rainfall is among the most consequential and least predictable quantities in atmospheric science. Both physics-based numerical weather prediction (NWP) and modern data-driven forecasting systems continue to struggle with the processes governing its variability.

A fundamental challenge lies in precipitation itself. Unlike large-scale atmospheric variables, rainfall is sparse, intermittent, highly localized and strongly heavy-tailed, reflecting the nonlinear nature of deep convection. In NWP these processes occur below the grid scale and must be parameterized, a long-recognized source of systematic bias (Stephens et al. 2010). Machine-learned weather models (Lam et al. 2023; Bi et al. 2023; Lang et al. 2024; Bodnar et al. 2025) achieve state-of-the-art skill on smooth continuous fields, but treat precipitation as one target among many, optimized with the same generic objectives despite its very different statistics.

The dominant source of subseasonal variability in the SAM is the Monsoon Intraseasonal Oscillation (MISO), a northward-propagating envelope of organized deep convection that governs the transition between active and break phases of the monsoon (Goswami et al. 2003; Suhas, Neena, and Goswami 2013). Although rainfall constitutes the most visible manifestation of this evolution, the underlying convective organization is expressed far more coherently in continuous atmospheric fields, particularly outgoing longwave radiation (OLR). Rainfall superimposes fine-scale intermittency on this slowly evolving state, obscuring the organization that governs its occurrence. This motivates a cross-variable strategy in which the dynamics are learned from continuous proxies first and the representation is then transferred to rainfall.

Because the representation never sees rainfall, its ability to transfer to precipitation is a stringent test of the dynamics encoded in the latent. If a model pretrained only on continuous proxies predicts sparse, intermittent rainfall better than one trained directly on rainfall, the representation has captured transferable physical structure rather than task-specific shortcuts. Establishing that requires probing the representation itself, not downstream skill alone. Joint-Embedding Predictive Architectures (JEPAs) are widely motivated as learning latent “world models”. What their rollouts encode has begun to be examined on video and robotics data, but not for atmospheric state, and not where the readout target is a physical variable absent from pretraining.

We address this question with **M-JEPA**, a multiscale autoregressive JEPA trained in two stages. In the upstream stage, a context encoder and an exponential-moving-average target encoder embed Lagrangian patches that follow tracked convective envelopes, and an autoregressive predictor rolls

*Corresponding author.

the context latent forward seven daily steps under latent regression; the five proxies are the only inputs, and rainfall is absent from the objective. In the downstream stage, the encoder and predictor are frozen and the latent rollout is read out to daily precipitation by a shared decoder trunk with a probabilistic and a deterministic branch, matched to rainfall’s zero-inflated statistics. Rainfall is therefore a transfer target throughout rather than a pretraining objective, and every reported skill number is a measurement of the frozen representation. Our contributions are fourfold:

- **A probing framework for atmospheric world models.** A frozen-backbone protocol examining predictive information, variable-specific subspace organization, and temporal evolution through Centered Kernel Alignment (CKA) (Section 5); a fourth probe, latent-to-pixel attribution, is in supplementary Section D.4.
- **Evidence for transferable atmospheric dynamics.** Two controls, the identical architecture pretrained on rainfall alone and a randomly initialized backbone, attribute the transfer to learning on continuous proxies, not to architectural inductive bias or the decoder (Section 5.6).
- **Skill concentrated at spatially sharp extremes.** On matched windows M-JEPA Ens attains a lower fair CRPS than the 51-member operational IFS Ens, by a resolved but small margin (-0.08 mm/day, 95% CI $[-0.16, -0.001]$), with higher Brier Skill Score, from a 15.4M-parameter model on one consumer GPU. The margin is not generic: member-FSS at the IMD Heavy and Very Heavy thresholds grows with lead, and the fine-scale power spectrum tracks observations more closely than IFS Ens’s own members (Section 5.1).
- **Characterization of representation limitations.** The rollout is dynamically damped, saturating by lead 5, and the decoded forecast under-continues the observed decay for the longest-lived systems, motivating a scaling hypothesis in which faithful propagating dynamics need proxy information denser in time, vertical levels and spatial extent (Section 6).

2 Related Work

JEPA and latent world models. Joint-embedding predictive architectures predict in representation space rather than pixel space (Assran et al. 2023, 2025). M-JEPA follows V-JEPA 2 (Assran et al. 2025) most directly, transposing its context encoder, exponential-moving-average target and latent autoregressive rollout from video to a co-moving atmospheric patch. Work asking what such rollouts encode (Zhang et al. 2026; Nam et al. 2026; Thil et al. 2026) sits almost entirely in video and robotics, where the prediction target shares the encoder’s modality. We ask what a latent encodes when it is read out to a *different physical variable* than any it was pretrained on.

Machine-learned weather models. Emulators trained on reanalysis now match or exceed operational skill on the large-scale fields (Lam et al. 2023; Bi et al. 2023; Price et al. 2025; Lang et al. 2024), pretrained models extend the recipe across variables and resolutions (Bodnar et al. 2025; Nguyen

et al. 2023), and benchmarks have followed at medium and subseasonal-to-seasonal range (Rasp et al. 2024; Nathaniel et al. 2024). All are evaluated by forecast score alone, leaving unexamined what their representations encode and whether it transfers off the training variables. That is the question this paper puts to a JEPA latent, with GraphCast, AIFS and the operational IFS as references.

Data-driven monsoon and precipitation prediction.

Subseasonal prediction of the South Asian monsoon has largely proceeded by reducing the system before predicting it: the oscillation is compressed to an index or a few oscillatory modes, these are forecast, and rainfall is recovered from them (Anirudh et al. 2025; Bach et al. 2024). That keeps the northward march of the envelope but compresses away the zonal structure determining where rain falls; the representation here carries the evolving envelope field in longitude, latitude and time. Related efforts span Earth-system emulation of monsoon variability (Banerjee et al. 2026), subseasonal machine learning (He et al. 2021), radar nowcasting (Shi et al. 2015; Wen et al. 2026; Lin et al. 2022), dynamics-informed diffusion (Rühling Cachay et al. 2023), quantile-aware subseasonal precipitation (Chen et al. 2026) and explicit heavy-tail distributions (Naveau et al. 2016; Wilson, Tan, and Luo 2022). All learn rainfall from rainfall; the representation studied here never observes it.

Atmospheric latent representations. Atmospheric latents have been inspected directly (Wisinski et al. 2026; Fan et al. 2025; Zhao et al. 2026), but always as *reconstruction* latents of the same variables they encode. None studies a self-supervised *world-model* latent transferred to a held-out target. Frozen-encoder readouts with probabilistic heads exist for precipitation (Dourado Filho et al. 2025); that recipe is not claimed as novel, and forecasting skill is treated here as a *probe* of representation quality. The conjunction is new: a JEPA latent learned on object-centric *Lagrangian patches*, transferred across variables to a held-out *sparse* target, and analyzed with a proxy→latent→pixel probing suite and controls attributing the transfer to the proxies.

3 Data and Problem Setup

Approach. We forecast the next seven daily precipitation fields over the South Asian monsoon domain (9.5°S – 29.5°N , 60.5 – 99.5°E) on a 1° grid from five daily proxy fields, with rainfall absent from pretraining. In place of a fixed Eulerian window the model learns on *Lagrangian patches* following one convective envelope at a time, so the encoder sees a single system’s evolution in its own co-moving frame.

Event tracking. Envelopes are located by a classical detect-and-track pipeline on OLR band-passed to the 20–90 day intraseasonal band, which isolates the organized convective envelope of the MISO from synoptic and seasonal variance (Wheeler and Kiladis 1999; Suhas, Neena, and Goswami 2013). Being retrospective and zero-phase, this filter selects and centers patches only; it is never a model input, and the encoder consumes raw, unfiltered fields. Tracks are retained at a 12-day minimum and tracking is direction-blind. Poleward motion is therefore a property of the retained




Figure 1: The two stages of M-JEPA. *Stage I* (top): context encoder f_θ , target encoder f_ξ , autoregressive predictor g_ϕ , pretrained on proxies with no rain objective. *Stage II* (bottom): the frozen rollout $\hat{z}_{t+1:t+7}$, rain history $y_{t-4:t}$ and static geography s condition one shared decoder trunk whose two branches emit a 16-member probabilistic forecast and a deterministic field.

population (60% of test tracks), not a selection criterion. Because the frame re-centers on the tracked centroid at every step, the envelope’s bulk translation is supplied by the tracker and is never predicted; the model forecasts intensity and structure within the moving patch, and no metric credits it for knowing the system’s future position. A sweep over the two tracker settings that fix patch geometry leaves the ordering against persistence and deterministic IFS unchanged in all ten configurations, though not the ensemble comparison, which we could not re-run per configuration (supplementary Section E.5).

Data and windows. The proxies are OLR (Lee 2025) and specific humidity q , zonal and meridional wind u, v , and geopotential z at 850 hPa from ERA5 (Hersbach et al. 2020); the verification target is GPM satellite rainfall (Huffman, Bolvin, and Nelkin 2015). All fields are regridded to the target grid, deseasonalized, and z -scored on training-year statistics only. Each retained centroid anchors a tight foreground and a wider background crop, and every five-day context plus seven-day horizon span along a track yields one window. Splits are by year block: pretraining 1979–2017, decoder training 1998–2013 (667 windows, trained with ± 2 -grid-cell spatial-jitter augmentation), validation 2014–2017, test 2018–2025 (337 windows). Detection runs on all calendar months, so the pretraining population is intraseasonal convective envelopes generally; evaluation is restricted to JJAS, when the northward-propagating MISO envelope dominates that population. We impose none of the strict propagation filters used to define MISO indices, so the retained tracks are a broader, on average slower set than a canonical MISO composite (supplementary Section A.1). Pretraining years overlap the decoder years, but both routes to leakage are closed: rainfall is absent from every Stage-1 input and objective, and pretraining ends in 2017, before the first test year. Full details: supplementary Section A.

4 Method

M-JEPA is trained in two stages: self-supervised multiscale autoregressive JEPA pretraining on the proxies (Stage 1), then a rainfall readout on the frozen latent rollout (Stage 2). Figure 1 fixes the notation used throughout.

4.1 Stage 1: upstream pretraining on proxies

Each input clip is a five-day sequence of Lagrangian patches $x_{t-4:t}$ anchored on the tracked convective centroid, carried in two streams that differ in extent and token footprint, not in grid spacing. The *foreground* is OLR alone on a 20×20 patch at 1° , tokenized in 4×4 -cell tubelets; the *background* carries q, u, v, z at 850 hPa plus a spatially smoothed OLR envelope on a 40×40 patch about the same center, tokenized in 8×8 tubelets. Both therefore yield a 5×5 grid of 256-dimensional tokens, resolving convective detail in the core and the synoptic-to-planetary environment around it at equal token cost. Each stream is tokenized and tagged separately, $u_d^{\text{fg}} = \mathcal{E}^{\text{fg}}(x_d^{\text{fg}}) + e_{\text{fg}}$ and u_d^{bg} likewise, where the $\mathcal{E}$ are the per-stream tokenizers and $e_{\text{fg}}, e_{\text{bg}}$ are learned stream tags. The context encoder T_θ , a six-block eight-head pre-norm transformer, then embeds each day independently,

$$z_d = f_\theta(x_d) = T_\theta([u_d^{\text{fg}} \| u_d^{\text{bg}}]) \in \mathbb{R}^{50 \times 256}, \quad (1)$$

so the two scales interact through self-attention over the joint 50-token set. All temporal structure is left to the predictor. The context latents are the resulting five-day sequence $z_{\text{ctx}} = (z_{t-4}, \dots, z_t)$.

The autoregressive predictor g_ϕ is a block-causal transformer (six blocks, eight heads) that emits a residual increment and is applied to its own outputs to roll the state forward,

$$\begin{aligned} \hat{z}_{t+d} &= \hat{z}_{t+d-1} + g_\phi(z_{\text{ctx}}, \hat{z}_{t+1:t+d-1}), \\ \hat{z}_t &\equiv z_t, \quad d = 1, \dots, 7, \end{aligned} \quad (2)$$

so each step conditions on the full sequence produced so far, and the zero-initialized output projection starts training at persistence. Regression targets are supplied by the target encoder f_ξ , an exponential-moving-average copy of f_θ that receives no gradient,

$$\begin{aligned} z_{\text{tgt}, t+d} &= f_\xi(x_{t+d}), \\ \xi &\leftarrow \mu \xi + (1 - \mu) \theta, \quad \mu = 0.996. \end{aligned} \quad (3)$$

The Stage-1 objective combines latent regression in two forms with variance-covariance regularization and an auxiliary reconstruction,

$$\begin{aligned} \mathcal{L}_1 &= \sum_j \ell(z_j + g_\phi(z_{\leq j}), z_{\text{tgt}, j+1}) \\ &\quad + \sum_{d \leq k} \ell(\hat{z}_{t+d}, z_{\text{tgt}, t+d}) \\ &\quad + \mathcal{V}(z) + 0.04 \mathcal{C}(z) \\ &\quad + 0.1 \ell(D_{\text{aux}}(z_d), x_d^{\text{fg}}), \end{aligned} \quad (4)$$

where ℓ is the smooth- L_1 loss. The first term is teacher-forced and regresses every observed one-day transition of the clip; the second is the free-running rollout at horizon k , raised on a curriculum from two to seven days. $\mathcal{V}$ and $\mathcal{C}$ are the hinged unit-variance and squared off-diagonal covariance terms of VICReg (Bardes, Ponce, and LeCun 2022) that prevent latent collapse, and D_{aux} is a reconstruction head that keeps the latent spatially decodable. Rainfall enters neither (4) nor any Stage-1 input. Trainable parameters are 4.8M in f_θ and 4.9M in g_ϕ ; f_ξ and D_{aux} are discarded after pretraining.

Both f_θ and g_ϕ are frozen for Stage 2, which makes the rollout a fixed operator rather than a tunable feature extractor. Every probe and every forecast score therefore measures the pretrained representation, not its adaptation to rainfall.

4.2 Stage 2: downstream rainfall readout

Stage 2 reads three inputs per window: the frozen rollout $\hat{z}_{t+1:t+7}$ from (2), the observed five-day rain history $y_{t-4:t}$ on the 20×20 target patch, and static geography s (land fraction, latitude, longitude). It has two parts, taken in turn below: a shared trunk that decodes the rollout to output resolution, and two branches that read the trunk. Writing $\hat{z}^{\text{fg}}$ and $\hat{z}^{\text{bg}}$ for the foreground and background token blocks, the shared decoder trunk projects both back onto the 5×5 token grid, fuses the two scales there with a 3×3 convolution, and upsamples to the output grid through two transposed-convolution blocks ($5 \rightarrow 10 \rightarrow 20$, 128 channels),

$$m_{t+d} = \mathcal{U}(W_{\text{fuse}}[p_{t+d}^{\text{fg}} \| p_{t+d}^{\text{bg}}]) \in \mathbb{R}^{128 \times 20 \times 20}, \quad (5)$$

where $p_{t+d}^{\text{fg}} = P_{\text{fg}}(\hat{z}_{t+d}^{\text{fg}})$ is the projected foreground token block and p_{t+d}^{bg} likewise. Fusing at token resolution combines the two scales while they are still spatially registered. Rain history and geography enter afterwards at output resolution through their own convolutional encoders E_y , applied to $\tilde{y} = \log(1 + y_{t-4:t})$ with a learned token for windows that have no history, and E_s , combined by 1×1 convolutions,

$$\begin{aligned} c &= h(\hat{z}_{t+d}, y_{t-4:t}, s) \\ &= W_s[W_y[m_{t+d} \| E_y(\tilde{y})] \| E_s(s)]. \end{aligned} \quad (6)$$

The E_y and E_s maps are computed once per window and broadcast over leads, so all lead-to-lead variation in c originates in the rollout. A 1×1 readout $\hat{o}_{t+d} = W_{\text{olr}}(m_{t+d})$ of the trunk feature also predicts the true future OLR field o_{t+d} , regularizing the trunk with

$$\mathcal{L}_{\text{OLR}} = \ell(\hat{o}_{t+d}, o_{t+d}), \quad (7)$$

the same smooth- L_1 ℓ as (4).

Two branches read c . The *probability* branch is a rectified flow with no deterministic anchor, unlike residual-correction designs such as CorrDiff (Mardani et al. 2025), and transports Gaussian noise to rain fields in square-root space: with $x_1 = \sqrt{y}$, $x_0 \sim \mathcal{N}(0, I)$, $\tau \sim \mathcal{U}(0, 1)$ and $x_\tau = (1 - \tau)x_0 + \tau x_1$,

$$\begin{aligned} \mathcal{L}_{\text{fm}} &= \mathbb{E} \|v_\theta(x_\tau, \tau, c) - (x_1 - x_0)\|^2, \\ \mathcal{L}_{\text{es}} &= \text{ES}(\{x^{(k)}\}_{k=1}^4, x_1), \end{aligned} \quad (8)$$

the auxiliary term is the energy score (Gneiting and Raftery 2007) of four differentiable draws, scoring whole sampled fields where $\mathcal{L}_{\text{fm}}$ constrains only the velocity at one interpolation time. The velocity field v_θ is a $20 \rightarrow 10 \rightarrow 5 \rightarrow 10 \rightarrow 20$ U-Net of width 96 (3.0M parameters) taking x_τ and a sinusoidal embedding of τ , with c pooled and re-concatenated at every scale so the conditioning reaches the bottleneck undiluted. The trunk and probability branch train jointly,

$$\mathcal{L}_2 = \mathcal{L}_{\text{OLR}} + \mathcal{L}_{\text{fm}} + 0.1 \mathcal{L}_{\text{es}}, \quad (9)$$

so the trunk’s only training signal is this distributional objective.

The *deterministic* branch $\hat{y} = h_\theta(c)$ shares the flow branch’s U-Net topology at width 32 (0.3M parameters) but takes neither noise nor τ , giving a single forward pass with a softplus output. It is trained afterward, as a separate readout on the same trunk now held frozen, on

$$\mathcal{L}_{\text{det}} = \mathcal{L}_{\text{FACL}}(\hat{y}, y) + \lambda \mathcal{L}_{\text{TDT}}(\hat{y}, y), \quad \lambda = 0.02, \quad (10)$$

where, writing $\mathcal{F}$ for the 2-D orthonormal DFT over the spatial dims of a lead’s field,

$$\mathcal{L}_{\text{FAL}} = \mathbb{E}[(|\mathcal{F}\hat{y}| - |\mathcal{F}y|)^2], \quad \mathcal{L}_{\text{FCL}} = 1 - \frac{\text{Re}\langle \mathcal{F}\hat{y}, \mathcal{F}y \rangle}{\|\mathcal{F}\hat{y}\| \|\mathcal{F}y\|}, \quad (11)$$

and $\mathcal{L}_{\text{FACL}}$ selects $\mathcal{L}_{\text{FAL}}$ with probability $p(\text{step})$, annealed $1 \rightarrow 0$ over the first 70% of training, and $\mathcal{L}_{\text{FCL}}$ otherwise: spectral sharpness early, placement late. Writing $d\hat{y}_{t+d} = \hat{y}_{t+d+1} - \hat{y}_{t+d}$ for the predicted lead-to-lead tendency and dy_{t+d} likewise for observed,

$$\begin{aligned} \mathcal{L}_{\text{TDT}} &= \mathbb{E}[w_{t+d} |d\hat{y}_{t+d} - dy_{t+d}|], \\ w_{t+d} &= 2 \text{ if } dy_{t+d} < 0, \quad d\hat{y}_{t+d} > dy_{t+d}, \text{ else } 1, \end{aligned} \quad (12)$$

i.e. an L_1 tendency penalty up-weighted by $2 \times$ specifically when the forecast under-decays relative to a truly decaying observation. Training the branches sequentially sidesteps the double-penalty problem structurally rather than by loss weighting (Subich et al. 2025), since a likelihood forecast and a squared-error point forecast are different functionals of the predictive distribution, so one trunk optimized for both at once is optimal for neither. Trunks trained under $\mathcal{L}_{\text{det}}$ itself

Table 1: Forecast skill on matched windows against the 51-member operational IFS ensemble ($n=169$, fair CRPS in mm/day; the IFS (det.) row is on its own single-member convention). The CRPS difference against IFS Ens is -0.08 mm/day, 95% paired-bootstrap CI $[-0.16, -0.001]$ over 10^4 resamples. Boldface marks the better ensemble. Methodology: supplementary Section C.

	Members	CRPS ↓	BSS ↑	FSS10 ↑	Spread/RMSE → 1	Rank dev. ↓
M-JEPA Ens	16	6.81	+0.05	0.53	0.73	0.60
IFS Ens	51	6.89	-0.04	0.63	0.45	0.95
IFS (det.)	1	9.76	-0.36	0.73	0.00	0.70

top out at pooled ACC 0.255–0.271 regardless of curriculum, whereas reading out from the frozen distributionally-trained trunk reaches 0.292 on that same pooled convention; the corresponding case-averaged scores, which are the convention used for every result we report, are in Section 5.2. Across Stage 2 the trunk holds 2.1M parameters and the conditioning encoders 0.4M, for 5.7M against the 9.7M frozen Stage-1 stack (components rounded independently, summing to 5.8M).

Each window therefore yields, at every lead $d = 1, \dots, 7$, a 16-member probabilistic rainfall field $\{\hat{y}_{t+1:t+7}^{(k)}\}_{k=1}^{16}$, obtained by integrating v_θ from independent noise and squaring back to mm day $^{-1}$, and one deterministic field $\hat{y}_{t+1:t+7}$, both used directly with no post-hoc recalibration. Architecture and optimization settings for both stages: supplementary Section B.

Every score below is computed in the patch frame and case-averaged, using the fair unbiased CRPS estimator (Ferro, Richardson, and Weigel 2008) and the NHC homogeneous-sample matching rule; these conventions, and member-FSS, Brier skill and ACC, are defined once in supplementary Section C and then used without restatement.

5 Results

We report forecast skill against the operational references first, then what the frozen representation contains that makes that skill possible. Token- and pixel-level attribution is deferred to supplementary Section D.4.

5.1 Probabilistic forecast skill

The probabilistic reference is IFS Ens, the 51-member operational ECMWF ensemble, and the same underlying model as the deterministic IFS run of Section 5.2. On matched windows ($n=169$), M-JEPA Ens attains CRPS 6.81 against 6.89 mm/day and positive Brier skill (+0.05 against -0.04), from a 15.4M-parameter model on one 16 GB consumer GPU. The CRPS margin is resolved but slight: $\Delta = -0.08$, 95% CI $[-0.16, -0.001]$ over $n=1183$ paired cases. In exchange, IFS Ens keeps a clear advantage on neighborhood skill (Table 1), the first appearance of a sharpness-versus-overlap trade-off that recurs throughout.

Figure 2a disaggregates M-JEPA Ens’s member-FSS advantage over IFS Ens (matched $n=169$) by lead, at the India Meteorological Department’s operational 24-hour rainfall classification (Falga and Wang 2022), Heavy (64.5 mm) and




Figure 2: Probability-branch diagnostics, matched $n=169$ population. (a) Member-FSS at IMD Heavy/Very Heavy thresholds by lead. (b) Radial power spectral density, M-JEPA Ens vs. IFS Ens vs. observed.

Very Heavy (115.6 mm); the margin is larger at long leads than at short (Heavy: $1.4\times$ at lead 1 to $1.8\times$ at lead 7; Very Heavy: $1.3\times$ to $4.1\times$), a placement-tolerant, neighborhood-skill effect distinct from raw point error at extreme intensities. At the third IMD tier, Extremely Heavy (≥ 204.5 mm, not shown), the ordering is lead-dependent (M-JEPA first in each pair): IFS Ens leads at the first two leads (0.069 vs. 0.196; 0.062 vs. 0.144), M-JEPA Ens from lead 3 onward by up to $6.6\times$ at lead 5 (0.046 vs. 0.007), on 27–38 events per lead; pooled, the short-lead deficit dominates the tier (0.056 vs. 0.077).

Figure 2b gives a candidate mechanism: the radial power spectral density of each ensemble’s individual members against observations, pooled over the matched population and all seven leads, reported descriptively rather than as a hypothesis test. Both systems miss the observed spectrum, but in opposite directions as wavenumber increases. IFS Ens is closest at the largest scale (48% of observed power) and falls to 17% at the finest resolved wavenumber, the smoothing signature in which mass-conserving blur spreads a sharp feature over a wider area at lower amplitude. M-JEPA Ens starts no better (46%) but its share *rises* with wavenumber to 71–72% at the finest scales, four times IFS Ens’s retention there. Matching fine-scale power shows only that the right *amount* of small-scale variance survives, not that it is placed correctly. Panel (a)’s member-FSS margin, which does credit placement, is the complementary evidence. Estimator and per-radius values: supplementary Section F.6.

5.2 Deterministic skill and metric sensitivity

Case-averaged over the test set ($n=337$ windows, seven leads), the deterministic branch attains RMSE 16.24 mm/day and spatial anomaly correlation 0.31 against persistence’s 23.46 and 0.18, exceeding it on both at every lead; persistence nonetheless retains a small advantage at the 3° and 5° neighborhood widths (0.49/0.56 against 0.47/0.53), the same sharpness-versus-overlap trade-off that separates M-JEPA Ens from IFS Ens. Against the operational archives, restricted to the cells the tracked patches cover and each on its own matched-coverage subset (266/127/127 distinct target days, from 161/66/66 tracked windows), M-JEPA’s RMSE is lower than IFS’s (18.23 vs. 18.45, leading at six of the seven leads); on ACC and FSS against all three references,

Table 2: Dry-area specificity (correct-negative rate on truly-dry cells, 2 mm threshold) by lead. AIFS and GraphCast share one matched subset, so a single M-JEPA row serves both.

System $\uparrow$	1	2	3	4	5	6	7
M-JEPA (IFS subset)	0.555	0.534	0.516	0.503	0.503	0.492	0.475
IFS	0.540	0.477	0.460	0.452	0.462	0.463	0.454
M-JEPA (AIFS/GC subset)	0.516	0.532	0.521	0.493	0.487	0.480	0.457
AIFS	0.526	0.497	0.479	0.471	0.443	0.427	0.430
GraphCast	0.424	0.414	0.400	0.387	0.378	0.366	0.377

and on RMSE against AIFS and GraphCast, the references lead (supplementary Table S12).

This split reflects metric sensitivity as much as skill: under a scale-invariant ACC a forecast that sheds fine-scale detail can score well by being smooth, and both learned archives show the signature. AIFS self-smooths progressively over the horizon (roughness ratio $0.396 \rightarrow 0.323$); GraphCast is uniformly over-smooth from lead 1 and gains instead as the truth simplifies, observed roughness falling 22% as the tracked system ages. At case level each (window, lead) case’s ACC correlates negatively with its own observed roughness (Pearson $r = -0.28$ IFS, -0.57 AIFS, -0.57 GraphCast), so rougher targets score lower regardless of lead (supplementary Section F.2). The FSS margin carries a further confound, since the archives are regridded from 0.25° onto our 1° grid, smoothing them before scoring. Two structural asymmetries remain regardless. Each reference initializes from an analysis-quality state at 6-hourly or finer cadence (Lam et al. 2023; Lang et al. 2024), against a frozen daily latent; and each forecasts the full domain, against one co-moving patch.

On metrics that score sharpness directly the ordering reverses. Measured by dry-area specificity, the correct-negative rate on truly-dry cells at the 2 mm threshold, the deterministic branch exceeds IFS and GraphCast at every lead and AIFS at six of seven, losing only at lead 1 (Table 2). Figure 3 shows the same contrast on a single day.

5.3 Event lifecycle and amplitude evolution

The decoded forecast tracks a convective event’s evolution, not merely its instantaneous intensity, on two independent axes. On *position*, comparing the *sign* of the precip-weighted centroid’s drift relative to lead 1, the forecast matches the observed direction more often than the 50% chance rate at every lead (pooled 60.5% northward and 59.0% zonal, $n=2022$, binomial $p < 10^{-4}$ on both axes), rising to 65.0%/64.0% once near-zero observed movement is excluded, so the signal strengthens rather than weakens on genuine motion. M-JEPA still clears chance on the $n=161$ subset where IFS also has a matched forecast (60.6% N/S, 56.1% E/W, $p < 10^{-3}$). IFS is nonetheless the more accurate of the two there (68.0%, 59.0%), as its full-physics initialization and sub-daily cadence would predict. On *amplitude*, the model reproduces the decay both across the forecast horizon (lead 7 relative to lead 1: $0.74\times$ forecast against $0.73\times$ observed) and across an event’s own lifecycle (-0.0253 per day against -0.0247 observed, a 2% over-decay, on a dry bias of 2.1 mm/day). Reproducing lifecycle decay matters here because it is the




Figure 3: Five-model spatial comparison on a matched target day (2024-08-04). Columns: GPM observations, M-JEPA, IFS, AIFS, GraphCast; rows: forecasts issued 1, 3, 5, 7 days ahead. Operational fields are single deterministic members cropped at the same Lagrangian patch, so this compares placement, not probabilistic skill; dashed box marks the native patch extent.

same aging signal that inflates the archives’ ACC: the model follows the observed simplification rather than being scored by it.

Against those two successes the latent’s own trajectory is more limited. Linear Centered Kernel Alignment (CKA; Kornblith et al. 2019) between the context and rolled latents falls from 0.94 at lead 1 to 0.79 by lead 5 and then plateaus, so the rollout transports the representation rather than copying it forward while its trajectory saturates (supplementary Section D.3). Consistently, beyond day 16, where the sample thins to the longest-lived tracks, the observed composite continues falling to $0.68\times$ its day-5 value while the decoded composite stops at $0.75\times$, so the forecast under-continues the decay of the longest-lived systems (supplementary Section F.5).

5.4 Latent variable structure

The five proxies survive pretraining in a readable, physically organized form. Linear probes on the token-averaged context latent recover each proxy’s patch-mean with R^2 between 0.67 (zonal wind) and 0.85 (geopotential), and the content is *linearly* accessible rather than hidden in a nonlinear code, since a multilayer probe tuned by cross-validation, with model selection never touching the test years, improves on the linear probe by at most $+0.02$, on none of the five significantly (supplementary Section D.2). The five probe directions are also close to mutually orthogonal (mean absolute off-diagonal cosine 0.15), and the two largest overlaps are physically interpretable: geopotential with the OLR envelope ($+0.38$) and

Table 3: Isolating the source of transfer. The first two rows share architecture, recipe and probability head, so only the pretraining *signal* differs; the last three are conventional networks given the same rainfall history. CRPS in mm/day, no recalibration, on the full test set ($n=337$) except the rain-only control, which rainfall sparsity leaves a smaller held-out population ($n=118$). Table 1’s 6.81 is this same model on the $n=169$ windows the IFS ensemble also covers.

Pretraining signal	Params	CRPS ↓	BSS ↑
Five proxies	9.7M	5.54	+0.18
Rainfall only, same recipe	9.6M	7.52	−0.04
U-Net (rain history)	351K	7.38	−0.62
ConvLSTM (rain history)	22K	8.33	−1.07
ViT (rain history)	836K	8.65	−0.99

humidity with the meridional wind (+0.30), corresponding to convection organizing beneath lowered geopotential and to the low-level southerly flow that moistens the envelope’s poleward flank, recovered without variable labels.

The same latent separates extreme-rain windows before any decoding. A logistic probe detecting whether a window reaches a widespread-heavy-rain day (area-mean ≥ 15 mm/day at any lead, 27% base rate, $n=337$) attains ROC-AUC 0.67 and average precision 0.41. Modest but real, that separation locates the extreme-skill result of Section 5.1 partly in the representation, not solely in the decoder.

5.5 Predictive information in the latent rollout

To test whether the rollout is a genuine *forecast* rather than a re-encoding of the present, we regress area-mean rainfall on day $t+d$ from the rolled latent $\hat{z}_{t+d}$ and from a same-capacity control, the frozen context-end latent z_t reused at every lead, which would match it if the rollout added nothing. Instead the rolled latent decodes future rainfall better at *every* lead (pooled R^2 0.174 against 0.084, margin +0.090, 95% paired-bootstrap CI [+0.011, +0.178], 10^4 resamples; per-lead breakdown, supplementary Section D.1), peaking at lead 3 (+0.139, where the static probe falls below the unconditional-mean baseline) and decaying to +0.048 by lead 7. Individually, only the pooled margin and lead 1 exclude zero: the per-lead intervals are wide at this test-window count, so the claim rests on the pooled statistic. Even so, the rollout carries decodable information about the future that the present-state encoding lacks, and is therefore predicting rather than autoencoding.

5.6 Attribution of the transfer

Four checks attribute the foregoing to the pretraining *signal* rather than the architecture (supplementary Section E): a randomly initialized backbone of identical architecture (trained exceeds random by +0.128 pooled R^2), a second held-out target in future OLR (margin +0.126), a leave-one-proxy-out ablation in which meridional wind is the only individually load-bearing proxy, and a tracker-sensitivity sweep.

The principal control replaces the pretraining signal itself. Applying the identical JEPA architecture and recipe to the

rainfall field, holding everything else fixed, raises CRPS error by 36% (7.52 against 5.54 mm/day) and collapses Brier skill from +0.18 to −0.04 (Table 3); the two are indistinguishable only on neighborhood skill (FSS10 0.53 and 0.54). The mechanism is data sparsity: rainfall is intermittent, so the same tracking yields only 226 valid pretraining windows against 3885 for the all-season proxies. Even so, the control outperforms three conventional rain-history networks on the same patches and splits, all of which have *negative* Brier skill across two orders of magnitude in parameter count. What makes the representation transfer is therefore the proxy choice, not the architecture or the parameter budget.

6 Limitations and Scaling Outlook

Our primary limitation concerns temporal rollout dynamics rather than representation capacity. The latent trajectory saturates by lead 5 (Section 5.3), so decoded forecasts under-predict the observed decay of long-lived systems. Such damping is expected under squared-error regression, which contracts toward the conditional mean as uncertainty grows; isolating loss-induced damping from representational limits would require an ensemble-valued rollout.

The Lagrangian frame also bounds the benchmark scope. Metrics are evaluated only over tracked convective patches, with operational references cropped into co-moving frames, and admitting only tracked systems caps the test set at 337 windows and 169 matched days on a single seed. The headline CRPS advantage is resolved on this sample; broader generalization requires further evaluation.

These constraints indicate an information-limited rather than an idea-limited regime. Under a $17\times$ reduction in pre-training data volume the rain-only control degrades gracefully, still outperforming three conventional rain-history networks (Section 5.6). Overcoming lead-5 saturation therefore calls for denser proxy inputs, at higher temporal frequency and over more vertical levels and a broader domain, rather than a larger parameter count (supplementary Section G).

7 Conclusion

Self-supervised pretraining on continuous atmospheric proxies enables M-JEPA to learn transferable subseasonal dynamics without rainfall supervision. Across a seven-day horizon, the frozen latent rollout decodes future rainfall better than a static present-state baseline (pooled R^2 margin +0.090), whereas direct precipitation training incurs 36% higher CRPS error (7.52 vs. 5.54 mm/day), confirming the value of proxy representations.

Against the 51-member ECMWF operational ensemble, M-JEPA attains a small but resolved advantage in fair CRPS (6.81 vs. 6.89 mm/day) and positive Brier skill (+0.05 vs. −0.04). These gains concentrate at heavy rainfall thresholds (member-FSS advantage growing $1.4\times \rightarrow 1.8\times$ over the horizon) and fine spatial scales (71–72% spectral power retention vs. 17%), while the ensemble retains the advantage on neighborhood skill and the deterministic references on point metrics. Taken together, this demonstrates that latent cross-variable learning can capture physical dynamics across disjoint targets.

References

- Anirudh, K. M.; Raj, P.; Sandeep, S.; Kodamana, H.; and Sabeerali, C. T. 2025. A Skillful Prediction of Monsoon Intraseasonal Oscillation Using Deep Learning. *Journal of Geophysical Research: Machine Learning and Computation*, 2(2): e2024JH000504.
- Assran, M.; Duval, Q.; Misra, I.; Bojanowski, P.; Vincent, P.; Rabbat, M.; LeCun, Y.; and Ballas, N. 2023. Self-Supervised Learning from Images with a Joint-Embedding Predictive Architecture. In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR)*.
- Assran, M.; et al. 2025. V-JEPA 2: Self-Supervised Video Models Enable Understanding, Prediction and Planning. arXiv:2506.09985.
- Bach, E.; Krishnamurthy, V.; Mote, S.; Shukla, J.; Sharma, A. S.; Kalnay, E.; and Ghil, M. 2024. Improved Sub-Seasonal Prediction of South Asian Monsoon Rainfall Using Data-Driven Forecasts of Oscillatory Modes. *Proceedings of the National Academy of Sciences*, 121(15): e2312573121.
- Banerjee, B. K.; Sharma, D.; Sujith, R. I.; et al. 2026. A Deep Learning Earth System Model Simulation of Indian Monsoon Intraseasonal and Interannual Variability. arXiv:2607.01676.
- Bardes, A.; Ponce, J.; and LeCun, Y. 2022. VICReg: Variance-Invariance-Covariance Regularization for Self-Supervised Learning. In *International Conference on Learning Representations (ICLR)*.
- Bi, K.; Xie, L.; Zhang, H.; Chen, X.; Gu, X.; and Tian, Q. 2023. Accurate medium-range global weather forecasting with 3D neural networks. *Nature*, 619(7970): 533–538.
- Bodnar, C.; Bruinsma, W. P.; Lucic, A.; Stanley, M.; Vaughan, A.; Brandstetter, J.; Garvan, P.; Riechert, M.; Weyn, J. A.; Dong, H.; Gupta, J. K.; Thambiratnam, K.; Archibald, A. T.; Wu, C.-C.; Heider, E.; Welling, M.; Turner, R. E.; and Perdikaris, P. 2025. A Foundation Model for the Earth System. *Nature*, 641: 1180–1187. ArXiv:2405.13063.
- Chen, L.; Su, X.; Zhong, X.; and Li, H. 2026. QuantWeather: Quantile-Aware Probabilistic Forecasting for Subseasonal Precipitation. arXiv:2605.10297.
- Dourado Filho, L. A.; da Silva Neto, A. M.; Miyaguchi, A.; Pereira David, R.; Tripodi Calumby, R.; and Picek, L. 2025. DINOv3 as a Frozen Encoder for CRPS-Oriented Probabilistic Rainfall Nowcasting. arXiv:2511.10894.
- Falga, R.; and Wang, C. 2022. The Rise of Indian Summer Monsoon Precipitation Extremes and Its Correlation with Long-Term Changes of Climate and Anthropogenic Factors. *Scientific Reports*, 12: 11985.
- Fan, H.; Bai, L.; Fei, B.; Xiao, Y.; Chen, K.; Liu, Y.; Qu, Y.; Ling, F.; and Gentile, P. 2025. Physically Consistent Global Atmospheric Data Assimilation with Machine Learning in Latent Space. *Science Advances*. ArXiv:2502.02884, doi:10.1126/sciadv.aea4248.
- Ferro, C. A. T.; Richardson, D. S.; and Weigel, A. P. 2008. On the Effect of Ensemble Size on the Discrete and Continuous Ranked Probability Scores. *Meteorological Applications*, 15(1): 19–24.
- Gneiting, T.; and Raftery, A. E. 2007. Strictly Proper Scoring Rules, Prediction, and Estimation. *Journal of the American Statistical Association*, 102(477): 359–378.
- Goswami, B. N.; Ajayamohan, R. S.; Xavier, P. K.; and Sengupta, D. 2003. Clustering of Synoptic Activity by Indian Summer Monsoon Intraseasonal Oscillations. *Geophysical Research Letters*, 30(8).
- He, S.; Li, X.; DelSole, T.; Ravikumar, P.; and Banerjee, A. 2021. Sub-Seasonal Climate Forecasting via Machine Learning: Challenges, Analysis, and Advances. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 35, 169–177.
- Henderson, P.; Islam, R.; Bachman, P.; Pineau, J.; Precup, D.; and Meger, D. 2018. Deep Reinforcement Learning that Matters. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 32.
- Hersbach, H.; et al. 2020. The ERA5 Global Reanalysis. *Quarterly Journal of the Royal Meteorological Society*, 146(730): 1999–2049.
- Huffman, G. J.; Bolvin, D. T.; and Nelkin, E. J. 2015. Integrated Multi-satellite Retrievals for GPM (IMERG). *NASA/GSFC Algorithm Theoretical Basis Document*.
- Kornblith, S.; Norouzi, M.; Lee, H.; and Hinton, G. 2019. Similarity of Neural Network Representations Revisited. In *International Conference on Machine Learning (ICML)*, volume 97 of PMLR, 3519–3529.
- Lam, R.; Sanchez-Gonzalez, A.; Willson, M.; Wirnsberger, P.; Fortunato, M.; Alet, F.; Ravuri, S.; Ewalds, T.; Eaton-Rosen, Z.; Hu, W.; Meroze, A.; Hoyer, S.; Holland, G.; Vinyals, O.; Stott, J.; Pritzel, A.; Mohamed, S.; and Battaglia, P. 2023. Learning Skillful Medium-Range Global Weather Forecasting. *Science*, 382(6677): 1416–1421.
- Lang, S.; Alexe, M.; Chantry, M.; Dramsch, J.; Pinault, F.; Raoult, B.; Clare, M. C. A.; Lessig, C.; Maier-Gerber, M.; Magnusson, L.; Boussetta, S.; Dueben, P. D.; Pappenberger, F.; and Rabier, F. 2024. AIFS: ECMWF's Data-Driven Forecasting System. *arXiv preprint arXiv:2406.01465*.
- Lang, S.; Alexe, M.; Chantry, M.; Dramsch, J.; Pinault, F.; Raoult, B.; Clare, M. C. A.; Lessig, C.; Maier-Gerber, M.; Magnusson, L.; et al. 2026. AIFS-CRPS: Ensemble Forecasting Using a Model Trained with a Loss Function Based on the Continuous Ranked Probability Score. *npj Artificial Intelligence*.
- Lee, H.-T. 2025. NOAA Climate Data Record (CDR) of Daily Outgoing Longwave Radiation (OLR), Version 2.0. NOAA National Centers for Environmental Information.
- Lin, H.; Gao, Z.; Xu, Y.; Wu, L.; Li, L.; and Li, S. Z. 2022. Conditional Local Convolution for Spatio-Temporal Meteorological Forecasting. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 36, 7470–7478.
- Mardani, M.; Brenowitz, N.; Cohen, Y.; Pathak, J.; Chen, C.-Y.; Liu, C.-C.; Vahdat, A.; Kashinath, K.; Kautz, J.; and Pritchard, M. 2025. Residual Corrective Diffusion Modeling for km-Scale Atmospheric Downscaling. *Communications Earth & Environment*, 6: 124.

- Meredith, E. P.; Ulbrich, U.; and Rust, H. W. 2023. Cell Tracking of Convective Rainfall: Sensitivity of Climate-Change Signal to Tracking Algorithm and Cell Definition (Cell-TAO v1.0). *Geoscientific Model Development*, 16(3): 851–867.
- Nam, H.; Le Lidec, Q.; Maes, L.; LeCun, Y.; and Balestrieri, R. 2026. Causal-JEPA: Learning World Models through Object-Level Latent Masking. In *International Conference on Machine Learning (ICML)*.
- Nathaniel, J.; Qu, Y.; Nguyen, T.; Yu, S.; Busecke, J.; Grover, A.; and Gentine, P. 2024. ChaosBench: A Multi-Channel, Physics-Based Benchmark for Subseasonal-to-Seasonal Climate Prediction. In *Advances in Neural Information Processing Systems (NeurIPS), Datasets and Benchmarks Track*, volume 37.
- Naveau, P.; Huser, R.; Ribereau, P.; and Hannart, A. 2016. Modeling Jointly Low, Moderate, and Heavy Rainfall Intensities Without a Threshold Selection. *Water Resources Research*, 52(4): 2753–2769.
- Nguyen, T.; Brandstetter, J.; Kapoor, A.; Gupta, J. K.; and Grover, A. 2023. ClimaX: A Foundation Model for Weather and Climate. In *Proceedings of the 40th International Conference on Machine Learning (ICML)*, volume 202 of *PMLR*.
- Prein, A. F.; Feng, Z.; Fiolleau, T.; Moon, Z. L.; Núñez Ocasio, K. M.; Kukulies, J.; Roca, R.; Varble, A. C.; Rehbein, A.; Liu, C.; Ikeda, K.; Mu, Y.; and Rasmussen, R. M. 2024. Km-Scale Simulations of Mesoscale Convective Systems Over South America—A Feature Tracker Intercomparison. *Journal of Geophysical Research: Atmospheres*, 129(8).
- Price, I.; Sanchez-Gonzalez, A.; Alet, F.; Andersson, T. R.; El-Kadi, A.; Masters, D.; Ewalds, T.; Stott, J.; Mohamed, S.; Battaglia, P.; Lam, R.; and Willson, M. 2025. Probabilistic Weather Forecasting with Machine Learning. *Nature*, 637(8044): 84–90.
- Rasp, S.; Hoyer, S.; Merose, A.; Langmore, I.; Battaglia, P.; Russell, T.; Sanchez-Gonzalez, A.; Yang, V.; Carver, R.; Agrawal, S.; et al. 2024. WeatherBench 2: A Benchmark for the Next Generation of Data-Driven Global Weather Models. *Journal of Advances in Modeling Earth Systems*, 16(6): e2023MS004019.
- Roberts, N. M.; and Lean, H. W. 2008. Scale-Selective Verification of Rainfall Accumulations from High-Resolution Forecasts of Convective Events. *Monthly Weather Review*, 136(1): 78–97.
- Rühling Cachay, S.; Zhao, B.; Joren, H.; and Yu, R. 2023. DYffusion: A Dynamics-Informed Diffusion Model for Spatiotemporal Forecasting. In *Advances in Neural Information Processing Systems (NeurIPS)*, volume 36.
- Shi, X.; Chen, Z.; Wang, H.; Yeung, D.-Y.; Wong, W.-K.; and Woo, W.-c. 2015. Convolutional LSTM Network: A Machine Learning Approach for Precipitation Nowcasting. In *Advances in Neural Information Processing Systems (NeurIPS)*.
- Stephens, G. L.; L'Ecuyer, T.; Forbes, R.; Gettelman, A.; Golaz, J.-C.; Bodas-Salcedo, A.; Suzuki, K.; Gabriel, P.; and Haynes, J. 2010. Dreary State of Precipitation in Global Models. *Journal of Geophysical Research: Atmospheres*, 115(D24): D24211.
- Subich, C.; Husain, S. Z.; Šeparović, L.; and Yang, Y. 2025. Fixing the Double Penalty in Data-Driven Weather Forecasting Through a Modified Spherical Harmonic Loss Function. In *International Conference on Machine Learning (ICML)*.
- Suhas, E.; Neena, J. M.; and Goswami, B. N. 2013. An Indian Monsoon Intraseasonal Oscillation (MISO) Index for Real-Time Monitoring and Forecast Verification. *Climate Dynamics*, 40: 2605–2616.
- Thil, L.; Read, J.; Kaddah, R.; and Doquet, G. 2026. Subspace-Decomposed JEPAs: Disentangling Progression and Content in Latent World Models. arXiv:2605.31111.
- Wen, P.; He, M.; Filippi, P.; Zhao, N.; Zhang, F.; Bishop, T. F. A.; Wang, Z.; and Hu, K. 2026. DuoCast: Duo-Probabilistic Diffusion for Precipitation Nowcasting. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 40, 39442–39450.
- Wheeler, M.; and Kiladis, G. N. 1999. Convectively Coupled Equatorial Waves: Analysis of Clouds and Temperature in the Wavenumber-Frequency Domain. *Journal of the Atmospheric Sciences*, 56(3): 374–399.
- Wilson, T.; Tan, P.-N.; and Luo, L. 2022. DeepGPD: A Deep Learning Approach for Modeling Geospatio-Temporal Extreme Events. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 36, 4245–4253.
- Wisinski, E. F.; Molina, M. J.; Hall, K. J. C.; Bao, H.; Mahajan, S.; Rosenbloom, N.; and Fasullo, J. 2026. What's in the latent space? Exploring coupled tropical Pacific variability within a Multi-branch β -Variational Autoencoder. arXiv:2604.07137.
- Zhang, Z.; Wang, Y.; Guan, Z.; Yang, Y.; Shi, B.; Zong, T.; Yi, H.; Chao, G.; Chen, X.; Yang, T.; Bao, C.; Yu, T.; Zhou, J.; and Xu, J. 2026. Delta-JEPA: Learning Action-Sensitive World Models via Latent Difference Decoding. arXiv:2606.31232.
- Zhao, S.; Liu, F.; Zhang, X.; Chen, H.; Han, T.; Gong, J.; Tao, R.; Xiao, P.; Bai, L.; and Ouyang, W. 2026. Transforming Weather Data from Pixel to Latent Space. In *International Conference on Machine Learning (ICML)*, volume 306 of *PMLR*.

Supplementary Material

A Data and Problem Setup

We predict the next seven daily precipitation fields over the South Asian monsoon domain (9.5°S–29.5°N, 60.5–99.5°E) on a 1° grid. Inputs are five daily proxy fields: outgoing longwave radiation (OLR) plus specific humidity, zonal and meridional wind, and geopotential at 850 hPa (Hersbach et al. 2020). The target is satellite precipitation (Huffman, Bolvin, and Nelkin 2015). All fields are regridded to the target grid, deseasonalized, and z -scored using training-year statistics only. Rather than a fixed Eulerian window, we extract *Lagrangian patches* that follow convective systems identified by a classical detection-and-tracking pipeline on band-passed OLR; the full detection and tracking algorithm and its settings are given in Section A.1, and the trajectories of the resulting test population in Figure S2. This retrospective, zero-phase (non-causal) bandpass filter is used *only* to define patch geometry (which convective system to follow, and where to center a window), sharing only the 20–90 day frequency band with the convention used to define standard real-time MISO monitoring indices (Suhas, Neena, and Goswami 2013), not their causal (forward-only) design; the tracking filter itself is never used to construct a model input value. The model forecasts the *intensity and structural evolution* of rainfall within the patch, not the patch trajectory. The tracker-supplied center coordinate fixes where each evaluation window sits but is never a predicted quantity, so it acts as an isolated oracle for window placement, and no forecast metric credits the model for knowing the system’s future position. All five daily proxy fields the encoder consumes are raw and unfiltered; an optional bandpass-filtered auxiliary channel exists in a strictly causal (forward-only) variant but is disabled by default. A tight foreground patch captures the convective core; a wider background patch captures the environmental circulation, with a purely spatial (not temporal) Gaussian smoothing of OLR that carries no information from other time steps. The analysis domain is itself 40×40 cells, the same nominal size as the background patch, so the co-moving crop is taken from a zero-padded field extended by half a patch width on every side rather than from the raw domain. A background patch therefore remains centered on the tracked centroid wherever the system sits, and coincides with the domain only for a system at dead center; for systems near an edge the out-of-domain portion is zero, which after deseasonalizing and z -scoring is the climatological mean rather than an artificial extremum. The foreground patch, at 20×20 , is well inside the domain for all retained tracks. The sensitivity of the comparative conclusions to the two tracker-geometry parameters is established separately in Section E.5.

A.1 Object Detection and Tracking

The detect-and-track pipeline runs on daily OLR over the all-variable box domain, all calendar months, at 1° resolution.

Signal. Each pixel’s OLR time series is band-pass filtered to the 20–90 day intraseasonal band with a third-order Butterworth filter applied zero-phase (forward–backward), the stan-

dard band for isolating the intraseasonal envelope (Wheeler and Kiladis 1999); missing values are linearly interpolated before filtering. This retrospective filter defines patch geometry only and never enters a model input.

Detection. On each day, deep-convective cores are the strongly negative anomalies: cells below the 10th-percentile quantile of that day’s band-passed field are selected, Gaussian-smoothed ($\sigma=1$ cell) to bridge fragmented cores, and grouped into connected components (`scipy.ndimage.label`); components smaller than 30 cells are discarded. Each object’s location is its *intensity-weighted* centroid, each cell weighted by its absolute OLR anomaly, so the center follows the deepest convection and not the surrounding stratiform anvil.

Tracking. Detections are linked into tracks by greedy nearest-cost assignment within each year. Matching object j on day t to an existing track i costs

$$c_{ij} = w_d \|\mathbf{x}_j - \hat{\mathbf{x}}_i\| + w_a \frac{|A_j - A_i|}{A_i} + w_t (\Delta t - 1) - w_n \Delta \text{lat},$$

with distance weight $w_d=1.0$, fractional-area-change weight $w_a=0.5$, time-gap weight $w_t=1.0$, and direction weight $w_n=0$. The direction term is identically zero, because tracking is direction-blind, so northward propagation is never rewarded in the assignment cost. Here $\hat{\mathbf{x}}_i$ is the track position linearly extrapolated from its last step, which bridges up to four missed detections (`gap_days=4`); candidate matches are restricted to a $5^\circ/\text{day} \times \Delta t$ search radius.

Retention. A track is kept if it spans at least 12 days, the five-day context plus seven-day forecast horizon. The strict-MISO physical filters of the original tracker (minimum net northward displacement, minimum and maximum propagation speed, and an ocean-genesis mask) are disabled here, which retains a larger and more varied training population. The test-year population (149 tracks, before the span filter) has median net northward displacement $+3.5^\circ$, or under 0.3° per day over the retained window, with 60% of tracks propagating poleward (2.4 binomial standard errors above chance at this sample size). Poleward motion therefore emerges from the data and not from the direction-blind tracking cost, as a tendency and not a selection criterion, and the median track is slower than a canonical MISO envelope. Each retained centroid then anchors the two-scale (tight foreground, wide background) co-moving crop described above. Figure S1 follows one long-lived envelope through this procedure, and Figure S2 shows the trajectories of the full test-year population.

Splits are by year block to avoid autocorrelation leakage: self-supervised pretraining on 1979–2017, decoder training on 1998–2013 (667 windows, trained with ± 2 -grid-cell spatial-jitter augmentation), validation on 2014–2017, and a held-out test set of 2018–2025 (337 windows).

B Model Architecture and Training

Pipeline overview and component diagram: main-paper Figure 1. Component structure, symbols and losses are given in main-paper Section 4; this section records the sizes and optimization settings behind them.




Figure S1: Lagrangian tracking of one long-lived convective envelope (track 2766, genesis 2021-09-07, ocean, equatorial Indian Ocean; 45-day duration; net northward displacement $+25.6^\circ$). **(a)** The tracked centroid trajectory from genesis to terminus, colored by elapsed time; diamonds mark the six days shown at right. **(b)** The $20^\circ \times 20^\circ$ bandpassed OLR anomaly patch centered on the moving centroid at each marked day, its border colored to match that day on the trajectory, illustrating how the Lagrangian frame re-centers on the system as it propagates from the equatorial Indian Ocean to the Bay of Bengal. This track is illustrative rather than typical, since the retained test population’s median net northward displacement is $+3.5^\circ$. This patch geometry, not the raw OLR values shown here, is what the tracker supplies; the encoder itself consumes unfiltered fields (Section A).

B.1 Component Sizes

The frozen Stage-1 stack holds 9.7M parameters: 4.8M in the context encoder f_θ (six pre-norm transformer blocks, eight heads, width 256, applied per day over the joint 50-token set) and 4.9M in the block-causal autoregressive predictor g_ϕ (six blocks, eight heads, width 256, residual output with a zero-initialized projection). The target encoder f_ξ is an architectural copy of f_θ updated only by the exponential moving average, and the auxiliary OLR reconstruction head is discarded after pretraining; neither is part of the released model. Stage 2 holds 5.7M parameters: a 2.1M shared trunk (two token projections, a 3×3 scale fusion at the 5×5 token grid, two transposed-convolution upsampling blocks to 20×20 at 128 channels, and a 1×1 OLR readout), 0.4M in the rain-history and geography conditioning encoders, a 3.0M probability branch (U-Net width 96, conditioning re-injected at every scale), and a 0.3M deterministic branch (the same U-Net topology at width 32, without the noise and time inputs); each component is rounded independently, so the parts sum to 5.8M against the 5.7M total. Inference cost is itemized in Section H.

The trunk and probability branch are trained jointly on main-paper Equation (9); only the Stage-1 stack is frozen. The deterministic branch is trained afterward, as a sepa-

rate readout on that same trunk now held frozen, on main-paper Equation (10); the two branches never share gradients into the trunk. This sequencing is deliberate, because every deterministic-branch trunk trained end-to-end under the deterministic loss alone tops out at pooled spatial ACC 0.255–0.271 regardless of curriculum, while the frozen, distributionally-trained trunk borrowed here reaches 0.292 on the same pooled convention, consistent with a distributional objective building more useful deterministic features than a deterministic loss does on its own. Both branches’ outputs are used directly, with no post-hoc recalibration. Retraining the deterministic branch from three random seeds on the fixed frozen trunk (checked under this project’s earlier pooled scoring convention, since superseded by the case-averaged convention used throughout the main text) moved the reported numbers within ± 0.007 mm/day on RMSE, ± 0.008 on spatial ACC, and ± 0.010 –0.020 on FSS at the 1° – 5° neighborhood widths, so RMSE and ACC gains do not rest on a single seed. The released checkpoint is seed 1410, which is also the best of the three; we report the spread above rather than that seed’s score alone, since selecting and reporting the best of several seeds inflates apparent skill (Henderson et al. 2018). The seed spread is small enough that the tabulated values stand whichever seed is used, with one exception noted




Figure S2: Lagrangian trajectories of all 149 convective OLR objects tracked in the held-out test population (genesis years 2018–2025, JJAS), the population underlying every forecast-skill number reported in this paper. Black circles mark genesis and red crosses mark terminus, with genesis concentrated across the equatorial Indian Ocean and Bay of Bengal; net northward displacement is positive on balance (median $+3.5^\circ$, Section A). The 1979–2017 self-supervised pretraining population (711 tracks, of which the 2014–2017 subset also serves as validation) shows the same qualitative spatial pattern.

below. Under the case-averaged convention, persistence out-right leads the deterministic branch’s FSS at 3° and 5° neighborhood width (main-paper Section 5.2; Table S10 below), superseding the earlier pooled-convention finding that this margin was merely seed-sensitive. That FSS comparison is the exception noted above: the seed spread was measured under the pooled convention and has not been re-measured under the case-averaged one, so we treat the persistence-versus-M-JEPA neighborhood comparison as seed-sensitive and do not claim it.

B.2 Optimization Settings

Table S1 lists the settings for both stages. Stage 2 uses a fixed epoch budget and the final-epoch checkpoint, with no validation-based selection, since on jointly trained trunks the validation neighborhood score swings between evaluations, so best-checkpoint selection returns half-trained models.

The controlled comparison of main-paper Table 3 uses the same flow-based probability head and the same Stage-2 recipe on both frozen backbones, so that any skill difference reflects the pretraining signal rather than the decoder. Because the rain-only control’s decoder-training population (226 windows, against 667) is small, its loader is given the same ± 2 -cell spatial-jitter augmentation used in the main pipeline. The three rain-history baselines have no frozen trunk to attach a branch to and are trained end to end instead,

	Stage 1 (pretraining)	Stage 2 (readout)
Objective	Eq. (4)	Eq. (9)
Optimizer	AdamW	AdamW
Learning rate	5×10^{-4}	10^{-3}
Weight decay	0.05	10^{-4}
Warmup	10 epochs	none
Epochs	100	80
Batch size	32	32
Gradient clipping	none	1.0
Checkpoint	final epoch	final epoch
Augmentation	± 2 -cell jitter	± 2 -cell jitter
Rollout horizon	curriculum $2 \rightarrow 7$	fixed at 7
VICReg (var/cov)	1.0 / 0.04	not applicable
Auxiliary weights	OLR 0.1	energy score 0.1
Trainable parameters	9.7M	5.7M
Windows	3885 (all season)	667 (rain-labeled)
Hardware	single NVIDIA RTX A4000 (16 GB)	

Table S1: Optimization settings for both training stages. Equation numbers refer to the main paper. Stage 1 sees no rainfall at any point; Stage 2 trains only the trunk and the two branches, on the frozen Stage-1 rollout.

with a standard zero-inflated occurrence-plus-intensity loss on the architectures listed in Section E.1.

B.3 Configuration Details for All Compared Systems

Table S2 itemizes the architecture and training configuration for every model compared in this paper. The three rain-history baselines (Section E.1) take a five-day rainfall history as input and predict occurrence and intensity jointly for all seven lead days; all three are trained with AdamW ($\text{lr} = 10^{-3}$, weight decay 10^{-4}) for 60 epochs on the identical tracked patches and splits as the primary model. IFS, AIFS, and GraphCast are not trained by us; they are public operational (IFS) or research (AIFS, GraphCast) forecast archives, cropped to the same Lagrangian patch and lead as our own predictions and scored identically as single deterministic members (Section C).

C Verification Protocol and Scoring Conventions

The main paper states these conventions once and then uses them without restatement.

Patch frame. Every score is computed on the cells a tracked Lagrangian patch covers, not on the full model domain. The operational archives are cropped to the same cells and the same lead before scoring, so all systems are compared on identical geometry. Which cells count is fixed by an *observation-derived* patch mask: the valid-observation footprint of the target field defines the scored cells, so the mask is set by the data rather than by any model’s output, and no system can be advantaged by the choice of scoring region. Full-domain rescoreing of the archives, for contrast, is in Section F.2.

Model	Architecture	Training
M-JEPA	Multiscale transformer context encoder (4.8M) + autoregressive predictor (4.9M), both frozen at readout; shared decoder trunk with two branches (5.7M)	Self-supervised on proxies only, no rain objective; trunk and branches trained on paired rain data, 667 train windows
Rain-only control	Identical architecture, backbone 9.6M against M-JEPA's 9.7M (only the tokenizer's input-channel count differs)	Same recipe as M-JEPA, applied to rainfall alone; 226 pretrain windows
U-Net	Encoder-decoder, 2 downsample / 2 upsample conv stages, skip connections, 32 base channels (351K params)	AdamW, lr 10^{-3} , weight decay 10^{-4} , 60 epochs, rain-history input only
ConvLSTM	Single ConvLSTM cell, 24 hidden channels, closed-loop autoregressive rollout (22K params)	AdamW, lr 10^{-3} , weight decay 10^{-4} , 60 epochs, rain-history input only
ViT	Patch embedding (patch 4, dim 128), 4-layer/4-head transformer encoder (836K params)	AdamW, lr 10^{-3} , weight decay 10^{-4} , 60 epochs, rain-history input only
IFS	Physics-based NWP, operational	Not trained by us; public open-data archive
AIFS	ML emulator (Lang et al. 2024); its ensemble variant is CRPS-trained (Lang et al. 2026)	Not trained by us; public research archive
GraphCast	ML emulator, graph neural network (Lam et al. 2023)	Not trained by us; public research archive

Table S2: Model and training configuration for every system compared in this paper. All M-JEPA, rain-only-control, U-Net, ConvLSTM and ViT models are trained on the identical Lagrangian patches and year-block splits (Section A) so that comparisons isolate the modeling choice, not the data. IFS, AIFS, and GraphCast are external references, evaluated but not trained.

Case-averaging. Each (window, lead) pair is scored individually and the resulting scores are averaged, rather than pooling every cell of every case into one score. Pooling lets the largest and wettest cases dominate; case-averaging weights each forecast situation equally. The project previously used the pooled convention, and where an older pooled number is quoted it is labeled as such.

Probabilistic scores. Brier skill is computed against a patch-local climatology at the 1 mm threshold, following the recommendation that a climatological reference be local to the verification sample rather than global. *Member-FSS* applies the neighborhood fraction skill score (Roberts and Lean 2008) to each ensemble member separately and then averages, so it credits a forecast that places rain within a spatial tolerance rather than requiring exact hits. FSS10 is that score at the 10 mm threshold; the neighborhood width is stated with each table that reports it.

Anomaly correlation. $ACC = \text{corr}(\text{pred} - \text{clim}, \text{obs} - \text{clim})$ is scale-invariant: it rewards spatial pattern match and is blind to amplitude, so a forecast that sheds fine-scale detail can score well precisely by being smooth. That property drives the comparison in main-paper Section 5.2 and is analyzed in Section F.2.

Matched-archive comparisons (M-JEPA vs. ECMWF 51-member) use the NHC (National Hurricane Center) homogeneous-sample convention: two forecasts are paired and compared only on calendar days where both forecasts have complete, valid output. This guards against archive-coverage biases. In this study, $n=169$ days (2018–2025 test period) have both M-JEPA 7-lead and ECMWF ensemble 7-lead output at the reference patch anchor. Windows whose system anchor lies near a domain edge are retained and scored on their observed cells only, instead of being discarded by a naive crop of the reference grid; this is why the matched population is not smaller than the archive extent alone would

suggest.

Fair CRPS estimator: we use the unbiased form $CRPS_{\text{fair}} = \frac{1}{K} \sum_i |x_i - y| - \frac{1}{2K(K-1)} \sum_{i,j} |x_i - x_j|$ (Ferro, Richardson, and Weigel 2008), where K is ensemble size and y is the observation. The biased “energy” form $CRPS_{\text{energy}} = \frac{1}{K} \sum_i |x_i - y| - \frac{1}{2K^2} \sum_{i,j} |x_i - x_j|$ systematically favors larger ensembles, and comparison of unequal-member forecasts with the biased form can flip the sign of the result, a pitfall relevant to any comparison of M-JEPA (16 members) vs. operational ensembles (51 members).

D Extended Probing Results

This section gives the full per-lead numbers and extended analysis behind the probes in main-paper Section 5: predictive information (Section D.1), variable decodability and subspace structure (Section D.2), and CKA trajectory evolution (Section D.3). Section D.4 gives the latent-to-rainfall attribution analysis in full; for space, this one is presented only here rather than summarized in the main paper.

D.1 Predictive Information: Full Results by Lead

Table S3 gives per-lead R^2 for the rolled latent vs. the static-latent control (the frozen context-end latent reused at every lead) across all 7 forecast days, and Figure S3 plots the two curves with their paired margin. The margin is hump-shaped, largest at lead 3 (where the static probe turns negative, below the unconditional-mean baseline), smallest at lead 7 (rollout advantage erodes at long range). The negative R^2 for the static probe at lead 3 indicates the initial latent state underperforms the unconditional-mean predictor at that lead.

D.2 Variable Decodability and Subspace Structure: Extended Analysis

Table S4 reports linear-probe and CV-tuned nonlinear-probe R^2 for the five proxy variables. The nonlinear probe is

Rolled latent decodes future rain better than a static snapshot, at every lead




Figure S3: Rain decodability (R^2 , linear probe) at each lead from the rolled latent (forecast) vs. the frozen present-state control (no forecast). The rolled latent decodes future rainfall better at every lead, by the widest margin at lead 3. Shaded region is the paired rolled-minus-frozen margin, whose pooled value and 95% bootstrap interval (10^4 resamples) are annotated; this paired margin, not the two marginal curves separately, is the statistic the claim rests on.

Lead (days)	Rolled $R^2 \uparrow$	Static $R^2 \uparrow$	Margin $\uparrow$	Margin 95% CI
1	0.341	0.261	+0.079	[+0.004, +0.169]
2	0.198	0.088	+0.110	[-0.003, +0.246]
3	0.105	-0.034	+0.139	[-0.003, +0.292]
4	0.092	0.022	+0.070	[-0.076, +0.227]
5	0.169	0.086	+0.083	[-0.051, +0.226]
6	0.169	0.070	+0.099	[-0.048, +0.263]
7	0.145	0.097	+0.048	[-0.096, +0.203]
Pooled	0.174	0.084	+0.090	[+0.011, +0.178]

Table S3: Rolled vs. static-latent R^2 per lead. Margin peaks at lead 3, consistent with a forecast whose advantage over persistence erodes at long range. The 95% confidence interval is from a 10^4 -replicate paired bootstrap over test windows; the pooled (lead-averaged) margin excludes zero, while individual leads overlap zero given the modest test-window count.

a multilayer perceptron whose architecture and optimizer are chosen by a cross-validated grid search over hidden width $\{128, 256, 512\}$, depth $\{1, 2\}$, learning rate $\{10^{-3}, 3 \times 10^{-4}\}$, and weight decay $\{10^{-3}, 10^{-4}\}$ (24 configurations), with model selection on the validation years (2014–2017) only; the reported nonlinear R^2 is the selected configuration’s *test*-year score, and the test years are never used for selection. The validation years do fall inside the self-supervised pretraining span, so probe selection is not fully independent of the encoder’s training data. That can only favor the tuned nonlinear probe, and the finding here is that the tuned probe does not exceed the linear one, so the contamination works against the conclusion we draw rather than for it. The proxies are strongly linearly decodable (0.67–0.85), and the tuned nonlinear probe does not significantly outperform the linear probe on any of them, and the paired-bootstrap gap confidence interval includes zero for four variables and is significantly *negative* for humidity, with the largest point gain only +0.02 (geopotential, OLR envelope). That the tuned

probe adds nothing indicates the variable content is stored in a linearly accessible form rather than a hidden nonlinear code.

Latent Dimension	Linear $R^2 \uparrow$	CV-tuned MLP $R^2 \uparrow$	Gap [95% CI]
q (humidity)	0.697	0.644	-0.052 [-0.105, -0.007]
u (zonal wind)	0.668	0.642	-0.026 [-0.082, +0.029]
v (merid. wind)	0.679	0.668	-0.011 [-0.062, +0.039]
z (geopotential)	0.849	0.865	+0.016 [-0.005, +0.037]
OLR (smoothed)	0.744	0.763	+0.018 [-0.021, +0.060]

Table S4: Linear vs. CV-tuned nonlinear probe R^2 for the five proxy readout targets, with the paired-bootstrap (10^4 -replicate) 95% CI on the tuned-minus-linear gap. Proxies are strongly and linearly decodable, and the tuned MLP does not significantly exceed the linear probe on any of them. Selected MLP configurations: q , u width 512/depth 1; v width 128/depth 1; z width 256/depth 1; OLR width 256/depth 2.

The five proxy directions occupy near-orthogonal subspaces (mean absolute off-diagonal cosine similarity 0.15 among the proxies alone). Overlaps are physically interpretable, with geopotential and the OLR envelope couple at +0.38, humidity and the meridional wind at +0.30, and the two horizontal wind components at -0.20: deep convection organized beneath lowered geopotential, and moisture advected by the northward monsoon flow.

D.3 CKA Evolution and Latent-Increment Structure

Centered Kernel Alignment (CKA) between the context latent and the rolled latent at each of the seven rollout steps shows the trajectory’s saturation (Table S5), with a rapid fall from 0.944 at lead 1 to 0.794 by lead 5, then plateau. This is an independent signature of damped intraseasonal dynamics in the learned representation.

Lead (days)	CKA (context vs. rollout)
1	0.944
2	0.872
3	0.825
4	0.804
5	0.794
6	0.793
7	0.794

Table S5: CKA between the context latent and the rolled latent at each of the seven rollout steps (leads 1–7). Rapid initial drop (evolution), then plateau (saturation).

Principal-component analysis of latent increment vectors $\hat{z}_{t+d} - z_t$ reveals distributed evolution: the top-3 PCs explain only 16%, 15%, 11% of increment variance respectively, totaling 42%, with no single dominant axis of change. This suggests the rollout evolves through many weakly-coupled modes, consistent with an environment interacting with the tracked system (wind shear, moisture convergence) rather than a single low-dimensional mode of variation.

D.4 Attribution and Saliency: Latent-Space Gradients

We trace how latent content becomes a rainfall prediction, on the current production decoder (shared flow trunk plus the det-on-flow-trunk deterministic head; an earlier version of this analysis used a since-retired decoder head and is superseded by the numbers below). We backpropagate the decoded expected-rain field to the rolled foreground latent tokens and aggregate the gradient magnitude per token on the native 5×5 token grid. The driving token sits one cell off the patch center (north and east of it) and is stable across leads 1–6, shifting to directly east of center at the longest lead (Figure S4a,b); the center token itself is high but never the maximum (91–99% of the peak across all seven leads). The grid’s contrast falls from a coefficient of variation of 0.272 at lead 1 to 0.240 at lead 7 (a $\approx 12\%$ reduction, not fully monotonic), consistent with the mild dispersion expected from the trajectory saturation reported in Section D.3. A complementary input-space saliency map at pixel resolution increases with distance from the patch center rather than peaking at a fixed ring (Figure S4c); this pattern is architecture-dependent and consistent with a convolutional boundary effect in the decoder trunk rather than necessarily reflecting the true convective footprint, so we do not read a specific radius from it. The token-level result is the more direct evidence, since the pretrained latent drives the readout through an off-center, spatially stable set of tokens rather than the exact centroid token alone.

E Controls and Ablations

Main-paper Section 5.6 isolates the source of the transfer with two controls (rainfall-only pretraining and a randomly initialized backbone), a further generalization test on a second held-out target, a leave-one-proxy-out ablation, and a check of sensitivity to the tracker’s detection settings, which establishes that the object-based evaluation frame itself is not an artifact of a particular choice of detection threshold or minimum object area. Each is given in full below, in that order.

E.1 Rainfall-Only Pretraining and Rain-History Baselines

To isolate the value of proxy-based pretraining, we retrain the identical JEPA encoder and predictor using the same self-supervised recipe on rainfall alone, holding architecture and training procedure constant while replacing the five proxy fields with rainfall only. This tests whether the proxy-learning hypothesis is driven by proxy availability (information-limited) or by something about the architecture or loss function (idea-limited).

The main paper’s claim that this rain-only control is stronger than conventional rain-history architectures requires ruling out that such an architecture, given only rainfall history, could do better if better designed or better parameterized. We therefore trained three standard architectures on the identical tracked patches and splits, each taking only a five-day rainfall history as input, with a standard zero-inflated




Figure S4: The readout draws on tokens around, not exactly on, the tracked convective centroid. (a,b) Token-level attribution (gradient of expected rainfall with respect to the rolled foreground latent tokens, native 5×5 token grid, no smoothing, arbitrary units, shared color scale) at leads 1 and 7; the star marks the peak token and the cross the patch’s geometric center; north is up and east is right. The peak sits one cell north-east of center at lead 1 and one cell east of center at lead 7. (c) Radially binned mean input-space saliency (gradient with respect to the raw 20×20 foreground field) against distance from the patch center, per lead: increases toward the patch edge rather than peaking at a fixed ring, a pattern we attribute to the decoder’s convolutional boundary handling rather than to convective structure.

occurrence-plus-intensity loss: a compact U-Net (351K parameters), a ConvLSTM (22K parameters), and a Vision Transformer (836K parameters). Table S6 gives the result, and expands main-paper Table 3 with the neighborhood-skill column.

Under-parameterization is not the cause of the standard architectures’ failure: they fail identically across two orders of magnitude in size, so rainfall history alone is an insufficient basis for this forecasting task at this lead range. The rain-only control’s self-supervised objective, though itself limited by the sparse target signal, still extracts more from that target than any of the three conventional networks.

The failure is invariant to capacity from two directions, which rules out the proxy model outperforming merely because it is larger. Downward, the three rain-history baselines fail identically across 22K–836K parameters, so adding parameters to a rain-only model does not help. Upward, the control’s backbone holds 9.6M parameters against M-JEPA’s 9.7M on the identical architecture and self-supervised recipe, the input-channel count of the tokenizer being the only difference, and it still trails the proxy model substantially (CRPS 7.52 against 5.54 mm/day, Brier skill -0.04 against $+0.18$).

Model	Params	CRPS (mm/day) ↓	FSS10 ↑	BSS ↑
M-JEPA (proxies)	9.7M	5.54	0.53	0.18
Rain-only control (same recipe)	9.6M	7.52	0.54	−0.04
U-Net (rain only)	351K	7.38	0.14	−0.62
ConvLSTM (rain only)	22K	8.33	0.00	−1.07
ViT (rain only)	836K	8.65	0.42	−0.99

Table S6: Rain-history baselines against the primary model and the same-recipe rain-only control, on the same tracked patches and year-block splits, case-averaged and without recalibration. Parameter counts are the self-supervised backbone for the two JEPA rows (context encoder plus predictor) and the full trained network for the three baselines. All rows are scored on the full test set ($n=337$ windows) except the rain-only control, whose sparse rain-valid data leave it a smaller held-out population ($n=118$), a consequence of the very sparsity the control is designed to expose. The three standard architectures span two orders of magnitude in parameter count (22K–836K) yet all have *negative* Brier skill, below the climatological reference at the 1 mm threshold, so their failure is one of information, not capacity. The rain-only control’s Brier skill is closest to zero among the rain-only models, which supports the main-paper claim that the self-supervised recipe contributes substantively once given adequate information.

Matching the parameter budget while holding the input to rainfall alone therefore does not eliminate the difference. What changes performance is the pretraining *signal* rather than the parameter count, which the control is designed to isolate. This graceful, information-tracking degradation is also the basis for the scaling hypothesis of Section G.

E.2 Randomly Initialized Backbone

A second control isolates *learning* from *architecture*: per-lead R^2 for the trained encoder vs. a randomly initialized backbone (same multiscale transformer + recurrent rollout architecture, no pretraining), in Table S7 and Figure S5a. The trained model separates at every lead; random skill is residual (≈ 0.10 by leads 6–7), reflecting target smoothness and not structure inherent to the untrained architecture. Ratios are computed from the unrounded per-lead values. The trained column comes from a separate execution of the same probe on the same checkpoint and the same 667/337 window split as Table S3; the probe fit is stochastic, so the per-lead values differ from that table by up to 0.037 and the pooled value by 0.003 (0.177 against 0.174). We report each run against its own paired control rather than mixing the two.

E.3 Cross-Target Transfer (OLR)

If the pretrained latent encodes general atmospheric state, its predictive advantage should not be specific to rainfall. Table S8 and Figure S5b give rolled vs. static-latent R^2 for the future-OLR readout at each lead. The margin is non-monotonic: small at lead 1 (+0.03), largest at leads 6–7 ($\approx +0.21$). Absolute skill is low: the rolled probe is positive only at leads 1–2 and negative beyond, and the static probe is negative from lead 2 onward, since forecasting the OLR anomaly that far ahead from a single latent is difficult. The

Lead (days)	Trained R^2 ↑	Random R^2 ↑	Ratio ↑	Gap ↑	Gap 95% CI
1	0.336	0.040	8.4×	+0.296	[+0.183, +0.415]
2	0.189	0.004	42.7×	+0.185	[+0.078, +0.294]
3	0.086	0.018	4.9×	+0.068	[−0.060, +0.200]
4	0.124	0.024	5.3×	+0.101	[−0.018, +0.218]
5	0.132	0.048	2.7×	+0.083	[−0.037, +0.206]
6	0.187	0.102	1.8×	+0.085	[−0.030, +0.205]
7	0.182	0.104	1.8×	+0.078	[−0.039, +0.202]
Pooled	0.177	0.049	3.6×	+0.128	[+0.055, +0.205]

Table S7: Trained vs. random-init backbone R^2 . The trained column is the same rolled-latent predictive-information probe reported in main-paper Section 5.5, re-run alongside its random-init control (see text for the run-to-run spread). Learning multiplies decodability by 5–43× over leads 1–4, converging to $\approx 1.8\times$ at the longest leads as random-feature decodability of a smooth, seasonally autocorrelated target rises; residual random skill is not attributed to the representation. The gap 95% CI is a 10^4 -replicate paired bootstrap over test windows computed under the original pairing; the pooled gap and the near-term leads exclude zero.

quantity of interest is therefore the *relative* advantage of the rollout over the static latent, which is positive at every lead and widens as both probes decay; this is what shows the learned state transfers to a second, held-out atmospheric field.

Lead (days)	Rolled R^2 ↑	Static R^2 ↑	Margin ↑	Margin 95% CI
1	0.454	0.424	+0.030	[−0.058, +0.121]
2	0.163	−0.017	+0.180	[+0.047, +0.322]
3	−0.145	−0.199	+0.053	[−0.124, +0.226]
4	−0.079	−0.170	+0.090	[−0.081, +0.256]
5	−0.125	−0.234	+0.110	[−0.096, +0.298]
6	−0.078	−0.293	+0.215	[−0.001, +0.419]
7	−0.045	−0.249	+0.204	[−0.009, +0.398]
Pooled	0.021	−0.105	+0.126	[+0.016, +0.233]

Table S8: Rolled vs. static-latent R^2 for future-OLR anomaly (cross-target transfer). Margin positive at every lead; absolute skill negative beyond lead 2, but the relative advantage of the rollout persists and widens. The 95% CI is a 10^4 -replicate paired bootstrap over test windows; the pooled margin excludes zero.

The widening margin is consistent with the CKA trajectory’s lead-5 plateau (Section D.3), despite an apparent tension between a representation whose structure saturates by lead 5 and a cross-target margin that continues to widen through leads 6–7. The CKA trajectory is computed once on the latent rollout itself (context latent versus rolled latent at each lead), independent of which downstream target (rain or future OLR) reads it out, so the lead-5 saturation is a property of the latent dynamics, not of the rain decoder. Decomposing the margin into its two arms resolves the apparent tension: the widening does not come from the rolled probe gaining skill (its R^2 stays approximately flat and low across leads 3–7) but from the static-latent control *collapsing*, from $−0.017$ at lead 2 to $−0.293$ at lead 6, as a single frozen snapshot becomes an increasingly poor predictor of the more distant OLR field. The growing margin therefore measures the decay of the persistence baseline, not growth




Figure S5: Controls isolating the source of transfer (main-paper Section 5.6), shown visually. **(a)** Trained encoder vs. randomly initialized backbone of identical architecture: both tested on the same predictive-information probe (rolled-vs-static R^2). The trained model separates from the random one at every lead; random features retain residual long-lead decodability (≈ 0.10 by leads 6–7, indicating target smoothness), while trained skill stays at least $1.4\times$ higher at every lead, confirming the transfer is a product of learning. **(b)** Cross-target transfer: rolled-vs-static R^2 margin for a held-out second target (future OLR anomaly, never seen during rain-decoder training). Margin larger than rainfall (+0.126 vs. +0.090) and broadens with lead (+0.03 near-term, +0.20 long-range), though absolute skill drops negative at long leads. Demonstrates the pretrained representation encodes general atmospheric state, not rain-specific shortcuts.

of rollout information, and is fully consistent with the lead-5 CKA plateau: the rollout saturates while the static control keeps degrading.

E.4 Leave-One-Proxy-Out Ablation

The controls above show that pretraining on the five proxies jointly transfers better than pretraining on rainfall alone, but not that information from one proxy specifically reaches the rainfall readout through the shared latent, as opposed to the five proxies simply supplying more raw pretraining volume. We test this directly: retrain the full pipeline with one background variable (q , u , v , or z) held out of the proxy set, everything else unchanged, and score each ablated model’s probability-branch forecast the same way as the full model (pooled CRPS, BSS, member-FSS10; matched full test population), in Table S9. None of q , u , v , z is itself the readout target, so a drop in rainfall skill when one is removed is evidence that its information was being combined with the others in the latent and used for rain prediction, not merely padding the pretraining sample count.

Every ablation costs CRPS, so every proxy contributes some non-redundant information, but the margins are small next to the 36% CRPS gap between the five-proxy and rain-only models (main-paper Table 3): no single proxy is doing the work of the whole set, consistent with redundant, distributed encoding across the five rather than one dominant channel. Meridional wind is the exception: its removal is the only one that degrades all three metrics together (+1.8% CRPS, -12.8% BSS, -6.1% FSS10), consistent with the humidity–meridional-wind coupling already identified in the latent’s subspace structure (main-paper Section 5.4): the

Proxy set	CRPS ↓	BSS ↑	FSS10 ↑
All five (full model)	5.78	0.18	0.60
– humidity (q)	5.80	0.17	0.58
– zonal wind (u)	5.81	0.17	0.59
– meridional wind (v)	5.88	0.16	0.56
– geopotential (z)	5.81	0.18	0.60

Table S9: Leave-one-proxy-out ablation, pooled convention, full test population. These runs predate the switch to case-averaged scoring and are internally comparable to each other, not to the case-averaged 5.54 of main-paper Table 3. Every removal costs CRPS, but the margins are modest (0.5–1.8%) rather than large. Percentages here and in the text are computed from the unrounded scores, not from the two-decimal entries shown, which round to slightly different ratios. Meridional wind is the one consistently load-bearing variable, the only ablation that also costs BSS and FSS10 together; geopotential’s removal leaves BSS and FSS10 essentially unchanged (-1.0% and $+0.1\%$).

moisture transport that organizes convection is carried jointly by the two, so removing either weakens that signal. Geopotential’s removal is the mildest on BSS and FSS10, which it leaves unchanged, though humidity costs marginally less on CRPS (+0.50% against +0.63%); either way geopotential’s contribution is more redundant with the other four than load-bearing on its own. OLR is not ablated, because it defines the tracking frame: removing it would change which windows exist, confounding an input ablation with a change of evaluation population.

E.5 Tracker and Detection-Setting Sensitivity

A recurring concern with any object-based, tracker-defined evaluation frame is that the reported ranking could be an artifact of the particular detection settings rather than a property of the model (Prein et al. 2024). We test this directly with a sensitivity grid over the two tracker parameters that fix patch geometry, detection quantile $\{8, 10, 12\}\%$ and minimum object area $\{0.75, 1, 1.5\}\times$ the default, re-tracking the held-out test years and re-scoring the frozen model and a dressed-persistence ensemble on each of the nine resulting window populations plus the default control (a configuration-grid sweep with window-bootstrap intervals, following the cell-tracking sensitivity protocol of Meredith, Ulbrich, and Rust (2023)). The model-versus-baseline ranking *direction* is invariant across all ten settings: the model outperforms persistence on CRPS and Brier skill in every configuration and outperforms the deterministic IFS in the same direction on every matched subset, while the wet-cell fraction spans only 0.014 across an order-of-magnitude change in the detection threshold. These comparative conclusions are thus robust to, not contingent on, the tracker choice. The sweep covers the persistence and deterministic-IFS comparisons only; re-running the 51-member ensemble comparison on all ten window populations was beyond the archive access available to us, so the headline probabilistic result of Section F rests on the default configuration alone.

Table S10: Deterministic skill on the full test set ($n=337$), case-averaged over the seven-day lead window (each window/lead scored individually, then averaged; ACC: spatial anomaly correlation; FSS at the 10 mm threshold (Roberts and Lean 2008)), M-JEPA scored with the deterministic branch. *Ensemble mean* averages the probability branch’s 16 raw members into a point forecast, scored identically, to check whether a dedicated point-forecast head earns its keep over simply averaging the generative samples. Boldface marks the best value. Matched-subset comparison against IFS, AIFS, and GraphCast: Table S12; per-lead detail below.

System	RMSE↓	ACC↑	FSS1°	FSS3/5°
M-JEPA	16.24	0.31	0.35	0.47/0.53
Ensemble mean	16.48	0.29	0.34	0.48/0.55
Persistence	23.46	0.18	0.33	0.49/0.56

F Forecast Skill: Extended Results

Main-paper Table 1 gives the full skill matrix (CRPS, BSS, FSS10, Spread/RMSE, rank histogram deviation) for M-JEPA Ens (16 members), IFS Ens (51 members), and the IFS deterministic control, on matched windows ($n=169$, homogeneous-sample rule, fair unbiased CRPS estimator, paired bootstrap with 10^4 resamples); not repeated here. Rank histogram deviation is the mean absolute departure from uniform across bins, lower better. An anomaly-correlation number is not reported for this $n=169$ population, since it was not computed on this matched set; the deterministic anomaly-correlation comparison against IFS, AIFS, and GraphCast on their own larger matched populations is reported separately below and favors the operational references.

F.1 Deterministic Skill by Lead

Dry-area specificity by lead is tabulated in main-paper Table 2, which keeps each reference on its own matched subset and therefore reports two M-JEPA rows; we do not additionally plot it, since collapsing those subsets into a single curve would average over non-homogeneous populations and violate the sampling rule of Section C. Table S11 gives the full per-lead breakdown of the deterministic comparison summarized in the main paper (RMSE, spatial anomaly correlation), for M-JEPA’s deterministic branch and the ensemble-mean point forecast (the probability branch’s 16 raw members averaged, Table S10’s ensemble-mean row) against persistence, case-averaged over the full $n=337$ -window test set for each lead, true patch frame throughout.

The same sharpness trade-off that shapes the probabilistic comparison (main-paper Table 1), where M-JEPA accepts a neighborhood-skill cost relative to the operational ensemble in exchange for a CRPS advantage, appears again here against persistence: the deterministic branch is uniformly better on point accuracy and pattern correlation, at 7–8 mm/day lower RMSE and roughly double the spatial ACC from lead 2 onward, while persistence retains a small advantage at the 5° neighborhood at lead 1 and from lead 4 onward. A field that relays the last observed rain distribution scores well on

Lead (days)	M-JEPA (det)	Ensemble mean	Persistence
<i>RMSE (mm/day) ↓</i>			
1	16.79	17.20	21.12
2	17.13	17.46	23.97
3	16.88	17.12	24.30
4	16.39	16.62	23.96
5	15.99	16.13	23.73
6	15.51	15.68	23.69
7	14.97	15.13	23.45
<i>Spatial ACC ↑</i>			
1	0.447	0.409	0.384
2	0.364	0.328	0.201
3	0.313	0.291	0.148
4	0.279	0.259	0.139
5	0.266	0.256	0.129
6	0.273	0.257	0.122
7	0.261	0.244	0.114
<i>FSS10 at 5° neighborhood ↑</i>			
1	0.721	0.726	0.756
2	0.633	0.641	0.606
3	0.560	0.573	0.548
4	0.493	0.513	0.536
5	0.451	0.486	0.513
6	0.438	0.476	0.493
7	0.414	0.450	0.473

Table S11: Per-lead breakdown of Table S10’s comparison against persistence, same convention and routines throughout (10 mm threshold, 5° neighborhood). Persistence is rebuilt here as the last observed context field at the fixed anchor; the lead-averaged values reproduce Table S10’s upper block exactly.

coarse-scale overlap and poorly on placement and amplitude, which is what the two metric families report here. Persistence being uncompetitive on RMSE and ACC at every lead confirms the rollout is not simply relaying the initial condition forward.

The matched-coverage comparison against IFS, AIFS, and GraphCast summarized in main-paper Section 5.2 (Table S12 below) restricts M-JEPA to the cells its tracked patches cover: the same patch-native frame as main-paper Table 1, not the full 40×40 domain the references forecast natively. Each system is compared on its own matched-coverage subset ($n = 266, 127, 127$; the all-five intersection is only ≈ 14 days, so the larger pairwise subsets are used and disclosed). Each lead is scored independently over every valid tracked window reaching that (calendar day, lead) pair, standard object-verification practice (e.g. tropical-cyclone track verification, MODE), not a requirement that one single tracked window supply all seven leads.

F.2 The Archives’ Rising-ACC Pattern: Mechanism and a Full-Domain Check

Table S13 shows AIFS and GraphCast’s own ACC *rising* and RMSE *falling* with lead (AIFS: ACC 0.376 $\rightarrow$ 0.458, RMSE 21.48 $\rightarrow$ 16.63; GraphCast: ACC 0.379 $\rightarrow$ 0.457, RMSE 20.86 $\rightarrow$ 16.46), the physically backwards direction

Table S12: Deterministic skill, case-averaged over the seven-day lead window, M-JEPA re-scored on each reference’s own matched-coverage subset (IFS: $n=266$; AIFS and GraphCast: $n=127$, an identical subset by construction, reported once). *Ensemble mean* as in Table S10. Every entry is the mean of the corresponding per-lead column of Table S13, each lead carrying equal window count, so the two tables reconcile term by term. The anomaly correlation uses a per-lead climatology throughout, as elsewhere in this paper; a climatology pooled across leads raises every ACC in this table by 0.005–0.010 without changing any ordering.

System	RMSE ↓	ACC ↑	FSS1°	FSS3/5°
<i>IFS matched subset ($n=266$); M-JEPA re-scored</i>				
M-JEPA	18.23	0.283	0.34	0.46/0.52
Ensemble mean	18.50	0.255	0.33	0.47/0.54
IFS	18.45	0.328	0.44	0.60/0.67
<i>AIFS/GraphCast matched subset ($n=127$); M-JEPA re-scored</i>				
M-JEPA	19.74	0.339	0.37	0.49/0.56
Ensemble mean	20.14	0.294	0.35	0.49/0.56
AIFS	18.92	0.419	0.49	0.63/0.69
GraphCast	18.81	0.418	0.46	0.60/0.65

for a forecast skill curve.

Mechanism. Because ACC is scale-invariant (Section C), a forecast that loses fine-scale detail relative to the truth as lead grows can show *rising* ACC even as real skill falls: the residual correlation concentrates on the retained large-scale pattern once high-frequency, largely uncorrelated detail is filtered out. We test this with two per-lead diagnostics on the same patch population: a damping ratio ($\text{std}(\text{pred})/\text{std}(\text{obs})$) and a roughness ratio (mean absolute nearest-neighbor gradient of forecast over obs, a blur signature amplitude alone can miss); Figure S6 shows the damping ratio by lead for all four systems. AIFS’s damping ratio falls $0.549 \rightarrow 0.488$ and its roughness ratio falls $0.396 \rightarrow 0.323$ over the horizon: a genuine, progressive self-smoothing that mechanically raises ACC under a scale-invariant metric. GraphCast’s roughness ratio is flat over the same horizon (≈ 0.27 – 0.29): it is uniformly over-smooth from lead 1 rather than increasingly so, while the observed field’s own roughness falls 22% ($9.41 \rightarrow 7.35$) over the horizon, matching the event-lifecycle decay already established in main-paper Section 5.3: as the tracked system ages, the truth itself simplifies toward something a persistently smooth forecast pattern-matches more easily, without GraphCast itself changing.

Case-level test of the mechanism. The lead-aggregated numbers above are only 7 points per archive and their ACC does not fall monotonically lead-to-lead, so we test the causal claim directly and with far more statistical power: every individual (window, lead) case’s ACC scattered against that same case’s own observed-field roughness, pooling across all leads (Figure S7). The relationship is negative and substantial at case-level resolution ($n=462$ – 1127 per archive): Pearson $r = -0.28$ (IFS), -0.57 (AIFS), -0.57 (GraphCast). Rougher, more complex target cases achieve lower




Figure S6: Amplitude-stability comparison among actual forecasting systems (persistence excluded, it relays a stale, systematically too-large field and runs to 1.31 by lead 7, off this axis): damping ratio $\text{std}(\text{pred})/\text{std}(\text{obs})$ by lead, patch frame, case-averaged. IFS retains the most amplitude (≈ 0.55 – 0.60 , closest to 1), M-JEPA and AIFS track a comparable, non-collapsing band (≈ 0.44 – 0.55), and GraphCast sits lowest throughout (≈ 0.41 – 0.46), consistent with its already-smooth-from-lead-1 signature. M-JEPA shows no distinct advantage on this diagnostic; the point is that it does not exploit the amplitude-collapse route to inflated ACC that AIFS’s declining ratio suggests.




Figure S7: The ACC-mechanism, tested directly: every (window, lead) case’s ACC vs. that case’s own observed-field roughness, pooled over leads, with fitted trend and Pearson r per archive.

ACC regardless of which lead they come from, and smoother, simpler cases achieve higher ACC regardless of lead; lead enters only because it correlates with case roughness (the lifecycle-decay effect), not because ACC directly rewards longer lead.

Full-domain check. Both patch-frame mechanisms above are consistent with the trend being a property of the *tracked, aging system* the patch follows, not of the archives’ skill at long lead in general. We test this directly: the same forecast files and target days, rescored on the full 40×40 domain

the reference systems natively forecast, instead of the patch crop around the track (Table S14). The rising-ACC / falling-RMSE pattern disappears entirely— ACC is flat and near zero for all three systems across the full horizon, and RMSE mildly *increases* with lead, the physically expected direction. Full-domain ACC is also far lower in absolute terms than the patch-frame values ($\lesssim 0.04$ vs. $0.29\text{--}0.46$), as expected: the patch frame’s climatology reference captures structure specific to an active convective envelope, while the full domain dilutes that signal with mostly quiescent surrounding area. The disappearance of the trend rules out a genuine improvement in archive skill at longer lead; it is consistent with an artifact of scoring one aging system’s patch through its life, the same effect underlying the observed amplitude decay reported in main-paper Section 5.3. Age and lead covary in this design and are not separately identified.

F.3 Ensemble Mean by Lead

Table S15 gives the per-lead detail for Table S10’s ensemble-mean row (the probability branch’s 16 raw members averaged into a point forecast), same conventions and populations as Tables S11–S13, true patch frame throughout.

The ensemble mean’s cross-model detail (vs. IFS/AIFS/GraphCast) is merged into Table S13 above as the “ens.” columns alongside the deterministic head.

F.4 Extreme-Event Skill of the Probability Branch

Grid-point categorical scores (POD: probability of detection; FAR: false alarm ratio; CSI: critical success index; ETS: equitable threat score) are strict single-cell measures with no neighborhood tolerance, so they isolate exact-hit skill from the placement credit FSS gives nearby cells. On the matched $n=169$ population in the true patch frame, member-averaged POD at Heavy is 0.053 against the operational ensemble’s 0.041, and at Very Heavy 0.023 against 0.015; the ordering reverses only at Extremely Heavy (0.006 vs. 0.013, 230 matched fields, the thinnest sample of the three tiers).

Member-FSS at that third tier behaves the same way and is the one threshold where the main paper’s extreme-skill result does not hold. The ordering is lead-dependent: IFS Ens leads at the first two leads (0.069 against 0.196 at lead 1; 0.062 against 0.144 at lead 2, M-JEPA first in each pair), M-JEPA Ens leads from lead 3 onward and by up to $6.6\times$ at lead 5 (0.046 against 0.007), and pooled over leads the short-lead deficit dominates (0.056 against 0.077, 230 matched fields). With 27–38 qualifying events per lead this tier is too thin to support either direction.

F.5 Directional Tracking and Rollout Velocity of the Decoded Forecast

Figure S8 shows one decoded forecast against observations across the horizon, as a qualitative reference for the pooled diagnostics that follow. Grid-point and neighborhood scores measure where rain is placed, not whether the forecast follows the *motion* of the convective envelope. To test motion at the decoded-forecast level, we take the precip-weighted centroid (cells > 1 mm/day) of the observed and forecast




Figure S8: GPM observations (left) and the M-JEPA deterministic field (right) at days 1, 3, 5, 7 of the seven-day forecast for the wettest test window by observed day-3 rainfall (target days 2023-07-02 to 2023-07-08). The forecast places a maximum on the observed coastal cell at short range and broadens into a smoother band by day 7. Dashed box marks the native patch extent.

rain field at each lead, for every held-out window ($n=337$), and evaluate a directional rather than a magnitude criterion: whether the forecast centroid’s drift relative to lead 1 has the same *sign* (northward vs. southward) as the observed drift, regardless of how far it moves (Figure S9a). At the loosest threshold (any nonzero observed drift), directional accuracy is 58–65% across leads, against a 50% chance rate ($n=2022$ pooled over leads 2–7, binomial $p < 10^{-4}$); restricting to observed drifts $> 0.5^\circ$ raises this to 61–69%, and $> 1.0^\circ$ to 63–71% ($n=1355$ pooled): the signal strengthens, not weakens, as near-zero-displacement noise is excluded, the signature of a genuine effect rather than an artifact of small splits. The same pattern holds on the zonal (east/west) axis: pooled accuracy 59.0% ($n=2022$, $p < 10^{-4}$), rising to 64.0% restricted to drifts $> 1.0^\circ$. This is a directional result only, and says nothing about displacement *magnitude*, which the model does not resolve well (see below).

Section D.3’s CKA plateau is a representational-structure diagnostic and does not by itself establish whether the latent is still *moving* once CKA saturates. Figure S9(b) answers

that directly, with the raw step-to-step displacement of the rollout in latent space, $\|\hat{z}_d - \hat{z}_{d-1}\|_2$ (pooled foreground latent, same pooling as the linear-probe features), at every autoregressive step. Displacement falls from 2.25 (context $\rightarrow \hat{z}_1$) to 1.70 (step into lead 2) and continues decaying toward a non-zero floor of 1.33–1.38 over leads 5–7, where the bootstrap CIs overlap and the curve is flat and no longer shrinking. The ratio of the first step to the last (context $\rightarrow \hat{z}_1$ against lead 6 $\rightarrow 7$) is $\approx 1.69\times$: rollout velocity falls to about 60% of its initial rate but does not vanish. This refines the rollout-saturation picture and does not contradict it: the CKA plateau shows the representation’s *structure* stops changing, while panel (b) shows the underlying latent state itself keeps moving at a damped, non-zero rate, so saturation asymptotes to a nonzero floor rather than collapsing the rollout to a fixed point.

Panel (c) tests amplitude over the event’s own lifecycle, which is the axis on which the forecast does track the observed evolution. For every held-out window we compute the day since the track’s genesis for each of its seven forecast-lead target days, and pool the observed and decoded area-mean rain intensity by that lifecycle day across the 49 distinct tracks that supply the 337 evaluation windows (a subset of the 149 detected test-year tracks in Figure S2: a track enters the evaluation set only where it yields a valid context-plus-horizon window at an unmasked anchor, and each contributing track supplies several overlapping windows). Context length fixes a floor of five days before any target day is reachable, so the earliest lifecycle day observable by this construction is day 5, a coverage limit of the windowing, not a selection choice. From day 5 through day 16 the decoded curve’s *shape* tracks the observed decay closely: each series normalized to its own day-5 value falls to $0.71\times$ (observed) and $0.71\times$ (forecast) by day 16, decay rates of -0.0247 and -0.0253 per day, on a dry bias of 2.1 mm/day over those days. Beyond day 16, in the thinner-sampled tail of longest-lived tracks ($n=84$ falling to $n=42$ by day 20), the observed curve continues falling to $0.68\times$ while the decoded curve stops declining at $0.75\times$: the forecast fails to continue the observed decay for the longest-lived systems, which is the same damping seen in the CKA plateau and the latent displacement floor, expressed in decoded amplitude at increasing lifecycle duration. Because the day 5–16 and post-day-16 ranges behave differently, both are reported; the split is not a sampling threshold, since day 17 ($n=84$) is better sampled than day 5 ($n=40$).

F.6 Radial Power Spectra of the Ensemble Members

Main-paper Figure 2b compares the spatial power spectrum of each ensemble’s members against observations, following the azimuthally-averaged spectral-realism check used for generative weather models (Price et al. 2025). Because a Lagrangian patch is a 20×20 window rather than a global field, we use a 2-D radial spectrum in place of a spherical-harmonic one. Each field is mean-subtracted and multiplied by a 2-D Hann window to suppress edge leakage, transformed with fft_2 , and its squared magnitude binned by integer radius $r = \lfloor \sqrt{(i - c_y)^2 + (j - c_x)^2} \rfloor$ about the shifted spectral

center; the $r=0$ (DC) bin is discarded, leaving radii 1–9. Each system’s curve is the ratio of its mean binned power to the observed mean binned power in the same bin, so a value of 1 means the forecast carries exactly the observed variance at that scale.

Averaging runs over every member, every lead and every window of the matched population: 1183 observed fields (169 windows $\times$ 7 leads), 18 928 M-JEPA member fields ($\times 16$) and 60 333 IFS Ens member fields ($\times 51$). Both systems lose power at every radius, so the informative quantity is the *direction* of the loss with wavenumber, not its level. IFS Ens declines monotonically (0.484 at $r=1$ to 0.170 at $r=9$), the expected signature of a mass-conserving smoother, which spreads a sharp feature over a wider area at lower amplitude. M-JEPA Ens begins slightly lower (0.464) and rises to 0.722 at $r=8$ and 0.711 at $r=9$, $4.2\times$ IFS Ens’s retention there. This is a descriptive comparison over pooled fields, not a hypothesis test, and we attach no confidence interval to the ratios. It also speaks only to the amount of small-scale variance, not to its placement, which is why the main text reads it together with the member-FSS result.

G Scaling Hypothesis: Information-Limited Signal

The rain-only control’s pretraining window count is 226 against 3885 for the all-season five-proxy pretraining signal, a $17\times$ reduction in information. Because the control is trained on rainfall alone, this same rain-valid window population serves both its pretraining and its decoder training, whereas M-JEPA pretrains on the 3885 all-season proxy windows and trains its decoder on the 667 rain-labeled subset. Despite this, the control’s skill degrades gracefully instead of collapsing (CRPS 7.52 against 5.54 mm/day, with Brier skill closest to zero among the rain-only models; main-paper Table 3). A collapse-scale degradation would point to an idea-limited method, in which the architecture or loss needs the specific target signal to work at all; the graceful degradation observed instead is consistent with an information-limited method, where performance tracks the richness of the pretraining signal rather than a narrow recipe requirement. We advance this as a hypothesis supported by the control comparison, not as a measured scaling law across multiple window-count settings; establishing a full scaling curve (pretraining skill as a function of proxy window count, proxy channel count, and temporal resolution) is the natural next step.

Temporal resolution is the other information axis. All three deterministic references that outperform M-JEPA on point-forecast metrics (main-paper Section 5.2) resolve atmospheric state well below the daily cadence of our pipeline: IFS’s dynamical core integrates continuously, and both AIFS and GraphCast step autoregressively at 6-hourly intervals (Lang et al. 2024; Lam et al. 2023). M-JEPA’s five proxies, by contrast, are resampled to daily means before the encoder ever sees them, and the rollout itself advances one calendar day per autoregressive step; any sub-daily structure in the underlying fields, including the diurnal cycle of convective initiation and the nocturnal propagation of mesoscale convec-




Figure S9: Pooled evolution diagnostics over the full test set ($n=337$), rain-centroid based (1 cell = 1° ; 95% window-bootstrap bands where applicable). **(a)** Northward/southward directional accuracy of the decoded centroid’s drift relative to lead 1 (sign match against observed), at three observed-drift thresholds: above the 50% chance line at every lead and every threshold, and rising as the threshold tightens (near-zero-displacement noise excluded). **(b)** Step-to-step latent rollout displacement $\|\hat{z}_d - \hat{z}_{d-1}\|_2$ against lead (pooled foreground latent; the open marker and dashed segment mark the context $\rightarrow \hat{z}_1$ step, a different regime from the subsequent $\hat{z}_{d-1} \rightarrow \hat{z}_d$ steps). Displacement decays from 2.25 to a non-zero floor of 1.33–1.38 by leads 5–7 ($\approx 1.69 \times$ damping, early against late step): the latent rollout is damped but not frozen. **(c)** Composite intensity lifecycle: observed and decoded area-mean rain by day since genesis, pooled over 49 test tracks (window-bootstrap bands; days with fewer than 20 pooled samples omitted). Normalized to each series’ own day-5 value, the decay rates over days 5–16 are -0.0247 and -0.0253 per day on a dry bias of 2.1 mm/day; in the thinner tail of the longest-lived tracks (day 17 onward) the decoded curve stops declining while the observed curve continues to fall. Panels (b) and (c) are two signatures of the same rollout damping, on undecoded latent velocity and decoded amplitude respectively; panel (a) is a directional result and does not speak to damping.

tive systems characteristic of the Bay of Bengal, is discarded at the data-ingestion stage, not merely under-modeled. This is consistent with denser temporal sampling being one of the information-ceiling levers named above, though the three references also differ from M-JEPA in initial-condition quality and training-corpus scale, so temporal resolution cannot be isolated as the cause from this comparison alone; a controlled ablation at matched initialization and data volume, varying only proxy cadence, is the direct test.

H Computational Cost and Runtime

Model size: encoder and autoregressive predictor 9.7M parameters combined (4.8M and 4.9M respectively), decoder 5.7M parameters (frozen Stage-1 stack, trainable decoder heads only at readout time), for the 15.4M total quoted in the main text. Inference cost splits by head. The deterministic point-forecast head is a single forward pass: a full seven-day forecast for a batch of 16 patches runs in 204 ms on a single GPU (12.8 ms per patch, amortized), with no iterative sampling. The ensemble-generating flow probability head used for the matched IFS-Ens comparison (“Forecast Skill” section, main paper) instead requires 30 rectified-flow integration steps per ensemble member, and is substantially more expensive per forecast than the deterministic head, though it still runs on the same single GPU; members are scored as generated, with no post-hoc recalibration step. No compara-

ble figure exists for the operational ensemble in a form we can measure, so the main text makes no quantitative compute comparison against IFS. We did not use the flow head for the probing suite (Section D), which reads out from the frozen encoder alone and does not invoke either decoder head.

Pretraining hardware and software. All training and inference for this work ran on a single NVIDIA RTX A4000 (16 GB) under Ubuntu 22.04.5 LTS, with Python 3.13.1, PyTorch 2.6.0 (CUDA 12.4, cuDNN 9.1.0), NumPy 2.2.2, SciPy 1.15.1, scikit-learn 1.6.1, xarray 2025.1.2, and pandas 2.3.3. Every run seeds Python, NumPy, and PyTorch (CPU and CUDA) from a single configured value (default 1410) and disables cuDNN autotuning in favor of deterministic kernels; the seed-sensitivity figures reported above use seeds 1410, 42, and 7. The self-supervised encoder pretraining recipe was: VICReg anti-collapse objective, auxiliary OLR-reconstruction weight 0.10, batch size 32, and 100 epochs; Stage 1 uses no spatial jitter, which is applied only when training the Stage-2 decoder (± 2 grid cells, $\approx \pm 2^\circ$). The VICReg coefficients were variance 1.0, covariance 0.04, and $\varepsilon = 10^{-4}$. Optimization used AdamW with learning rate 5×10^{-4} , weight decay 0.05, and a 10-epoch warmup.

Lead	RMSE ↓			ACC ↑			FSS ↑		
	det.	ens.	ref	det.	ens.	ref	det.	ens.	ref
<i>vs. IFS (n=161)</i>									
1	19.03	19.49	20.18	0.410	0.366	0.367	0.725	0.723	0.759
2	19.27	19.64	19.49	0.326	0.284	0.355	0.624	0.636	0.717
3	18.97	19.19	18.98	0.282	0.259	0.349	0.535	0.555	0.695
4	18.41	18.72	18.57	0.248	0.220	0.319	0.481	0.492	0.666
5	17.87	18.04	18.07	0.234	0.222	0.305	0.447	0.485	0.640
6	17.22	17.41	17.26	0.240	0.219	0.292	0.435	0.477	0.609
7	16.86	17.02	16.62	0.238	0.217	0.308	0.410	0.434	0.593
<i>vs. AIFS (n=66)</i>									
1	21.02	21.61	21.48	0.376	0.319	0.376	0.726	0.719	0.755
2	21.07	21.66	20.30	0.343	0.287	0.414	0.650	0.648	0.734
3	20.58	21.10	19.55	0.357	0.305	0.430	0.587	0.594	0.718
4	19.84	20.21	19.01	0.349	0.305	0.422	0.544	0.542	0.689
5	19.11	19.25	18.35	0.332	0.308	0.409	0.503	0.529	0.679
6	18.25	18.61	17.14	0.321	0.276	0.425	0.460	0.480	0.635
7	18.28	18.52	16.63	0.294	0.261	0.458	0.421	0.418	0.618
<i>vs. GraphCast (n=66)</i>									
1	21.02	21.61	20.86	0.376	0.319	0.379	0.726	0.719	0.729
2	21.07	21.66	20.24	0.343	0.287	0.416	0.650	0.648	0.694
3	20.58	21.10	19.78	0.357	0.305	0.416	0.587	0.594	0.682
4	19.84	20.21	19.14	0.349	0.305	0.412	0.544	0.542	0.656
5	19.11	19.25	18.25	0.332	0.308	0.409	0.503	0.529	0.648
6	18.25	18.61	16.95	0.321	0.276	0.435	0.460	0.480	0.602
7	18.28	18.52	16.46	0.294	0.261	0.457	0.421	0.418	0.571

Table S13: Cross-model skill by lead (per-lead detail behind Table S12 above), restricted to the cells M-JEPA’s tracked patches cover, the same patch-native frame as main-paper Table 1 and main-paper Figure 3. RMSE (mm/day) ↓, ACC ↑, FSS ↑ (10 mm, 5°), case-averaged. “det.” is the deterministic head; “ens.” is the probability branch’s 16 raw members averaged into a point forecast (Table S10’s ensemble-mean row). Window count per lead: 161 (IFS), 66 (AIFS/GraphCast); pooled across leads these give the 266/127 distinct target days of Table S12. All three references lead on ACC and FSS at every lead. On RMSE, the deterministic head is lower than IFS at leads 1–6 (IFS regains it only at lead 7) but never leads AIFS or GraphCast; the ensemble mean’s RMSE advantage over IFS is narrower still (leads 1 and 5 only) and never leads AIFS/GraphCast.

Table S14: Full-domain (40×40) rescoring of the reference archives, case-averaged, on the Table S13 target days for which a complete full-domain reference field exists; that requirement drops roughly a tenth of the windows (145/63/63 against 161/66/66), since a patch-frame score needs only the patch cells whereas a full-domain score needs the whole grid. RMSE (mm/day) $\downarrow$, spatial anomaly correlation $\uparrow$. Unlike the patch-frame comparison, no patch is centered on the tracked system here, so any lifecycle-decay effect specific to the tracked system cannot appear.

Lead	IFS ($n=145$)		AIFS ($n=63$)		GraphCast ($n=63$)	
	RMSE	ACC	RMSE	ACC	RMSE	ACC
1	19.73	-0.020	20.52	0.020	19.24	0.032
2	19.78	-0.017	20.14	0.029	19.19	0.040
3	20.03	-0.026	19.96	0.025	19.19	0.035
4	20.26	-0.035	19.95	0.007	19.36	0.023
5	20.34	-0.030	20.21	0.003	19.89	0.017
6	20.69	-0.023	20.35	0.010	20.26	0.022
7	20.74	-0.020	20.32	0.013	20.34	0.023

Lead (days)	Ensemble mean	Persistence
<i>RMSE (mm/day) $\downarrow$</i>		
1	17.20	21.12
2	17.46	23.97
3	17.12	24.30
4	16.62	23.96
5	16.13	23.73
6	15.68	23.69
7	15.13	23.45
<i>Spatial ACC $\uparrow$</i>		
1	0.409	0.384
2	0.328	0.201
3	0.291	0.148
4	0.259	0.139
5	0.256	0.129
6	0.257	0.122
7	0.244	0.114
<i>FSS10 at 5° neighborhood $\uparrow$</i>		
1	0.726	0.756
2	0.641	0.606
3	0.573	0.548
4	0.513	0.536
5	0.486	0.513
6	0.476	0.493
7	0.450	0.473

Table S15: Ensemble mean vs. persistence by lead, full test set ($n=337$), case-averaged. Unlike the pooled convention, the ensemble mean's ACC now exceeds persistence's at every lead, including lead 1. FSS10 favors the ensemble mean at leads 2–3 only and persistence at leads 1 and 4–7, the same crossover pattern as the deterministic branch (Table S11).