



Prof. Dr.-Ing. A. Bruhn
Institute for Visualization and Interactive Systems
Department Intelligent Systems
University of Stuttgart

Homework Assignment 3

H 3.1 (Variational Methods)

10 Points

Instead of using the grey value constancy assumption, let us assume that the y -derivative of the image f remains constant over time, i.e.

$$f_y(x + u, y + v, t + 1) = f_y(x, y, t) \quad (1)$$

- (a) Linearise the constancy assumption (1) w.r.t the flow functions u and v .
 - (b) Write down an energy functional similar to Horn and Schunck based on the linearised constancy assumption from (a).
 - (c) Compute the Euler-Lagrange equations for this energy functional.
 - (d) Discretise the Euler-Lagrange equations from (c).
 - (e) Starting from the discrete equations computed in (d), derive an iterative scheme that computes the minimiser.
-

H 3.2 (Stereo)

6 Points

Consider a camera with focal distance 2. Its image coordinate system is orthogonal with square pixels of size 1. The principal point in this coordinate system is located in $(2, 3)^\top$. Furthermore, the position of the world coordinate system relative to the camera coordinate system is given by a rotation around the z -axis by an angle of 90° and a translation by the vector $(5, 0, -1)^\top$.

- (a) Compute the intrinsic matrix A_{int} (including the focal length f).
 - (b) Compute the extrinsic matrix A_{ext} .
 - (c) Compute the full projection matrix P .
-

P 3.3 (Horn and Schunck)

Please download the required file `cv19_ex03.tgz` from ILIAS. To unpack the data, use `tar xvfz cv19_ex03.tgz`.

In the routine `flow`, supplement the missing code such that it computes one Jacobi iteration step. Make sure the image boundaries are treated correctly. To compile the program, type

```
gcc -O3 -o hsTemplate hsTemplate.c -lm
```

The filling-in effect that is characteristic for variational methods can be studied with the image pair `pig1.pgm`, `pig2.pgm`. To this end, investigate the result for different numbers of iterations. What is a good value for the regularisation parameter α in this case?

Submission:

The theoretical problem(s) have to be submitted in handwritten form before the next tutorial (December 06th).

Deadline for Submission is: Friday, December 06th, 9:45 am (before the tutorial)



Prof. Dr.-Ing. A. Bruhn
Institute for Visualization and Interactive Systems
Department Intelligent Systems
University of Stuttgart

Classroom Assignment 3

C 3.1 (Eigenvalue Analysis)

Let $\mathbf{J} \in \mathbb{R}^{n \times n}$ be a symmetric $(n \times n)$ matrix with real components. We consider its corresponding quadratic form given by

$$E : \mathbb{R}^n \longrightarrow \mathbb{R}, \quad E(\mathbf{v}) = \mathbf{v}^\top \mathbf{J} \mathbf{v}$$

Show that among all vectors $\mathbf{v} \in \mathbb{R}^n$ with $|\mathbf{v}| = 1$, the function value $E(\mathbf{v})$ is minimal for the eigenvector of \mathbf{J} corresponding to its smallest eigenvalue. What can we say about E if \mathbf{J} is positive definite?