POLICE LABORATORY REPORT ON SEDIMENT SAMPLES OBTAINED FROM THE VICTIM

- the beach where the body was found [Sample V1]
- the inside of the shattered skull and matted hair and a pocket in the victim's clothing [Sample V2]
- a filthy handkerchief that was found at the scene. Nasal mucus on the handkerchief contained mica, hornblende, quartz and feldspar.
- the victim's watch, which had stopped at 1735.
- the campsite in the dunes where the victim and suspects were staying [Sample V3]
-

Sample Descriptions:

	Sample V1	Sample V2	Sample V3
Grain shape	Mineral grains: spherical to subspherical Shelly material: Variable – plates, rods, subspherical Rock (lithic) grains: subspherical , platey	Mineral grains: subspherical Shelly material: : platey Rock (lithic) grains: subspherical platey	Mineral grains: spherical
Angularity	Shelly material: rounded-well rounded Mineral grains - Quartz: subangular-subrounded	Mineral grains: Sub rounded to subangular Rock: v.angular-subangular Shelly material: rounded to angular	Mineral grains: 90% rounded-well rounded 10% subangular (tend to be the smaller grains
Composition	50% quartz 48% shelly material incl. in order of relative abundance: shell fragments (bivalves and gastropods), whole gastropods (3 types), disarticulated bivalves, forams (2 types), bryozoa (2 types), echinoid spines (green and purple), and fish teeth. 2% basic igneous (including basalt) and metamorphosed basic intrusive igneous	40% shell fragments 10% coralline algae 40% quartz 10% basic igneous (including gabbro)	95% quartz 5% calcite (occur at fine end of grain size distribution)
Other observations	Polished shell fragments- high degree of abrasion		Grain surface is frosted, Fe oxide staining present