



**The comparison of fitness of
mechanistic vs. phenomenological
microbial population growth models
based on an empirical dataset**

MSc Computational Methods in Ecology and Evolution

Department of life Science

Imperial College London

Yuan Zhang

yuan.zhang119@imperial.ac.uk

Word count: 3055

Contents

1	Introduction	2
2	Methods	5
2.1	Data and preparation before the modelling	5
2.2	Models	5
2.2.1	Mechanistic Model	5
2.2.2	Phenomenological Model	6
2.3	Model fitting and Visualization	7
2.4	Model selection criteria	7
2.5	Computing languages	7
3	Results	8
3.1	Fitting percentage of 4 models	8
3.2	Best fitting model assessment	8
3.3	Four model fittings on one species with different temperatures	9
4	Discussion	10
4.1	Difference between phenomenological and mechanistic models	10
4.2	Comparison of 2 phenomenological models (Gompertz model and Baranyi model)	11
4.3	Temperature factor	12
4.4	Limitation and suggestions for future studies	12
5	Conclusion	13

1 Abstract

2 There is an essential function of the fluctuations in single population abundance for the
3 ecosystem dynamic and functional characteristics. In this experiment, focusing on 4 mod-
4 els (Logistic model, Gompertz model, Baranyi model, as well as the Buchanan model),
5 fit probability and the probability of becoming the best fit model were tested based on
6 275 data sets. As a result, it was found that the Logistic model has a large probability
7 of fit but 2 phenomenological models (Gompertz model, Baranyi model) have better fit-
8 ness. The Buchanan model shows very low levels in both the probability of fit and the
9 probability of best fit. Comparisons between the models are discussed. Moreover, tem-
10 perature as an environmental factor has also been studied and discussed. Finally, based
11 on this experiment and other researches, ideas and suggestions for future modeling were
12 proposed.

13 1 Introduction

14 As an indispensable part of population research, microbial growth has been extensively
15 studied. There is an essential function of the fluctuations in single population abundance
16 for the ecosystem dynamic and functional characteristics. Additionally, population growth
17 was described as a reliable index to reflect biological responses (Pörtner et al. 2010).
18 Some issues could be investigated and predicted by focusing on microbial growth. No-
19 tably, in terms of risks of food poisoning, caused by food spoilage, pathogen growth will
20 consequently raise a common issue of public health risk (Peleg & Corradini 2011). Be-
21 sides, tumor growth pasterns also play a crucial role in the ongoing battle against cancer
22 (Araujo & McElwain 2004, d'Onofrio 2009). A miscellany of new strategies, experimental
23 techniques and theoretical approaches can be integrated into the struggle against the
24 cancer. However, to explain and interpret the experimental discoveries, researchers usu-
25 ally make the best use of recourse to mathematical models based on previous cancer re-
26 searches in many aspects and fields. Therefore, scientists has become conscious of the
27 possibilities of mathematical models, realizing that the existing medical techniques and
28 experimental methods can not be able to tell the differences between a variety of possible

mechanisms underlying key aspects of tumour evolution(Araujo & McElwain 2004).

The population is usually growing in an exponential way at the beginning, based on the low abundance and unlimited resources on the basis of the Malthusian principle (Costabile & Rowthorn. 1985). Before the population grows fast, there exists a time lag. As time goes by, the population is growing more and more slowly and finally at a relatively stable balanced point because of limitations in several aspects. It is essential to pay much attention to microbial growth rates. Particularly, the latter one in batch culture usually has a specific series of phases. The first one is named as the lag phase, in the course of which a series of transcriptional machinery can be stimulated, such as the genes referring to the nutrient uptake and metabolic change for the preparations for bacteria growth. The second phase is called the exponential growth phase, in which the number of bacteria at a steady speed and the amount will double with every generation. The growing speed will not slow down until it reaches the carrying capacity of the media. Then it will step into a stable status for the population, entering the followed phase called the stationary phase.

In previous studies, one of the measurements to determine the microbial growth rate is widely used (Peleg & Corradini 2011, Zwietering et al. 1990). As Figure1 presents, explicitly, it plots the cell number or density against time on a semi-log graph and to fit a straight line through the exponential growth phase – the slope of the line shows the highest growth rate. It is also practical to develop the models to depict the entire sigmoidal bacterial growth curve. Bacterial growth usually reflects a phase where the specific growth rate begins at the point of zero. Next, it will speed up to a high-level value within one particular period of time, leading to a lag time(t_{lag}). Moreover, the growth curve will cover a final phase where the rate comes down and eventually to zero; that is, it will reach an asymptote (N_{max}) (Halimi & Shukor 2014). In Figure 1, the growth curve can be defined as the logarithm of the number of organisms plotted against time, and the results vary to the changes of growth rates in a sigmoidal curve, having a lag phase merely behind $t=0$ followed by an exponential phase and next by a stationary phase. In addition to the lag period and the asymptotic value, there is some other valuable parameter of the growth curve, meaning the maximum specific growth rate(r_{max}). The slope of the line determines r_{max} on the condition that the organisms have an exponential growth since




Figure 1: A growth curve of semi-log density against time (this is a modified version from Zwietering et al. (1990))

59 the logarithm of the number.

60 Mathematical models have been applied and complicated with various parameters in
 61 bacterial growth studies, while different models have different results to some degree.
 62 Most models can be classified into two types: mechanistic and phenomenological ones
 63 and applications (Peleg & Corradini 2011). Given that a mechanistic model provides a
 64 tentative kinetic explanation of growth, it can be extensively applied in a large class of
 65 cases (Peleg et al. 2007). Particularly, a classical mechanistic model is called the logistic
 66 (Verhulst) equation, which has various modifications in former researches (Fujikawa et al.
 67 2004, Kacena et al. 1999, Peleg et al. 2007). Buchanan model (Buchanan et al. 1997)
 68 is also mechanistic as it could be considered as a 'three-phase logistic mode'. Different
 69 from the mechanistic ones, phenomenological models often are not derived from first
 70 principles, describing the empirical connection of phenomena. Several phenomenological
 71 growing models can be referred from literature, including the modified Gompertz model
 72 (Zwietering et al. 1990) and Baranyi model (Baranyi 1993, Grijspeerd & Vanrolleghem
 73 1999). To explain the curve and reduce the measured samples to a limited number of
 74 valuable parameters, the adequate models are necessary for the researchers.

75 This project aims to compare the fitness of 4 mathematical models (two mechanistic

76 model: Logistic equation and the Buchanan model; two phenomenological models: the
77 Gompertz model and the Baranyi model) based on a large dataset of microbial popula-
78 tion growth cases. Moreover, this paper will also discuss if there is an existence of the
79 interaction between temperature and model fitting, given that the temperature factor was
80 not written in any model. The hypotheses are: (1) Mechanistic models fit better than
81 phenomenological ones in a large number of cases (2) Gompertz model fit better than
82 the Baranyi model (3) Temperature does not affect which population growth rate model
83 fits best

84 **2 Methods**

85 **2.1 Data and preparation before the modelling**

86 The original dataset called *LogisticGrowthData.csv* was used in this project. Collected
87 from different studies, it contains measurements of change in biomass or number of cells
88 of microbes over time. These data were collected through lab experiments across the
89 world. The field names were defined in a file called *LogisticGrowthMetaData.csv*, also
90 in the data directory. The two main fields of interest are *PopBio(abundance)* and *Time*.

91 Data were modified with unique single IDs and unsuitable and low-quality data were
92 deleted. Single population growth rate curves can be identified as unique temperature-
93 species-medium-citation-replicate combinations (concatenate them to get a new string
94 variable that identifies unique growth curves). In addition, NA, negative, duplicated data
95 were filtered and removed. Besides, datasets with less than 5 data points were also fil-
96 tered out and removed where 5 is the minimum number of data points needed to fit the
97 models. After the modification, the dataset composed by 4225 observations, 45 species
98 with 275 unique IDs and temperature.

99 2.2 Models

100 2.2.1 Mechanistic Model

101 A classical mechanistic model, is the logistic equation:

$$N_t = \frac{N_0 * N_{max} * e^{r_{max}*t}}{N_{max} + N_0 * (e^{r_{max}*t} - 1)} \quad (1)$$

102 Here N_t is population size at time t , N_0 is initial population size, r is maximum growth
103 rate (AKA r_{max}), and N_{max} is carrying capacity.

104 The Buchanan model, considered as a 'three-phase logistic mode', is relatively simple:

$$N_t = \begin{cases} N_0 & \text{if } t \leq t_{lag} \\ N_{max} + r_{max} * (t - t_{lag}) & \text{if } t_{lag} < t < t_{max} \\ N_{max} & \text{if } t \geq t_{max} \end{cases} \quad (2)$$

105 2.2.2 Phenomenological Model

106 One of popular phenomenological model is called modified Gompertz model(Zwietering
107 et al. 1990):

$$N_t = A * e^{-e^{\frac{r_{max} * e * (t_{lag} - t)}{A} + 1}} \quad (3)$$

108 where r_{max} maximum growth rate, the tangent to the inflection point. t_{lag} is the x-axis
109 intercept to this tangent (duration of the delay before the population starts growing expo-
110 nentially) and A is the asymptote ($A = \ln(N_{max}/N_0)$), N_0 is initial cell culture (Population)
111 density, N_{max} is maximum population density.

112 Another phenomenological model, called as Baranyi model, introduces a new dimension-
113 less parameter h_0 which represents the initial physiological state of the cells. The length
114 of the lag phase is determined by the value of h_0 at inoculation and the post-inoculation
115 environment. Thus the definition of lag is independent from the shape of the growth curve,
116 and the effect of the previous environment is separated from the effects of the present
117 environment. This allows modeling growth without a lag period following inoculation from

media favorable to growth to new media also favorable to growth. r_{max} and h_0 can be related to obtain the lag time, t_{lag} . The formulation given is:

$$N_t = N_0 + r_{max} * A_t - \ln(1 + \frac{e^{r_{max}*A_t} - 1}{e^{N_{max}-N_0}}) \quad (4)$$

where:

$$A_t = t + (\frac{1}{r_{max}}) * \ln(\frac{e^{-r_{max}*t} + h_0}{1 + h_0}) \quad (4.2)$$

$$t_{lag} = \frac{\ln(1 + \frac{1}{h_0})}{r_{max}} \quad (4.3)$$

2.3 Model fitting and Visualization

A R script was written to get the model fitting named *Model_fitting.R*. Before fitting models to the dataset, the starting value of the parameters were calculated. For added complication, note that the biomass are written in log to the base 10, which means that data transformed to log base 10. These could also be written in the natural log too. The r_{max} was derived from linear model. Start N_{max} value was calculated by maximum of Log10N, while Start N_0 value was calculated by minimum of Log10N. Then 4 models mentioned above were used to investigate the model fitting with a loop. In the meanwhile, AIC and BIC for each model were calculated and stored in a file named *statistic_data.csv* in the Data folder. Consequently, dataset of each ID was plotted to figure. Those figures were saved as *respectiveplots* to the *Results* folder.

2.4 Model selection criteria

4 models were compared and evaluated by Akaike information criteria(AIC) (Johnson & Omland 2004) and Bayesian information criterion (BIC) criteria (Ahmadi 2008). The best model is the model with most support with lowest AIC or BIC. The best fitted models were counted and calculated the percentage for all datasets using the R script *results_plotting.R*. Finally, the data after criteria were plotted and saved to the folder *Results*.

138 **2.5 Computing languages**

139 Bash 3.2 : A file *compile.sh* written in bash was used to compile the report into pdf format
140 with bibliography. Another file *run_miniproject.sh* written in bash aimed to run all the files
141 and get the final results and report.

142 R 3.6.1: It was applied to get the data preparation, calculate the criteria value, as well as
143 plot the results. Some helpful packages were used. *dplyr* was used to manage tables,
144 *ggplot2* was used for plotting, and *minpack.lm* was used to conduct non-linear models.

145 It needs to be pointed that, to take the most efficient work, in this research Python was
146 not used, but It is undeniable that python still has related similar functions.

147 **3 Results**

148 **3.1 Fitting percentage of 4 models**

149 The four models are compared with respect to the probability of fitting to 275 data sets.
150 It can be clearly seen in Figure 2 that the mechanistic model (Logistic) has the highest
151 match rate (98.5%). Two phenomena models (Baranyi and Gompertz) are also very
152 high, which have 76.7% and 88.7% respectively. However, the fitting percentage of the
153 Buchanan model is exceptionally low, indicating that less than half of the data set can be
154 adapted to this model (41.8%).

155 **3.2 Best fitting model assessment**

156 Figure 3 shows the percentage of each model that was rated as the best fitting. It can be
157 seen that both phenomena models (Baranyi and Gompertz) are higher than mechanical
158 models (Logistic and Buchanan), of which the Gompertz model (37.5% of AIC and 36.7%
159 of BIC) has the remarkably highest optimal rate. Whereas, the Buchanan model still gets
160 a very deficient optimal percentage(10.2% of AIC and 9.5% of BIC). There are some
161 differences between the fitness of the models and neither can consistently offer the best
162 fit to almost every growth curve.




Figure 2: Figure 2. Percentage of model fitting

3.3 Four model fittings on one species with different temperatures

One species with different temperatures was selected randomly as an example to present the model fitting situations about this unique ID. The species is *Staphylococcus spp.*, whose database is focusing on the bacterial growth on the raw Chicken Breast (Galarz et al. 2015). This measurement has 6 types of temperature. Figure 4 shows that in this example, the result is that most models still have a relatively good fit at different temperatures. Therefore, specifically from this example, it is undeniable that the effect of temperature on the fitness of the model is basically non-existent. Moreover, it can be seen from Figure 4. b that when the temperature is 4 degrees Celsius, the data is relatively small, and the Buchanan model deviates significantly from the actual situation. It is worth mentioning that both in Figure 4.b and 4.c that there is a downward trend at the end period of each curve, but each model does not fit well and does not reflect the decreasing part. Furthermore, it also could be found that the Logistic model has an upturned part at the beginning of Figure 4.a, c, d, and e. Compared with other models, it does not fit well in the lag-time duration.




Figure 3: Percentage of Best model fitting for each model

4 Discussion

4.1 Difference between phenomenological and mechanistic models

The results in 3.1 show that when studying a large number of data sets, the Logistic model has the highest capacity for fitting probability. The reason is a mechanical model starting from the most basic growth theory. The results of 3.2 show that though the model is not as high as the mechanistic model, the Gompertz model still obtains the highest optimal adaptation rate, which is consistent with previous research that phenomenological models often perform better than mechanistic ones, for the former ones have less restrictive assumptions and uncertainty (Bokulich 2011, Chowell et al. 2016, Eskola & Parvinen 2007). Thus, it will not tend to be applied on a large scale in modeling microbial growth and casualty. In addition, it is possible that although it totally lacks the mechanistic interpretation, the equation or the other empirical 4-parameter models, can be applied to fitting the experimental growth (Transtrum & Qiu 2016).

One of the standards of evaluating the phenomenological model is whether it is convenient to apply and accurate to offer a fit, without the basis of any basic principles or mechanistic considerations (Peleg & Corradini 2011). For instance, the parameters of the




Figure 4: Four model fittings on one species (*Staphylococcus* spp.) with different temperatures

modified Gompertz equation have not been directly linked with microbial death kinetics when using the logarithmic number of cells (Xiong et al. 1999). The ‘three-phase logistic model’, The Baranyi-Roberts Model presents easy mechanism and intuitive meanings, but shows low fitness and precision. Due to the two ‘If’ statements, the linear model could not be assessed that conveniently for the description and prediction of growth on dynamic conditions and then the transition between one phase and the next one will not tend to be sharp just like those in the other isothermal curves.

4.2 Comparison of 2 phenomenological models (Gompertz model and Baranyi model)

The results also show that the Gompertz model is better than the Baranyi model, both in the possibility of fitting and best fitting rate. However, it is different from a study illustrating that taking the model generated data into consideration in a broad way, the results by Baranyi’s model can be more ideal than the modified Gompertz equation (Juneja et al. 2007, Xiong et al. 1999). Nevertheless, more literature argues the modified Gompertz equation was used primarily in modelling the asymmetrical sigmoidal shape of microbial

209 growth curves (Halmi & Shukor 2014, Lim et al. 1998, McMeekin et al. 2013). The Gom-
210 pertz model can offer a description of the growth curves with a short or long lag time
211 or neither combined with the parameters' adjustments (Peleg & Corradini 2011). How-
212 ever, the Baranyi model is better than the Gompertz one at higher growth rate samples
213 (Baranyi 1993).

214 There are some problems in the Baranyi model, one of which can be the meaning that
215 the lag time defined has a close relationship with the maximum specific growth rate. In
216 some favorable situation for the organism, someone can expect the growth not merely
217 to be more vigorous, but also to begin earlier (manifested in a shorter lag time). Thus,
218 it has discovered a negative relation between that lag time and maximum (Brown 2007).
219 The Baranyi model can make a prediction that the extensions of almost every isothermal
220 (semi-logarithmic) tangent of the growth curve at the inflection point must meet at one
221 point. It is not merely inconsistent by some reported growth data in the article (Huang
222 2008, Xiong et al. 1999), but also proven not to be accurate by simulating realistic growth
223 curves with the Gompertz as a model violating the rule.

224 **4.3 Temperature factor**

225 Although on the basis of the frequent observation, there are not any reasons in the as-
226 pect of theories that the growth parameters are always rising and falling in unison, keep-
227 ing pace with changes in the external condition. Therefore, to quantify the effects of the
228 environmental factors on microbial growth needs, the growth parameter should be as-
229 sessed, not merely considering the maximum specific growth rate. The results vary to
230 the selected methods with determining the "lag time" often yield different values, showing
231 that it is not an inadequate definition of growth parameters. The simultaneous effects of
232 the factors, including T, pH, or a_w , should not be multiplicative but be evaluated via ex-
233 periments (Peleg & Corradini 2011). However, the result 3.3 shows that the temperature
234 has little effect on fitness. The reason may be that several models can be performed at
235 different temperatures. For instance, when the temperature dependence was assessed
236 via a series of experiments, the Buchanan model will be applied to the generation of non-
237 isothermal growth curve to some degree alike that in the biphasic survival curve (Mishra

et al. 2017). Similarly, other models can make predictions about non-isothermal growth patterns according to isothermal data. Therefore, it is practical to select a relatively simple and convenient model (Peleg & Corradini 2011).

4.4 Limitation and suggestions for future studies

This article has many limitations. First of all, there are some data from other literature, and there are strange distributions, which may be due to simultaneous mortality during bacterial growth. Moreover, although AIC and BIC are good candidates for modelling, the criteria used are limited. Moreover, in this experiment, the difference between AIC and BIC has not been thoroughly explored. In the case of extensive data, it is difficult to show all the adapted p-values. Those limitations could be considered in future studies. Furthermore, some research directed that the future of predictive microbiology may be altered from combining a phenomenological and mechanistic basis to complicated systems (McMeekin et al. 2013).

5 Conclusion

In conclusion, the hypothesis that the mechanistic model is better than the phenomenon model is unilateral. It requires different analysis in different situations. In this experiment, a large number of data sets were researched and found that although the logical model has a high probability of being fit, the phenomenon model can better fit the data set and better fit the degree. In addition, the Gompertz model fits best in phenomenological models in a large number of cases.

Moreover, although the temperature did not show a significant impact on model adaptation in the study of a single example, it is recommended to consider the influence of environmental factors in future experiments. Generally, all models are optimized to generate a mortality phase by timing the original growth term by a decay factor, and it will be logistic, exponential, stretched exponential or any algebraic term that begins at 1 and drops to 0. A typical instance can be the model built for complicated chemical reactions and biological processes following the competing mechanisms (Peleg et al. 2009). Al-

though it is evident that the observed growth pattern can be influenced by almost every factor above and their interactions, the shape of the growth curve cannot cover adequate data to assess the physiological influences of the changes singularly (Peleg 2006).

References

- Ahmadi, H., G. A. (2008), 'Non-linear hyperbolastic growth models for describing growth curve in classical strain of broiler chicken', *Research Journal of Biological Sciences* **3**(11), 1300–1304.
- Araujo, R. P. & McElwain, D. L. S. (2004), 'A history of the study of solid tumour growth: the contribution of mathematical modelling', *Bulletin of Mathematical Biology* **66**(5), 1039—1091.
- Baranyi, J., R. T. A. M. P. (1993), 'A non-autonomous differential equation to model bacterial growth', *Food Microbiology* **10**, 43–59.
- Bokulich, A. (2011), 'How scientific models can explain', *Synthese* **180**(1), 33–45.
URL: <https://doi.org/10.1007/s11229-009-9565-1>
- Brown, S. (2007), 'Two implications of common models of microbial growth', *ANZIAM Journal* **49**.
- Buchanan, R. L., Whiting, R. C. & Damert, W. C. (1997), 'When is simple good enough: a comparison of the gompertz, baranyi, and three-phase linear models for fitting bacterial growth curves', *Food Microbiology* **14**(4), 313–326.
- Chowell, G., Hincapie-Palacio, D., Ospina, J., Pell, B., Tariq, A., Dahal, S., Moghadas, S., Smirnova, A., Simonsen, L. & Viboud, C. (2016), 'Using phenomenological models to characterize transmissibility and forecast patterns and final burden of zika epidemics', *PLoS currents* **8**.
- Costabile, L. & Rowthorn., B. (1985), 'Malthus's theory of wages and growth.', *The Economic Journal* **95**(378), 418–437.
- d'Onofrio, A. (2009), 'Fractal growth of tumors and other cellular populations: Linking

the mechanistic to the phenomenological modeling and vice versa', *Chaos, Solitons & Fractals* **41**(2), 875—880.

Eskola, H. T. M. & Parvinen, K. (2007), 'On the mechanistic underpinning of discrete-time population models with allee effect', *Theoretical Population Biology* **72**(1), 41–51.

Fujikawa, H., Kai, A. & Morozumi, S. (2004), 'A new logistic model for escherichia coli growth at constant and dynamic temperatures', *Food Microbiology* **21**(5), 501–509.

Galarz, L. A., Fonseca, G. G. & Prentice, C. (2015), 'Predicting bacterial growth in raw, salted, and cooked chicken breast fillets during storage', *Food Science and Technology International* **22**(6), 461–474.

Grijnspeerdt, K. & Vanrolleghem, P. (1999), 'Estimating the parameters of the baranyi model for bacterial growth', *Food Microbiology* **16**(6), 593–605.

Halmi, M. I. E, S. M. S. J. W. L. W. & Shukor, M. Y. (2014), 'Modeling the growth curves of acinetobacter sp. strain dry12 grown on diesel', *Journal of Environmental Bioremediation & Toxicology* **2**(1), 33—37.

Huang, L. (2008), 'Growth kinetics of listeria monocytogenes in broth and beef frankfurters—determination of lag phase duration and exponential growth rate under isothermal conditions', *Journal of Food Science* **73**(5), E235–E242.

Johnson, J. B. & Omland, K. S. (2004), 'Model selection in ecology and evolution', *Trends in Ecology & Evolution* **19**(2), 101–108.

Juneja, V. K., Valenzuela Melendres, M., Huang, L., Gumudavelli, V., Subbiah, J. & Thippareddi, H. (2007), 'Modeling the effect of temperature on growth of salmonella in chicken', *Food Microbiology* **24**(4), 328–335.

Kacena, M. A., Merrell, G. A., Manfredi, B., Smith, E. E., Klaus, D. M. & Todd, P. (1999), 'Bacterial growth in space flight: logistic growth curve parameters for escherichia coli and bacillus subtilis', *Applied Microbiology and Biotechnology* **51**(2), 229–234.

Lim, C. C. . . ., Arora, R. . . . & Townsend, E. C. . . . (1998), 'Comparing gompertz and richards functions to estimate freezing injury in rhododendron using electrolyte leakage', *Journal of the American Society for Horticultural Science* **123**(2), 246–252.

- 319 McMeekin, T., Olley, J., Ratkowsky, D., Corkrey, R. & Ross, T. (2013), 'Predictive microbi-
320 ology theory and application: Is it all about rates?', *Food Control* **29**(2), 290—299.
- 321 Mishra, A., Guo, M., Buchanan, R. L., Schaffner, D. W. & Pradhan, A. K. (2017), 'Develop-
322 ment of growth and survival models for salmonella and listeria monocytogenes during
323 non-isothermal time-temperature profiles in leafy greens', *Food Control* **71**, 32–41.
- 324 Peleg, M. (2006), *Advanced Quantitative Microbiology for Food and Biosystems: Models
325 for Predicting Growth and Inactivation*, CRC Press.
- 326 Peleg, M. & Corradini, M. G. (2011), 'Microbial growth curves: What the models tell us
327 and what they cannot', *Critical Reviews in Food Science and Nutrition* **51**(10), 917—
328 945.
- 329 Peleg, M., Corradini, M. G. & Normand, M. D. (2007), 'The logistic (verhulst) model for
330 sigmoid microbial growth curves revisited', *Food Research International* **40**(7), 808–
331 818.
- 332 Peleg, M., Corradini, M. G. & Normand, M. D. (2009), 'Isothermal and non-isothermal
333 kinetic models of chemical processes in foods governed by competing mechanisms',
334 *Journal of Agricultural and Food Chemistry* **57**(16), 7377–7386.
335 **URL:** <https://doi.org/10.1021/jf9012423>
- 336 Pörtner, H. O., Schulte, P. M., Wood, C. M. & Schiemer, F. (2010), 'Niche dimensions in
337 fishes: An integrative view', *Physiological and Biochemical Zoology* **83**(5), 808—826.
- 338 Transtrum, M. K. & Qiu, P. (2016), 'Bridging mechanistic and phenomenological models of
339 complex biological systems', *PLoS computational biology* **12**(5), e1004915–e1004915.
- 340 Xiong, R., Xie, G., Edmondson, A. S., Linton, R. H. & Sheard, M. A. (1999), 'Compar-
341 ison of the baranyi model with the modified gompertz equation for modelling thermal
342 inactivation of listeria monocytogenes scott a', *Food Microbiology* **16**(3), 269—279.
- 343 Zwietering, M. H., Jongenburger, I., Rombouts, F. M. & van 't Riet, K. (1990), 'Modeling
344 of the bacterial growth curve', *Applied and Environmental Microbiology* **56**(6), 1875.