

# Causal inference in degenerate systems: An impossibility result

Yue Wang

Nov. 29, 2021

University of California, Los Angeles

Accepted by AISTATS 2020


# Authors

Yue Wang, Postdoctoral Fellow, Department of Computational Medicine, University of California, Los Angeles.  
Email: [yuew@g.ucla.edu](mailto:yuew@g.ucla.edu)



Linbo Wang, Assistance Professor, Department of Statistical Sciences, University of Toronto, Canada.

Email: [linbo.wang@utoronto.ca](mailto:linbo.wang@utoronto.ca)



- What is causal effect.
- Existing causal quantities and their problems.
- Criteria for a “good” causal quantity
- An impossibility theorem.
- Algorithms and simulations.

# What is causal effect

- Heating with fire causes water to boil.
- HIV exposure causes AIDS.
- Smoking causes lung cancer.

# What is causal effect

- If A and B are related, does A cause B?
- A real (indirect) causation: Driving without a seat belt can reduce the risk of cancer.
- Correlation does not imply causation.
- When the rooster crows, the sun rises. (B causes A)
- As ice cream sales increase, the rate of drowning deaths increases sharply. (A hidden factor C causes both A and B)

# What is causal effect

- When A directly causes B, the causal relation can be further classified or quantified.
- Heating with fire causes water to boil. (Deterministic)
- HIV exposure causes AIDS. (Stochastic, strong effect)
- Smoking causes lung cancer. (Stochastic, weak effect)

# What is causal effect

- Skip philosophical discussions (Hume et al.) of causal effect...



# Purpose

- We have some random variables  $X_1, X_2, \dots, X_n, Y$ .
- $X_1, \dots, X_n$  (cause variables) are exactly all the possible direct causes of  $Y$  (result variable). We assume  $Y$  does not cause any  $X_i$ , and there is no hidden cause of  $Y$ .
- Our purpose is to quantify the effect of a direct causal relationship  $X_1 \rightarrow Y$ , based on the joint probability distribution of  $X_1, X_2, \dots, X_n, Y$ .

# Causal quantities

- There are various quantities that measure a causal relation.
- Correlation coefficient and its variations.
- Average treatment effect and its variations.
- Mutual information and its variations.
- Granger causality and transfer entropy (for stochastic processes).

- Idea: if  $X$  causes  $Y$ , then  $X$  contains information of  $Y$ .  $X$  has predict power on  $Y$ . Use information to quantify causal effect.
- Measure of information: entropy.





$$H(X) = - \sum_i p_i \log p_i,$$

where  $p_i = \mathbb{P}(X = x_i)$ .

- $H(X) \geq 0$ . Equality holds if and only if  $X$  is deterministic.

# Mutual information (MI)

$$MI(X, Y) = H(X) + H(Y) - H(X, Y).$$



# Mutual information (MI)

- Intuition: the information shared between  $X$  and  $Y$ . The information gain of  $Y$  if we know  $X$ . The predict power of  $X$  on  $Y$ .
- If  $X$  causes  $Y$ , then  $MI(X, Y)$  can be used to describe the causal effect of  $X \rightarrow Y$ .
- $MI(X, Y) \geq 0$ . Equality holds if and only if  $X$  and  $Y$  are independent.

# Conditional Mutual information (CMI)


- Generalize MI for more variables.

$$\text{CMI}(X_1, Y \mid X_2) = \text{MI}(X_1 X_2, Y) - \text{MI}(X_2, Y).$$

- Conditioned on the knowledge of  $X_2$ , how much extra information of  $Y$  could  $X_1$  provide.
- Can be used to describe the causal effect of  $X_1 \rightarrow Y$  if  $X_1$  and  $X_2$  cause  $Y$ .
- $\text{CMI}(X_1, Y \mid X_2) \geq 0$ . Equality holds if and only if  $X_1$  and  $Y$  are independent conditioned on  $X_2$ . This means that with the knowledge of  $X_2$ ,  $X_1$  contains no new knowledge of  $Y$ .

# Conditional Mutual information (CMI)

$$\text{CMI}(X_1, Y | X_2) = \text{MI}(X_1 X_2, Y) - \text{MI}(X_2, Y).$$




# Conditional Mutual information (CMI)

- Venn diagram does not work.
- $X_1, X_2$  are independent variables, with equal probabilities to take 0 or 1.  $Y = X_1 + X_2 \bmod 2$ .
- Any two of  $X_1, X_2, Y$  are independent, but these two could determine the third variable.
- $MI(X_1, Y) = 0$ ,  $CMI(X_1, Y | X_2) > 0$ .



# Problem of CMI


- What if  $\mathbb{P}(X_1 = X_2) \approx 1$ ?



$$X_2 = X_1 + \epsilon, Y = X_2 + \delta$$

$$X_1 \perp\!\!\!\perp Y \mid X_2$$

$\epsilon$  and  $\delta$  are independent, equal 0 with high probabilities.



$$X_2 = X_1 + \epsilon, Y = X_1 + \delta$$

$$X_2 \perp\!\!\!\perp Y \mid X_1$$

- $\text{CMI}(X_1, Y \mid X_2)$  is 0 in the first case, and 0.0065 in the second case.

# Problem of CMI

**Table:** Joint distributions of  $X_1$ ,  $X_2$ ,  $Y$  in two cases

$X_1$	$X_2$	$Y$	Case 1	Case 2
0	0	0	0.4990005	0.4990005
0	0	1	0.0004995	0.0004995
1	1	0	0.0004995	0.0004995
1	1	1	0.4990005	0.4990005
0	1	0	0.0000005	0.0004995
0	1	1	0.0004995	0.0000005
1	0	0	0.0004995	0.0000005
1	0	1	0.0000005	0.0004995

- Utilize the slight difference between  $X_1$  and  $X_2$ .
- Causal strength (CS):  
D. Janzing, D. Balduzzi, M. Grosse-Wentrup, and B. Schölkopf. Quantifying causal influences. *Ann. Stat.*, 41(5):2324–2358, 2013.
- 


$$\text{CS}(X_1, Y) = \sum_{x_1, x_2, y} \mathbb{P}(x_1, x_2, y) \log \frac{\mathbb{P}(y \mid x_1, x_2)}{\sum_{x'_1} \mathbb{P}(y \mid x'_1, x_2) \mathbb{P}(x'_1)}.$$

- Part mutual information (PMI):  
J. Zhao, Y. Zhou, X. Zhang, and L. Chen. Part mutual information for quantifying direct associations in networks. Proc. Natl. Acad. Sci., 113(18):5130–5135, 2016.



$$\text{PMI}(X_1, Y \mid X_2) = \sum_{x_1, x_2, y} \mathbb{P}(x_1, x_2, y) \log \frac{\mathbb{P}(x_1, y \mid x_2)}{\sum_{x'_1} \mathbb{P}(y \mid x'_1, x_2) \mathbb{P}(x'_1) \sum_{y'} \mathbb{P}(x_1 \mid x_2, y') \mathbb{P}(y')}.$$


# New methods



$$X_2 = X_1 + \epsilon, Y = X_2 + \delta$$

$$X_1 \perp\!\!\!\perp Y \mid X_2$$


In the first case,  $\text{CMI}(X_1, Y \mid X_2)$ ,  $\text{CS}(X_1, Y)$  and  $\text{PMI}(X_1, Y \mid X_2)$  are 0. In the second case,  $\text{CMI}(X_1, Y \mid X_2)$ ,  $\text{CS}(X_1, Y)$  and  $\text{PMI}(X_1, Y \mid X_2)$  are 0.0065, 0.6852, 0.9661.



$$X_2 = X_1 + \epsilon, Y = X_1 + \delta$$


$$X_2 \perp\!\!\!\perp Y \mid X_1$$

# Problem of new methods



$$X_2 = X_1 + \epsilon, Y = X_2 + \delta$$

$$X_1 \perp\!\!\!\perp Y \mid X_2$$



$$X_2 = X_1 + \epsilon, Y = X_1 + \delta$$


$$X_2 \perp\!\!\!\perp Y \mid X_1$$

- These two joint distributions are almost the same, but the resulting causal effects are very different.
- CS and PMI may not be continuous with joint distribution (under total variation distance) when both  $\{X_1\}$  and  $\{X_2\}$  have all the information of  $Y$  contained in  $\{X_1, X_2\}$ .

# Markov boundary (MB)

- Markov chain  $Z(t-2) \rightarrow Z(t-1) \rightarrow Z(t) \rightarrow Z(t+1)$ . Use  $\mathcal{S} = \{Z(t-2), Z(t-1), Z(t)\}$  to predict  $Y = Z(t+1)$ :  $\{Z(t)\}$  is enough.
- Markov random field: neighbors are enough to predict one variable.

## Markov Random Fields



Can be generalized to any **undirected** graphs (nodes, edges)

**Neighborhood system:** each node is connected to its neighbors  
neighbors are reciprocal

**Markov property:** each node only depends on its neighbors

Note: the black lines on the left graph are illustrating the 2D grid for the image pixels  
they are not edges in the graph as the blue lines on the right

# Markov boundary (MB)

Markov boundary of  $Y$  within  $\mathcal{S} = \{X_1, \dots, X_n\}$ :  $\mathcal{S}_1 \subset \mathcal{S}$ , which is minimal, and has the same information on  $Y$ . (Remove all redundant variables with no predict power.)

$$\text{CMI}(\mathcal{S} \setminus \mathcal{S}_1, Y \mid \mathcal{S}_1) = 0,$$

$$\forall \mathcal{S}_2 \subsetneq \mathcal{S}_1, \quad \text{CMI}(\mathcal{S} \setminus \mathcal{S}_2, Y \mid \mathcal{S}_2) > 0.$$

This means  $Y \perp\!\!\!\perp \mathcal{S} \setminus \mathcal{S}_1 \mid \mathcal{S}_1$ .



# Markov boundary (MB)

- MB may not be unique. (Set  $X_1 = X_2$  in the above example, then both  $\{X_1\}$  and  $\{X_2\}$  are MB of  $Y$  within  $\{X_1, X_2\}$ .)
- Statnikov et al. (2013) studied 13 benchmark data sets from reality, and found that five of them have multiple MB.
- Assume MB is unique. A cause variable inside MB has non-zero unique information of  $Y$ , and a cause variable outside MB has zero unique information of  $Y$ . Therefore the unique MB should be exactly all the cause variables with non-zero causal effect.

# Problem of new methods

- When there are multiple MB (called “degenerate”), both CS and PMI are not directly defined. (Contains  $0/0$  in the expression.)
- Try to use continuation: choose a sequence of distributions (for which CS and PMI are defined) converging to the original distribution, and check whether the corresponding CS and PMI converge.

# Problem of new methods

Consider a probability distribution  $p$  on  $\mathcal{S} \cup Y$ , under which  $Y$  has multiple MB in  $\mathcal{S}$ , and  $X_1$  is in at least one, but not all of such MB. Now  $\text{CMI}(X_1, Y \mid \mathcal{S} \setminus \{X_1\}) = 0$ .

## Theorem (Wang & Wang, 2020)

*There exist constants  $c_1 < c_2$ , such that in any arbitrarily small neighborhood of  $p$ ,  $\text{CS}(X_1 \rightarrow Y)$  takes any value between  $c_1$  and  $c_2$ . The same applies to PMI.*

# Problem of new methods

- When there are multiple MB, both CS and PMI cannot be well-defined.
- Similar to the behavior of a complex analytical function near an essential singularity (Picard's great theorem).
- It is not numerically feasible to calculate CS and PMI in such cases.

# Sketch of the proof

## Lemma (Wang & Wang, 2020)

*Assume  $Y$  has multiple MB,  $X_1$  is inside some MB but not all MB, and set  $X_2$  to be all other variables. Then there exist  $x'_1, x_2$  such that  $\mathbb{P}(x'_1) > 0$ ,  $\mathbb{P}(x_2) > 0$ ,  $\mathbb{P}(x'_1, x_2) = 0$ .*

Under small perturbations,  $\mathbb{P}(x'_1, x_2)$  is very small but positive, so that CS and PMI can be defined, but  $\mathbb{P}(y \mid x'_1, x_2)$  can change significantly.

# Sketch of the proof



$$CS(X_1, Y) = \sum_{x_1, x_2, y} \mathbb{P}(x_1, x_2, y) \log \frac{\mathbb{P}(y \mid x_1, x_2)}{\sum_{x'_1} \mathbb{P}(y \mid x'_1, x_2) \mathbb{P}(x'_1)}.$$



$$\begin{aligned} \text{PMI}(X_1, Y \mid X_2) = \\ \sum_{x_1, x_2, y} \mathbb{P}(x_1, x_2, y) \log \frac{\mathbb{P}(x_1, y \mid x_2)}{\sum_{x'_1} \mathbb{P}(y \mid x'_1, x_2) \mathbb{P}(x'_1) \sum_{y'} \mathbb{P}(x_1 \mid x_2, y') \mathbb{P}(y')} \end{aligned}$$

- If  $\mathbb{P}(x'_1)$  and  $\mathbb{P}(x_2)$  are not small, but  $\mathbb{P}(x'_1, x_2)$  is very small, then changing  $\mathbb{P}(y \mid x'_1, x_2)$  has a significant impact on CS and PMI, but the whole distribution is perturbed slightly.

# Sketch of the proof

- Construct two sequences of distributions, both of which converge to the original distribution.
- CS (or PMI) of two sequences always exist, but converge to different values.
- Distribution of one sequence can continuously transform into distribution of the other sequence, during which CS (or PMI) is always defined.
- When there are multiple MB, CS and PMI are not well defined, and CMI is zero.
- Questions?

# Purpose

- We have cause variables  $\mathcal{S} = \{X_1, X_2, \dots, X_n\}$  and result variable  $Y$ .
- Our purpose is to quantify the effect of a causal relationship  $X_1 \rightarrow Y$ , based on the joint distribution of  $X_1, X_2, \dots, X_n, Y$
- Existing causal quantities have different problems.
- We propose several criteria for a “good” causal quantity.
- We focus on the case where MB is unique. In such case, the unique MB should be exactly all the variables with non-zero causal effect.



# Criteria for quantifying causal effect

- C1. The strength of  $X \rightarrow Y$  is a continuous function of the joint distribution of  $Y$  and  $S$ , under the total variation distance.
- C2. If there is a unique MB  $\mathcal{M}$  of  $Y$  within  $S$ , and  $X \notin \mathcal{M}$ , then the strength of  $X \rightarrow Y$  is 0.
- C3. If there is a unique MB  $\mathcal{M}$  of  $Y$  within  $S$ , and  $X \in \mathcal{M}$ , then the absolute value of the strength of  $X \rightarrow Y$  is at least  $\text{CMI}(X, Y \mid \mathcal{M} \setminus \{X\})$ .
- In C3,  $\text{CMI}(X, Y \mid \mathcal{M} \setminus \{X\})$  can be replaced by any positive constant, which only depends on  $X, Y, \mathcal{M} \setminus \{X\}$ .

# Criteria for quantifying causal effect

- CMI satisfies C1 and C2.
- CS and PMI satisfies C2 and C3.
- A naive causal effect measure that takes a large positive constant value (such as  $2H(Y)$ ) satisfies C1 and C3.

# An impossibility theorem

Consider a probability distribution  $p$  on  $\mathcal{S} \cup Y$ , under which  $Y$  has multiple MB in  $\mathcal{S}$ , and  $X$  is in at least one, but not all of such MB. We want to define a quantity for  $X \rightarrow Y$  that can be calculated from  $p$ .

## Theorem (Wang & Wang, 2020)

*In any neighborhood  $\mathfrak{N}$  of  $p$ , all identifiable measures of the strength of  $X \rightarrow Y$  must violate at least one of the criteria in C1 – C3.*

# Sketch of proof

For variable  $X_1$ , we define  $X_1$  with  $\epsilon$ -noise to be  $X_1^\epsilon$ , which equals  $X_1$  with probability  $1 - \epsilon$ , and equals an independent noise with probability  $\epsilon$ . Denote all cause variables by  $\mathcal{S}$ .

Lemma (Strict Data Processing Inequality, Wang & Wang, 2020)

*$\mathcal{S}_1$  is a group of variables without  $X_1$ ,  $Y$ . If we add  $\epsilon$ -noise on  $X_1$  to get  $X_1^\epsilon$ , then  $\text{CMI}(X_1^\epsilon, Y \mid \mathcal{S}_1) \leq \text{CMI}(X_1, Y \mid \mathcal{S}_1)$ , and the equality holds if and only if  $\text{CMI}(X_1, Y \mid \mathcal{S}_1) = 0$ .*

Lemma (Wang & Wang, 2020)

*Assume  $Y$  has multiple MB. For one MB  $\mathcal{M}_0$ , if we add  $\epsilon$  noise on all variables of  $\mathcal{S} \setminus \mathcal{M}_0$ , then in the new distribution,  $\mathcal{M}_0$  is the unique MB.*

# Sketch of proof

- Assume  $X_1 \in \mathcal{M}_0$ ,  $X_1 \notin \mathcal{M}_1$  for MB  $\mathcal{M}_0, \mathcal{M}_1$ .
- We can add  $\epsilon$ -noise on  $\mathcal{S} \setminus \mathcal{M}_1$ , such that  $\mathcal{M}_1$  is the unique MB. Criterion C2 shows that the effect of  $X_1^\epsilon \rightarrow Y$  is 0.
- We can add  $\epsilon$ -noise on  $\mathcal{S} \setminus \mathcal{M}_0$ , such that  $\mathcal{M}_0$  is the unique MB. Criterion C3 shows that the absolute value of the effect of  $X_1 \rightarrow Y$  is at least  $\text{CMI}(X_1, Y \mid \mathcal{M}_0 \setminus \{X_1\}) > 0$ .
- Let  $\epsilon \rightarrow 0$ . Criterion C1 shows that the absolute value of the effect of  $X_1 \rightarrow Y$  should be at least  $\text{CMI}(X_1, Y \mid \mathcal{M}_0 \setminus X)$ , and should be 0.

- Quantifying causal effect with multiple MB is an essentially ill-posed problem.
- When a distribution with unique MB is close to another distribution with multiple MB, a reasonable causal quantity can be very small (CMI) or fluctuate violently (CS, PMI). Therefore in such case, quantitative method is not feasible.
- Questions?

# Uniqueness of MB

- In practice, whether should we apply quantitative methods?
- We need practical methods to determine the uniqueness of MB.
- First step: algorithms that can produce one MB.
- Known methods: IAMB, KIAMB, Semi-Interleaved HITON-PC, MBOR, BLCD, PCMB, GLL-PC. All of them have additional requirements on the joint distribution (to exclude morbid cases).
- We propose Algorithm 1, using the idea of discarding redundant variables one by one.

# Uniqueness of MB

Algorithm 1: An assumption-free algorithm for producing one MB

(1) **Input**

Joint distribution of  $\mathcal{S} = \{X_1, \dots, X_k\}$  and  $Y$

(2) **Set**  $\mathcal{M}_0 = \mathcal{S}$

(3) **Repeat**

**Set**  $X_0 = \arg \min_{X \in \mathcal{M}_0} \text{CMI}(X, Y \mid \mathcal{M}_0 \setminus \{X\})$

**If**  $X_0 \perp\!\!\!\perp Y \mid \mathcal{M}_0 \setminus \{X_0\}$

**Set**  $\mathcal{M}_0 = \mathcal{M}_0 \setminus \{X_0\}$

**Until**  $X_0 \not\perp\!\!\!\perp Y \mid \mathcal{M}_0 \setminus \{X_0\}$

(4) **Output**  $\mathcal{M}_0$  is a MB



# Uniqueness of MB

- Theoretical results on the uniqueness of MB.
- Construct the “essential set”  $\mathcal{E}$ :  $X_i \in \mathcal{E}$  if and only if  $MI(\mathcal{S} \setminus \{X_i\}, Y) < MI(\mathcal{S}, Y)$ .

Lemma (Wang & Wang, 2020)

*$\mathcal{E}$  is the intersection of all MB.*

Theorem (Wang & Wang, 2020)

*MB is unique if and only if  $\mathcal{E}$  is an MB.*

- Based on such results, we have several algorithms of determining the uniqueness of MB.

# Uniqueness of MB

- Algorithm 2: use another algorithm to produce one MB, and check whether each variable in this MB is essential.
- The checking is not very straightforward, since we want to avoid testing conditional independence for too many variables.
- This is equivalent to executing TIE\* algorithm until producing the second MB, or finishing with the unique MB.

# Uniqueness of MB

Algorithm 2: A general algorithm for determining uniqueness of MB

(1) **Input**

Joint distribution of  $\mathcal{S} = \{X_1, \dots, X_k\}$  and  $Y$   
An algorithm  $\Omega$  which could produce one MB correctly

(2) **Set**  $\mathcal{M}_0 = \{X_1, \dots, X_m\}$  to be the result of Algorithm  $\Omega$  on  $\mathcal{S}$

(3) **For**  $i = 1, \dots, m,$

**Set**  $\mathcal{M}_i$  to be the result of Algorithm  $\Omega$  on  $\mathcal{S} \setminus \{X_i\}$

**If**  $Y \perp\!\!\!\perp \mathcal{M}_0 \mid \mathcal{M}_i$

**Output**  $Y$  has multiple MB

**Terminate**

(4) **Output**  $Y$  has a unique MB

# Uniqueness of MB

- Use Algorithm 1 as “ $\Omega$ ” in Algorithm 2, to get “Alg. 2-AF”, an assumption free algorithm for determining MB uniqueness.
- Use KIAMB as “ $\Omega$ ” in Algorithm 2, to get “Alg. 2-KI”, an algorithm for determining MB uniqueness (requires composition property).
- We propose two more algorithms for better comparison.

# Uniqueness of MB

Algorithm S1: An assumption-free algorithm for determining the uniqueness of MB

Using Algorithm 1 to find one MB, and directly check whether each variable in this MB is essential.

(1) **Input**

Joint distribution of  $\mathcal{S} = \{X_1, \dots, X_k\}$  and  $Y$

(2) **Set**  $\mathcal{M}_0 = \{X_1, \dots, X_m\}$  to be the result of  
Algorithm 1 on  $\mathcal{S}$

(3) **For**  $i = 1, \dots, m,$

**If**  $X_i \perp\!\!\!\perp Y \mid \mathcal{S} \setminus \{X_i\}$

**Output**  $Y$  has multiple MB

**Terminate**

(4) **Output**  $Y$  has a unique MB

# Algorithms

Algorithm S2: An assumption-free algorithm for determining the uniqueness of MB

Directly build the essential set, and check if it is an MB.

(1) **Input**

Joint distribution of  $\mathcal{S} = \{X_1, \dots, X_k\}$  and  $Y$

(2) **Set**  $\hat{\mathcal{E}} = \emptyset$

(3) **For**  $i = 1, \dots, k,$

**If**  $X_i \not\perp\!\!\!\perp Y \mid \mathcal{S} \setminus \{X_i\}$   
 $\hat{\mathcal{E}} = \hat{\mathcal{E}} \cup \{X_i\}$

(4) **If**  $Y \perp\!\!\!\perp \mathcal{S} \mid \hat{\mathcal{E}}$

**output:**  $Y$  has a unique MB


**Else**

**output:**  $Y$  has multiple MB

# Simulation setup

- Implement Alg. 2-AF, Alg. 2-KI, Alg. S1 and Alg. S2.
- Test on four artificial cases.
- In Case 1 and Case 3, MB is unique. In Case 2 and Case 4, MB is not unique.
- In Case 3 and Case 4, the assumption of KIAMB, i.e. the composition property, is failed (thus Alg. 2-KI is failed).


# Algorithms performances



Case 1. Success rates of Alg. 2-AF (blue '+'); Alg. 2-KI (green 'x'); Alg. S1 (black '◇'); Alg. S2 (red '○') with different numbers of observations. Number of observations is in logarithm.




# Algorithms performances




Case 2. Success rates of Alg. 2-AF (blue '+'); Alg. 2-KI (green '×'); Alg. S1 (black '◇'); Alg. S2 (red '○') with different numbers of observations. Number of observations is in logarithm.

# Algorithms performances



Case 3. Success rates of Alg. 2-AF (blue '+'); Alg. 2-KI (green 'x'); Alg. S1 (black '◇'); Alg. S2 (red 'o') with different numbers of observations. Number of observations is in logarithm.

# Algorithms performances



Case 4. Success rates of Alg. 2-AF (blue '+'); Alg. 2-KI (green 'x'); Alg. S1 (black '◇'); Alg. S2 (red 'o') with different numbers of observations. Number of observations is in logarithm.

# Algorithms performances

- Performance: In Case 1 and Case 2, where the composition property holds (KIAMB is valid), Alg. 2-KI is slightly better than Alg. 2-AF, and both are much better than Alg. S1 and Alg. S2.
- Performance: In Case 3 and Case 4, where the composition property fails, Alg. 2-KI fails to produce correct results, while Alg. 2-AF exhibits the best performance.
- In practice, if one has a strong belief in the composition property, then we recommend Alg. 2-KI. Otherwise Alg. 2-AF is preferable.
- Questions?

- When there are multiple MB, CS and PMI are not well-defined.
- When there are multiple MB, we cannot define a “good” causal quantity. (Use qualitative methods instead.)
- Theoretical and practical methods for determining the uniqueness of MB.
- Be aware of multiple MB!

- Our work: Wang, Y., & Wang, L. (2020). Causal inference in degenerate systems: An impossibility result. In International Conference on Artificial Intelligence and Statistics (pp. 3383-3392). PMLR.
- CMI: DOBRUSHIN, R. L. (1963). General formulation of Shannon's main theorem in information theory. Amer. Math. Soc. Trans. 33, 323–438.
- CS: JANZING, D., BALDUZZI, D., GROSSE-WENTRUP, M. & SCHÖLKOPF, B. (2013). Quantifying causal influences. Ann. Stat. 41, 2324–2358.

- PMI: ZHAO, J., ZHOU, Y., ZHANG, X. & CHEN, L. (2016). Part mutual information for quantifying direct associations in networks. *Proc. Natl. Acad. Sci.* 113, 5130–5135.
- MB: PEARL, J. (1988). *Probabilistic Reasoning in Intelligent Systems: Networks of Plausible Inference*.
- KIAMB: PEÑA, J.M., NILSSON, R., BJORKEGREN, J. & TEGNER, J. Towards scalable and data efficient learning of Markov boundaries. (2007). *Int. J. Approx. Reason.*, 45(2): 211–232.
- TIE\*: STATNIKOV, A., LYTKIN, N.I., LEMEIRE, J. & ALIFERIS, C.F. Algorithms for discovery of multiple Markov boundaries. (2013). *J. Mach. Learn. Res.* 14, :499–566.

# Thank you!