

Assignment 1 for CMPUT 652 RL with Robots

October 5, 2019

Total 110 points. Complete within the given space using \LaTeX or handwritten notes. Show the derivations or the steps for obtaining partial points. On the other hand, mistakes, missteps or bad reasoning behind a correct answer will result in points deducted. Submit at armahmood@ualberta.ca

- 1** Consider the supervised regression problem of predicting an output Y from a vector input \mathbf{x} using a linear function with a vector parameter \mathbf{w} . We found in the lecture that the Mean Squared Error (MSE) $E[(Y - \mathbf{x}^\top \mathbf{w})^2]$ is a reasonable objective for this problem. Derive the Least-Squares (LS) method, defined below, from MSE (5 points).

$$\mathbf{w}_{t+1}^{\text{LS}} = \left(\frac{1}{t} \sum_{k=1}^t \mathbf{x}_k \mathbf{x}_k^\top \right)^{-1} \left(\frac{1}{t} \sum_{k=1}^t \mathbf{x}_k Y_k \right). \quad (1)$$

$$\begin{aligned} E[(Y - \mathbf{x}^\top \mathbf{w})^2] &= E[Y^2 + \mathbf{w}^\top \mathbf{x} \mathbf{x}^\top \mathbf{w} - 2\mathbf{w}^\top \mathbf{x} Y] \\ &= E[Y^2] + \mathbf{w}^\top E[\mathbf{x} \mathbf{x}^\top] \mathbf{w} - 2\mathbf{w}^\top E[\mathbf{x} Y] \end{aligned}$$

The exact solution can be obtained by setting the derivative equal to 0 and the Least-Square solution can be obtained by substituting the expected values with sample average:

$$\begin{aligned} \nabla_{\mathbf{w}} E[(Y - \mathbf{x}^\top \mathbf{w})^2] &= 2E[\mathbf{x} \mathbf{x}^\top] \mathbf{w} - 2E[\mathbf{x} Y] = 0 \\ \mathbf{w}^* &= E[\mathbf{x} \mathbf{x}^\top]^{-1} E[\mathbf{x} Y] \\ \mathbf{w}_{t+1}^{\text{LS}} &= \left(\frac{1}{t} \sum_{k=1}^t \mathbf{x}_k \mathbf{x}_k^\top \right)^{-1} \left(\frac{1}{t} \sum_{k=1}^t \mathbf{x}_k Y_k \right) \end{aligned}$$

- 2** Continuing with the supervised regression problem, derive the Stochastic Gradient Descent (SGD) method, defined below, from MSE (5 points).

$$\mathbf{w}_{t+1} = \mathbf{w}_t + 2\alpha_t (Y_t - \mathbf{x}_t^\top \mathbf{w}_t) \mathbf{x}_t. \quad (2)$$

$$\begin{aligned} E[(Y - \mathbf{x}^\top \mathbf{w})^2] &= E[Y^2 + \mathbf{w}^\top \mathbf{x} \mathbf{x}^\top \mathbf{w} - 2\mathbf{w}^\top \mathbf{x} Y] \\ &= E[Y^2] + \mathbf{w}^\top E[\mathbf{x} \mathbf{x}^\top] \mathbf{w} - 2\mathbf{w}^\top E[\mathbf{x} Y] \end{aligned}$$

Stochastic Gradient Descent can be obtained from approximate the exact gradient with one sample:

$$\begin{aligned} \nabla_{\mathbf{w}} E[(Y - \mathbf{x}^\top \mathbf{w})^2] &= 2E[\mathbf{x} \mathbf{x}^\top] \mathbf{w} - 2E[\mathbf{x} Y] \\ &\approx 2(\mathbf{x}_t \mathbf{x}_t^\top \mathbf{w} - \mathbf{x}_t Y_t) \\ &\approx -2(Y_t - \mathbf{x}_t^\top \mathbf{w}_t) \mathbf{x}_t \\ \mathbf{w}_{t+1} &= \mathbf{w}_t + 2\alpha_t (Y_t - \mathbf{x}_t^\top \mathbf{w}_t) \mathbf{x}_t \end{aligned}$$

- 3 According to the strong law of large numbers, sample mean is a consistent estimator of the expected value (See Lemma 3, pg 18, of [Mahmood 2017](#)). Use this fact to argue that the sample MSE $\frac{1}{t} \sum_{k=1}^t (Y_k - \mathbf{x}_k^\top \mathbf{w})^2$ objective is a consistent estimator of MSE (5 points).

- 4 Similarly, argue that LS is a consistent estimator of the solution to MSE (5 points).

- 5** Establish a relationship between MSE and the following: $E_{\mathbf{x}} \left[(E_{Y|\mathbf{x}}[Y] - \mathbf{x}^\top \mathbf{w})^2 \right]$. Note that MSE can be written as $E[(Y - \mathbf{x}^\top \mathbf{w})^2] = E_{\mathbf{x}} E_{Y|\mathbf{x}}[(Y - \mathbf{x}^\top \mathbf{w})^2]$. Also, the expectations E_X and $E_{Y|X}$ can be elaborated as $E_X f(X) = \int_{\mathcal{X}} p^*(x) f(x) dx$ and $E_{Y|X} f(Y) = \int_{\mathcal{Y}} p^*(y|x) f(y) dy$ (5 points).

$$\begin{aligned}
E[(Y - \mathbf{x}^\top \mathbf{w})^2] &= E_{\mathbf{x}} E_{Y|\mathbf{x}}[(Y - \mathbf{x}^\top \mathbf{w})^2] \\
&= \int_{\mathcal{X}} p^*(x) \int_{\mathcal{Y}} p^*(y|x) (y - \mathbf{x}^\top \mathbf{w})^2 dy dx \\
&= \int_{\mathcal{X}} p^*(x) \int_{\mathcal{Y}} p^*(y|x) y^2 - 2yx^\top \mathbf{w} + \mathbf{w}^\top x x^\top \mathbf{w} dy dx \\
&= \int_{\mathcal{X}} p^*(x) [E_{Y|\mathbf{x}}[Y^2] - 2E_{Y|\mathbf{x}}[Y] \mathbf{x}^\top \mathbf{w} + \mathbf{w}^\top x x^\top \mathbf{w}] dx \\
&= \int_{\mathcal{X}} p^*(x) [E_{Y|\mathbf{x}}[Y]^2 - 2E_{Y|\mathbf{x}}[Y] \mathbf{x}^\top \mathbf{w} + \mathbf{w}^\top x x^\top \mathbf{w}] dx \\
&\quad \text{the fact that } Y \text{ is independent with itself given } \mathbf{x} \text{ is used} \\
&= \int_{\mathcal{X}} p^*(x) (E_{Y|\mathbf{x}}[Y] - \mathbf{x}^\top \mathbf{w})^2 dx \\
&= E_{\mathbf{x}} \left[(E_{Y|\mathbf{x}}[Y] - \mathbf{x}^\top \mathbf{w})^2 \right]
\end{aligned}$$

- 6** Objective functions serve many purposes. They bring specificity to the problem formulation, provide a criteria for evaluation and can be used as a basis for deriving new algorithms. Accordingly, the same problem may have different associated objectives. For example, the sample MSE is widely used to evaluate the performance of a learning method, but not all methods ensue from MSE. Consider the following objective MSE_{L2} , which is a slight modification to MSE: $\text{MSE}_{L2}(\mathbf{w}) = E[(Y - \mathbf{x}^\top \mathbf{w})^2 + \lambda \mathbf{w}^\top \mathbf{w}]$. Derive the following method know as ridge regression from MSE_{L2} (5 points).

$$\mathbf{w}_{t+1}^{\text{RR}} = \left(\frac{1}{t} \sum_{k=1}^t \mathbf{x}_k \mathbf{x}_k^\top + \lambda \mathbf{I} \right)^{-1} \left(\frac{1}{t} \sum_{k=1}^t \mathbf{x}_k Y_k \right), \text{ where } \mathbf{I} \text{ is the identity matrix.} \quad (3)$$

$$\text{MSE}_{L2}(\mathbf{w}) = E[Y^2 - 2Y\mathbf{x}^\top \mathbf{w} + \mathbf{w}^\top \mathbf{x} \mathbf{x}^\top \mathbf{w} + \lambda \mathbf{w}^\top \mathbf{w}]$$

\mathbf{w} can be solved by setting the derivative equal to 0 and the required result can be obtained by substituting the expected value with sample average:

$$\begin{aligned}
\nabla_{\mathbf{w}} \text{MSE}_{L2}(\mathbf{w}) &= E[-2\mathbf{x}Y + 2\mathbf{x} \mathbf{x}^\top \mathbf{w} + 2\lambda \mathbf{w}] = 0 \\
E[\mathbf{x}Y] &= E[\mathbf{x} \mathbf{x}^\top + \lambda \mathbf{I}] \mathbf{w} \\
\mathbf{w} &= E[\mathbf{x} \mathbf{x}^\top + \lambda \mathbf{I}]^{-1} E[\mathbf{x}Y] \\
\mathbf{w}_{t+1}^{\text{RR}} &= \left(\frac{1}{t} \sum_{k=1}^t \mathbf{x}_k \mathbf{x}_k^\top + \lambda \mathbf{I} \right)^{-1} \left(\frac{1}{t} \sum_{k=1}^t \mathbf{x}_k Y_k \right)
\end{aligned}$$

- 9 The expected policy gradient update for contextual bandit problem can be given as follows: $E[-\log \pi_\theta(A|X)R]$, where X is a given context, π_θ is the parameterized policy distribution maintained by the agent, A is the action chosen by the agent according to π_θ , and R is the subsequent reward. Show that this update can be derived by calculating the gradient of the cross entropy objective: $CE(\theta) = -\int_{\mathcal{X}} p(x) \int_{\mathcal{A}} \pi_\theta(a|x) \log \pi^*(a|x) da dx$, where p , which is not a function of θ , is the distribution of context X . Also, π^* is the ideal policy distribution defined as the Boltzmann distribution $\pi^*(a|x) = Z^{-1} e^{q(x,a)\tau^{-1}}$, where $q(x,a) = E[R|A=a, X=x]$ is the contextual action value. (10 points).

Similar to Q8, we insert the Boltzmann distribution $\pi^*(a|x) = Z^{-1} e^{q(x,a)\tau^{-1}}$ in to the Cross-Entropy objective:

$$\begin{aligned}
CE(\theta) &= -\int_{\mathcal{X}} p(x) \int_{\mathcal{A}} \pi_\theta(a|x) \log \pi^*(a|x) da dx \\
&= -\int_{\mathcal{X}} p(x) \int_{\mathcal{A}} \pi_\theta(a|x) \log Z^{-1} e^{q(x,a)\tau^{-1}} da dx \\
&= -\int_{\mathcal{X}} p(x) \int_{\mathcal{A}} \pi_\theta(a|x) (\tau^{-1} q(x,a) - \log Z) da dx \\
&= \log Z - \tau^{-1} \int_{\mathcal{X}} p(x) \int_{\mathcal{A}} \pi_\theta(a|x) q(x,a) da dx
\end{aligned}$$

We can obtain the same result if we ignore the constant term:

$$\begin{aligned}
\nabla_\theta CE(\theta) &= -\tau^{-1} \int_{\mathcal{X}} p(x) \int_{\mathcal{A}} \nabla_\theta \pi_\theta(a|x) q(x,a) da dx \\
&= -\tau^{-1} \int_{\mathcal{X}} p(x) \int_{\mathcal{A}} \pi_\theta(a|x) \frac{\nabla_\theta \pi_\theta(a|x)}{\pi_\theta(a|x)} \nabla_\theta \pi_\theta(a|x) q(x,a) da dx \\
&= -\tau^{-1} \int_{\mathcal{X}} p(x) \int_{\mathcal{A}} \pi_\theta(a|x) \nabla_\theta \log \pi_\theta(a|x) q(x,a) da dx \\
&= -\tau^{-1} \int_{\mathcal{X}} p(x) E[\nabla_\theta \log \pi_\theta(A|x) q(x,A)] dx \\
&= -\tau^{-1} E[\nabla_\theta \log \pi_\theta(A|X) q(X,A)] \\
&= \tau^{-1} E[-\nabla_\theta \log \pi_\theta(A|X) R]
\end{aligned}$$

- 10 We wrote a PyTorch code for a discrete action bandit problem. Consider a continuous action bandit problem, where actions are distributed normally: $A \sim N(\mu, \sigma^2)$, and the reward is a quadratic function of the action: $R = -(A - 10)^2$. What is the optimal policy in terms of mean μ and standard deviation σ (5 points)? Write a PyTorch code for learning the mean and the standard deviation using policy gradient. Attach the code here together with the curve for estimated mean and standard deviation over time for 100K time steps. The code should run without error in colab and bring the plots. (15 points, total 20).

- 11** For discrete MDP, Derive the following Bellman equation for action value function from its definition: $q_\pi(s, a) = \mathbb{E}[G_t | S_t = s, A_t = a, A_k \sim \pi, \forall k > t]$, where S_t is the state at t , A_t is the action at t , G_t is the return after t , π is the policy, γ is the discount factor, $r(s, a) = \sum_r r \sum_{s'} p(s', r | s, a)$ and $p_\pi(s', a' | s, a) = \sum_r p(s', r | s, a) \pi(a' | s')$ (5 points).

$$q_\pi(s, a) = r(s, a) + \gamma \sum_{s', a'} p_\pi(s', a' | s, a) q_\pi(s', a'). \quad (4)$$

$$\begin{aligned} q_\pi(s, a) &= \mathbb{E}[G_t | S_t = s, A_t = a, A_k \sim \pi, \forall k > t] \\ &= \mathbb{E}_\pi[G_t | S_t = s, A_t = a] \\ &= \mathbb{E}_\pi[R_t + \gamma G_{t+1} | S_t = s, A_t = a] \\ &= \mathbb{E}[R_t | S_t = s, A_t = a] + \gamma \mathbb{E}_\pi[G_{t+1} | S_t = s, A_t = a] \\ &= \sum_r r \sum_{s'} p(s', r | s, a) + \gamma \mathbb{E}[\mathbb{E}_\pi[G_{t+1} | S_{t+1}, A_{t+1}] | S_t = s, A_t = a] \\ &= r(s, a) + \gamma \mathbb{E}[q_\pi(S_{t+1}, A_{t+1}) | S_t = s, A_t = a] \\ &= r(s, a) + \gamma \sum_{s', a'} \sum_r p(s', r | s, a) \pi(a' | s') q_\pi(s', a') \\ &= r(s, a) + \gamma \sum_{s', a'} p_\pi(s', a' | s, a) q_\pi(s', a') \end{aligned}$$

- 12** Write the above Bellman equation in matrix form (5 points).

$$\begin{aligned} v_\pi(s) &= \sum_a \pi(a | s) \sum_{s', r} p(s', r | s, a) (r + \gamma v_\pi(s')) \\ &= \sum_a \pi(a | s) \sum_{s', r} p(s', r | s, a) r + \gamma \sum_a \pi(a | s) \sum_{s', r} p(s', r | s, a) v_\pi(s') \\ &= \sum_a \pi(a | s) r(s, a) + \gamma \sum_{s'} \left[\sum_a \pi(a | s) p(s' | s, a) \right] v_\pi(s') \\ &= r_\pi(s) + \gamma \sum_{s'} p_\pi(s' | s) v_\pi(s') \end{aligned}$$

$$\mathbf{v}_\pi = \mathbf{r}_\pi + \gamma \mathbf{P}_\pi \mathbf{v}_\pi$$

$$\begin{bmatrix} v_\pi(s_0) \\ v_\pi(s_1) \\ \vdots \end{bmatrix} = \begin{bmatrix} r_\pi(s_0) \\ r_\pi(s_1) \\ \vdots \end{bmatrix} + \gamma \begin{bmatrix} p_\pi(s_0 | s_0) & p_\pi(s_1 | s_0) & \dots \\ p_\pi(s_0 | s_1) & p_\pi(s_1 | s_1) & \dots \\ \vdots & \vdots & \ddots \end{bmatrix} \begin{bmatrix} v_\pi(s_0) \\ v_\pi(s_1) \\ \vdots \end{bmatrix}$$

- 13** Consider a policy π that is not optimal, that is $v_\pi(s) < v_{\pi^*}(s), \exists s$. Also consider the greedy-policy operator g that gives a policy gq which is greedy with respect to a given function q , that is, $(gq)(s) = \arg \max_b q(s, b)$, assuming there is no tie. Show that, $v_{gq_\pi}(s) > v_\pi(s), \exists s$ (5 points).

$$\begin{aligned}
v_\pi(s) &= \sum_a \pi(a|s) q_\pi(s, a) \\
v_{gq_\pi}(s) &= \sum_a gq_\pi(a|s) q_\pi(s, a) \\
&= q_\pi(s, \arg \max_b q_\pi(s, b)) \\
&= \max_a q_\pi(s, a) \\
v_{gq_\pi}(s) &= \max_a q_\pi(s, a) \geq v_\pi(s) = \sum_a \pi(a|s) q_\pi(s, a), \forall s
\end{aligned}$$

The condition of equality under all states is that optimal policy, contradict

$$v_{gq_\pi}(s) > v_\pi(s), \exists s$$

- 14** As we have seen, the action value iteration method can be written as: $\mathbf{q}_{k+1} = \mathbf{q}_k - \alpha((\mathbf{I} - \gamma \mathbf{P}_{g\mathbf{q}_k}) \mathbf{q}_k - \mathbf{r})$, where \mathbf{q} is the estimate of action value in vector form, and the vector \mathbf{r} contains $[\mathbf{r}]_{sa} = r(s, a)$ for different s, a . Also, the matrix $\mathbf{P}_{g\mathbf{q}_k}$ contains state-action-transition probabilities, $[\mathbf{P}_{g\mathbf{q}_k}]_{sa, s'a'} = p(s'|s, a)(g\mathbf{q}_k)(a'|s')$, where next actions are taken under policy $g\mathbf{q}_k$, which is the greedy policy with respect to \mathbf{q}_k . Find the condition under which the action-value iteration method converges, and convince that the condition is satisfied in general. (5 points).

- 15** Write the following state-value iteration method in matrix form:
 $v_{k+1}(s) = v_k(s) + \alpha (\max_b [r(s, b) + \gamma \sum_{s'} p(s'|s, b)v_k(s')] - v_k(s))$ (5 points).

- 16** Consider the two-state MDP example from lecture, where $\gamma = 0.9$, reward $r(s, a)$ is 1 for state 1 and action 1, and zero for all other state action pairs, and the state transition probabilities $p(s'|s, a)$ are as follows: $p(0|0, 0) = p(1|0, 1) = p(0|1, 0) = p(1|1, 1) = 1$, and zero for any other transition. (5 points). Now, consider a policy π which takes action 0 at state 0 with probability 0.25 but takes the same action at state 1 with probability 0.75. Find $r_\pi(s) = \sum_a \pi(a|s)r(s, a)$, $p_\pi(s'|s) = \sum_a \pi(a|s)p(s'|s, a)$, and $v_\pi(s)$ for all s and s' . Mention the method you used for finding v_π or show calculation. (5 points).

$$\begin{aligned}
r_\pi(0) &= \pi(0|0) * r(0, 0) + \pi(1|0) * r(0, 1) = 0.25 * 0 + 0.75 * 0 = 0 \\
r_\pi(1) &= \pi(0|1) * r(1, 0) + \pi(1|1) * r(1, 1) = 0.75 * 0 + 0.25 * 1 = 0.25 \\
p_\pi(0|0) &= \pi(0|0) * p(0|0, 0) + \pi(1|0) * p(0|0, 1) = 0.25 * 1 + 0.75 * 0 = 0.25 \\
p_\pi(1|0) &= 1 - p_\pi(0|0) = 0.75 \\
p_\pi(0|1) &= \pi(0|1) * p(0|1, 0) + \pi(1|1) * p(0|1, 1) = 0.75 * 1 + 0.25 * 0 = 0.75 \\
p_\pi(1|1) &= 1 - p_\pi(0|1) = 0.25
\end{aligned}$$

with p and r , we can write v_π in the matrix form $\mathbf{v}_\pi = \mathbf{r}_\pi + \gamma \mathbf{P}_\pi \mathbf{v}_\pi$ and solve the linear system to obtain v_π :

$$\begin{bmatrix} v_\pi(0) \\ v_\pi(1) \end{bmatrix} = \begin{bmatrix} r_\pi(0) \\ r_\pi(1) \end{bmatrix} + \gamma \begin{bmatrix} p_\pi(0|0) & p_\pi(1|0) \\ p_\pi(0|1) & p_\pi(1|1) \end{bmatrix} \begin{bmatrix} v_\pi(0) \\ v_\pi(1) \end{bmatrix}$$

solve $\mathbf{v}_\pi = (\mathbf{I} - \gamma \mathbf{P}_\pi)^{-1} \mathbf{r}_\pi$, we have $v_\pi(0) \approx 1.16, v_\pi(1) \approx 1.34$

- 17** For the same MDP, find q_π , mention the method you used for finding q_π or show calculation, and describe gq_π , that is the greedy policy over q_π for the policy π described in the previous question (5 points).

Similar to Q16, we can obtain the matrix form of q_π and solve the linear system

$$\begin{aligned}
 q_\pi(s, a) &= \sum_{s', r} p(s', r|s, a) \left[r + \gamma \sum_{a'} \pi(a'|s') q_\pi(s', a') \right] \\
 &= \sum_{s', r} p(s', r|s, a) r + \gamma \sum_{s', r} p(s', r|s, a) \sum_{a'} \pi(a'|s') q_\pi(s', a') \\
 &= \sum_r p(r|s, a) r + \gamma \sum_{s'} \sum_{a'} \pi(a'|s') p(s'|s, a) q_\pi(s', a') \\
 &= r(s, a) + \sum_{a', s'} p_\pi(s', a'|s, a) q_\pi(s', a')
 \end{aligned}$$

$$q_\pi = \mathbf{r}_\pi + \gamma \mathbf{P}_\pi q_\pi$$

$$\begin{bmatrix} q_\pi(0, 0) \\ q_\pi(0, 1) \\ \vdots \end{bmatrix} = \begin{bmatrix} r_\pi(0, 0) \\ r_\pi(0, 1) \\ \vdots \end{bmatrix} + \gamma \begin{bmatrix} p_\pi(0, 0|0, 0) & p_\pi(0, 1|0, 0) & \dots \\ p_\pi(0, 0|0, 1) & p_\pi(0, 1|0, 1) & \dots \\ \vdots & \vdots & \ddots \end{bmatrix} \begin{bmatrix} q_\pi(0, 0) \\ q_\pi(0, 1) \\ \vdots \end{bmatrix}$$

fix the matrix

thus $q_\pi = (\mathbf{I} - \gamma \mathbf{P}_\pi)^{-1} + \mathbf{r}_\pi$, the solutions are:

- 18** Consider the nonlinear function approximator for supervised regression $Y \approx \theta^\top \phi(\mathbf{W}\mathbf{x})$, where Y is the target scalar output, \mathbf{x} is the vector input, θ is the output weight vector, \mathbf{W} is the input weight matrix, and ϕ is the element-wise activation function for the hidden layer. Consider the activation function to be sigmoid. Show the backpropagation update for input weights for this case in matrix form. (5 points).

As this is a regression problem, we use Mean Squared Error $(Y - \theta^\top \phi(\mathbf{W}\mathbf{x}))^2$ as the objective:

$$\begin{aligned}
 \theta_{t+1} &= \theta_t - \alpha_t \nabla_{\theta_t} (Y_t - \theta_t^\top \phi(\mathbf{W}_t \mathbf{x}_t))^2 \\
 &= \theta_t + 2\alpha_t (Y_t - \theta_t^\top \phi(\mathbf{W}_t \mathbf{x}_t)) \phi(\mathbf{W}_t \mathbf{x}_t)
 \end{aligned}$$

$$\begin{aligned}
 \mathbf{W}_{t+1} &= \mathbf{W}_t - \alpha_t \nabla_{\mathbf{W}_t} (Y_t - \theta_t^\top \phi(\mathbf{W}_t \mathbf{x}_t))^2 \\
 &= \theta_t + 2\alpha_t (Y_t - \theta_t^\top \phi(\mathbf{W}_t \mathbf{x}_t)) \nabla_{\mathbf{W}_t} \phi(\mathbf{W}_t \mathbf{x}_t) \\
 &= \theta_t + 2\alpha_t (Y_t - \theta_t^\top \phi(\mathbf{W}_t \mathbf{x}_t)) \phi(\mathbf{W}_t \mathbf{x}_t) \circ (1 - \phi(\mathbf{W}_t \mathbf{x}_t)) \nabla_{\mathbf{W}_t} \mathbf{W}_t \mathbf{x}_t \\
 &= \theta_t + 2\alpha_t (Y_t - \theta_t^\top \phi(\mathbf{W}_t \mathbf{x}_t)) \phi(\mathbf{W}_t \mathbf{x}_t) \circ (1 - \phi(\mathbf{W}_t \mathbf{x}_t)) \mathbf{x}_t^\top
 \end{aligned}$$

where \circ indicates elementwise multiplication