Summary

Report ID:	ac089661-c438-4f52-9a70-e8120ad7d362	**Generated (UTC):**	2026-02-22T02:39:08.326255Z
Document Type:	AI Summary	**Reviewer:**	MedDecode AI (Automated)

Report ID: ac089661-c438-4f52-9a70-e8120ad7d362

Generated: 2026-02-22T02:39:08.326255Z

Most likely possibilities:

- Oblique fracture of the proximal phalanx of the 4th digit (ring finger) — Likelihood: High
- Possible soft tissue swelling associated with the fracture — Likelihood: Medium
- Possible occult/non-displaced fracture in other hand bones cannot be excluded — Likelihood: Low

Supporting visual cues:

- A clear oblique fracture line is visible in the mid-shaft of the proximal phalanx of the 4th digit.
- There is cortical disruption and mild angulation of the fracture fragments.
- No other obvious acute fractures or dislocations are seen in the visualized bones of the hand.

What would confirm:

- Orthogonal X-ray views (e.g., lateral, oblique) of the hand and specifically the 4th digit.
- Correlation with clinical examination findings (pain, swelling, deformity).
- A formal radiology report.

Limits: Single X-ray view, no clinical history provided.

Next step: Obtain additional X-ray views and clinical correlation.

CONFIDENTIAL

AI-generated summary for informational use only. Confirm with a clinician.