Summary

Report ID:	ce837662-b693-45b9-81ed-7d6101a91acf	**Generated (UTC):**	2026-02-22T03:27:03.547140Z
Filename:	broken finger xray.pdf	**Reviewer:**	MedDecode AI (Automated)

Report ID: ce837662-b693-45b9-81ed-7d6101a91acf

Filename: broken finger xray.pdf

Generated: 2026-02-22T03:27:03.547140Z

IMPRESSION:

Possible oblique fracture of the proximal phalanx of the second digit (index finger).

Confidence: High

FINDINGS:

- An oblique lucent fracture line is identified through the mid-shaft of the proximal phalanx of the second digit.
- Mild angulation and subtle displacement of the fracture fragments are noted at the fracture site.
- The remaining metacarpals and phalanges appear intact.
- Carpal bones demonstrate preserved alignment and no obvious acute abnormalities.
- No other obvious acute osseous pathology is seen.

WHAT THIS MEANS (plain language):

- It appears there is a break in the bone of your index finger, specifically in the first bone closest to your hand.
- The break runs diagonally across the middle part of this bone.
- The pieces of the bone might be slightly angled or shifted.
- This finding will need further medical assessment and guidance.

LIMITATIONS:

- Only a single radiographic view (PA) of the hand is provided, which may not fully demonstrate the extent of displacement or angulation.
- No clinical history or patient symptoms were available for correlation.
- Soft tissue assessment is limited on this single projection.

CONFIDENTIAL

AI-generated summary for informational use only. Confirm with a clinician.

RECOMMENDED NEXT STEP:

- Correlation with clinical findings and patient symptoms is essential.
- Obtain additional radiographic views (e.g., lateral and oblique views) of the hand/digit for complete fracture characterization.
- Clinical evaluation by an orthopedic specialist or emergency department physician for definitive diagnosis and management.
- Consider temporary immobilization of the affected digit.

CONFIDENTIAL

AI-generated summary for informational use only. Confirm with a clinician.