

**STAT 510 Homework 1****Due Date:** 11:00 A.M., Wednesday, January 23

1. Consider the matrix

$$\mathbf{X} = \begin{bmatrix} 1 & 7 & 0 & -3 \\ 0 & -2 & 1 & 2 \\ -1 & -5 & -1 & 1 \end{bmatrix}.$$

- Show that the columns of  $\mathbf{X}$  are linearly dependent.
  - Find the rank of  $\mathbf{X}$ .
  - Use the generalized inverse algorithm in slide set 1 to find a generalized inverse of  $\mathbf{X}$ .
  - Use the R function `ginv` in the MASS package to find a generalized inverse of  $\mathbf{X}$ . (To load the MASS package into your R workspace use the command `library(MASS)`. If the MASS package is not already installed, you will need to install it before loading. The command `install.packages('MASS')` should install the MASS package if necessary.)
  - Provide one matrix  $\mathbf{X}^*$  that satisfies both of the following characteristics:
    - $\mathbf{X}^*$  has full-column rank (i.e.,  $\text{rank}(\mathbf{X}^*)$  is equal to the number of columns of  $\mathbf{X}^*$ ), and
    - $\mathbf{X}^*$  has column space equal to the column space of  $\mathbf{X}$ ; i.e.,  $\mathcal{C}(\mathbf{X}^*) = \mathcal{C}(\mathbf{X})$ .
  - What property do all the three-dimensional vectors in the column space of  $\mathbf{X}$  share?
- Prove that any set of  $n$ -dimensional vectors that contains the  $n$ -dimensional zero vector is linearly dependent.
  - Imagine extending a string from  $(0, 0)$ , the origin in  $\mathbb{R}^2$ , to a random point  $(x, y)$  in  $\mathbb{R}^2$ , where  $x \sim N(2, 1)$  independent of  $y \sim N(1, 1)$ . Use R to find the probability that the string will need to be longer than 6 units to reach from  $(0, 0)$  to  $(x, y)$ .
  - Suppose  $z_1, z_2 \stackrel{iid}{\sim} N(0, 1)$ . Find the distribution of the following random variables and prove that your answer is correct.
    - $(z_1 - z_2)^2/2$
    - $(z_1 + z_2)/|z_1 - z_2|$
  - Suppose  $y_1, \dots, y_n \stackrel{iid}{\sim} N(\mu, \sigma^2)$ . Let  $\mathbf{y} = [y_1, \dots, y_n]'$ , and let  $\bar{y} = \frac{1}{n} \sum_{i=1}^n y_i$ .
    - Show that  $s^2 = \frac{1}{n-1} \sum_{i=1}^n (y_i - \bar{y})^2$  can be written as  $\mathbf{y}'\mathbf{B}\mathbf{y}$  for some matrix  $\mathbf{B}$ .
    - Prove that  $(n-1)s^2/\sigma^2 \sim \chi_{n-1}^2$  using the “Important Distributional Result about Quadratic Forms” in slide set 1.
  - Prove that a matrix  $\mathbf{A}$  is  $\mathbf{0}$  if and only if  $\mathbf{A}'\mathbf{A} = \mathbf{0}$ . (Hint: What are the diagonal elements of  $\mathbf{A}'\mathbf{A}$ ?)
  - Prove that  $\mathbf{X}'\mathbf{X}\mathbf{A} = \mathbf{X}'\mathbf{X}\mathbf{B}$  if and only if  $\mathbf{X}\mathbf{A} = \mathbf{X}\mathbf{B}$ . Note that the “if” part of the proof, i.e.,

$$\mathbf{X}\mathbf{A} = \mathbf{X}\mathbf{B} \implies \mathbf{X}'\mathbf{X}\mathbf{A} = \mathbf{X}'\mathbf{X}\mathbf{B},$$

holds trivially. Thus, proving the converse, i.e.,

$$\mathbf{X}'\mathbf{X}\mathbf{A} = \mathbf{X}'\mathbf{X}\mathbf{B} \implies \mathbf{X}\mathbf{A} = \mathbf{X}\mathbf{B},$$

is the challenging part. One proof makes use of the result in problem 6.