

# COMM8102 ECONOMETRIC ANALYSIS

## ASSIGNMENT 3

YUHAO LIU

z5097536

July 17, 2021

### 1 Exercise 7.1

$$\begin{aligned}\hat{\beta}_1 &= \left( \frac{1}{n} \sum_{i=1}^n X_{1i} X'_{1i} \right)^{-1} \left( \frac{1}{n} \sum_{i=1}^n X_{1i} Y_i \right) \\&= \left( \frac{1}{n} \sum_{i=1}^n X_{1i} X'_{1i} \right)^{-1} \left( \frac{1}{n} \sum_{i=1}^n X_{1i} (X'_{1i} \beta_1 + X'_{2i} \beta_2 + e_i) \right) \\&= \left( \frac{1}{n} \sum_{i=1}^n X_{1i} X'_{1i} \right)^{-1} \left( \beta_1 \frac{1}{n} \sum_{i=1}^n X_{1i} X'_{1i} + \beta_2 \frac{1}{n} \sum_{i=1}^n X_{1i} X'_{2i} + \frac{1}{n} \sum_{i=1}^n X_{1i} e_i \right) \\&\quad \xrightarrow{P} E[X_1 X'_1]^{-1} (E[X_1 X'_1] \beta_1 + E[X_1 X'_2] \beta_2 + E[X_1 e]) \\&= \beta_1 + E[X_1 X'_1]^{-1} E[X_1 X'_2] \beta_2 + E[X_1 X'_1]^{-1} E[X_1 e] \\&= \beta_1 + E[X_1 X'_1]^{-1} E[X_1 X'_2] \beta_2 + 0 \\&= \beta_1 + E[X_1 X'_1]^{-1} E[X_1 X'_2] \beta_2\end{aligned}$$

by WLLN and CMT.  $E[X_1 e] = 0$ , since  $E[Xe] = 0$ . Since  $\hat{\beta}_1 \xrightarrow{P} \beta_1 + E[X_1 X'_1]^{-1} E[X_1 X'_2] \beta_2$ , it is not consistent for  $\beta_1$ . When  $E[X_1 X'_2] = \mathbf{0}$  or  $\beta_2 = \mathbf{0}$ ,  $\hat{\beta}_1 \xrightarrow{P} \beta_1$ .

## 2 Exercise 7.15

$$\begin{aligned}
\sqrt{n}(\hat{\beta} - \beta) &= \sqrt{n} \left( \frac{\sum_{i=1}^n X_i^3 Y_i}{\sum_{i=1}^n X_i^4} - \beta \right) \\
&= \frac{\sqrt{n} \frac{1}{n} \sum_{i=1}^n X_i^3 Y_i}{\frac{1}{n} \sum_{i=1}^n X_i^4} - \sqrt{n} \beta \\
&= \frac{\sqrt{n} \frac{1}{n} \sum_{i=1}^n X_i^3 (X_i \beta + e_i)}{\frac{1}{n} \sum_{i=1}^n X_i^4} - \sqrt{n} \beta \\
&= \frac{\sqrt{n} \frac{1}{n} \sum_{i=1}^n X_i^4 \beta}{\frac{1}{n} \sum_{i=1}^n X_i^4} + \frac{\sqrt{n} \frac{1}{n} \sum_{i=1}^n X_i^3 e_i}{\frac{1}{n} \sum_{i=1}^n X_i^4} - \sqrt{n} \beta \\
&= \underbrace{\frac{\beta \sqrt{n} \frac{1}{n} \sum_{i=1}^n X_i^4}{\frac{1}{n} \sum_{i=1}^n X_i^4} - \sqrt{n} \beta}_0 + \frac{\sqrt{n} \frac{1}{n} \sum_{i=1}^n X_i^3 e_i}{\frac{1}{n} \sum_{i=1}^n X_i^4} \\
&= \frac{\sqrt{n} \left( \frac{1}{n} \sum_{i=1}^n \left( X_i^3 e_i - \underbrace{E[X_i^3 e]}_0 \right) \right)}{\frac{1}{n} \sum_{i=1}^n X_i^4} \\
&\xrightarrow{d} \frac{\mathcal{N}(0, E[X_i^6 e_i^2])}{E[X_i^4]} \\
&= \mathcal{N}\left(0, \frac{E[X_i^6 e_i^2]}{(E[X_i^4])^2}\right)
\end{aligned}$$

## 3 Exercise 9.4

### 3.1 (a)

The size of a test is the type I error.

$$\begin{aligned}
P(\text{Reject } H_0 | H_0 \text{ is true}) &= P(\text{Reject } H_0 | H_0) \\
&= P(W < c_1 \text{ or } W > c_2 | H_0)
\end{aligned}$$

Under  $H_0$ ,  $W \xrightarrow{d} \chi_q^2$ ,  $P(W < c_1 \text{ or } W > c_2 | H_0) = P(W < c_1 | H_0) + P(W > c_2 | H_0) = \frac{\alpha}{2} + 1 - (1 - \frac{\alpha}{2}) = \alpha$ .

### 3.2 (b)

This is not a good test of  $H_0$  vs  $H_1$ . Wald test is the extension of t-test to multivariate case. Wald test statistics is the squared version of t statistics in the scalar case. Only the right tail captures the extreme deviation from 0 (both positive and negative). We do not need to look at the left tail of the  $\chi_q^2$  distribution.

## 4 Exercise 9.12

This interpretation is not correct.

The power of a hypothesis test is the probability that reject  $H_0$  when the  $H_1$  is true.  $p$  value is a quantity depending on the value of test statistics and the asymptotic null distribution  $G$ . The power depends on the critical value and the distribution of test statistic under the alternative hypothesis.

Although a larger sample will make the test have more power, it does not necessarily reduce the  $p$  value.  $p$  value depends on both of the test statistic and the asymptotic null distribution  $G$ . The extra samples can

make the value of test statistic increase, but it can also make the value of test statistic decrease. Therefore, it cannot be guaranteed that a larger sample will make the test reject the null.

## 5 Exercise 9.16

### 5.1 (a)

$$\hat{\theta} = \frac{1}{n} \left( \sum_{i=1}^n e_{1i}^2 - e_{2i}^2 \right)$$

### 5.2 (b)

By central limit theorem,

$$\left( \begin{array}{c} \sqrt{n} \left( \frac{1}{n} \sum_{i=1}^n e_{1i}^2 - \sigma_1^2 \right) \\ \sqrt{n} \left( \frac{1}{n} \sum_{i=1}^n e_{2i}^2 - \sigma_2^2 \right) \end{array} \right) \xrightarrow{d} \mathcal{N} \left( \left( \begin{array}{c} 0 \\ 0 \end{array} \right), V \right),$$

where

$$V = \left( \begin{array}{cc} E \left[ (e_1^2 - \sigma_1^2)^2 \right] & E \left[ (e_1^2 - \sigma_1^2) (e_2^2 - \sigma_2^2) \right] \\ E \left[ (e_1^2 - \sigma_1^2) (e_2^2 - \sigma_2^2) \right] & E \left[ (e_2^2 - \sigma_2^2)^2 \right] \end{array} \right).$$

Define a function  $r : \mathbb{R}^2 \rightarrow \mathbb{R}$ ,  $(x, y) \mapsto x - y$ . Then we have  $\theta = r(\sigma_1^2, \sigma_2^2)$ ,  $\hat{\theta} = r\left(\frac{1}{n} \sum_{i=1}^n e_{1i}^2, \frac{1}{n} \sum_{i=1}^n e_{2i}^2\right)$ . By delta method,

$$\begin{aligned} \sqrt{n}(\hat{\theta} - \theta) &= \sqrt{n} \left( r \left( \frac{1}{n} \sum_{i=1}^n e_{1i}^2, \frac{1}{n} \sum_{i=1}^n e_{2i}^2 \right) - r(\sigma_1^2, \sigma_2^2) \right) \\ &\xrightarrow{d} \mathcal{N}(0, R' V R), \end{aligned}$$

where

$$\begin{aligned} R &= \left( \begin{array}{c} \frac{\partial}{\partial \sigma_1^2} r(\sigma_1^2, \sigma_2^2) \\ \frac{\partial}{\partial \sigma_2^2} r(\sigma_1^2, \sigma_2^2) \end{array} \right) \\ &= \left( \begin{array}{c} 1 \\ -1 \end{array} \right). \end{aligned}$$

Finally, we have

$$\sqrt{n}(\hat{\theta} - \theta) \xrightarrow{d} \mathcal{N}(0, V_\theta),$$

with

$$\begin{aligned} V_\theta &= E \left[ (e_1^2 - \sigma_1^2)^2 \right] - 2 \cdot E \left[ (e_1^2 - \sigma_1^2) (e_2^2 - \sigma_2^2) \right] + E \left[ (e_2^2 - \sigma_2^2)^2 \right] \\ &= E \left[ ((e_1^2 - \sigma_1^2) - (e_2^2 - \sigma_2^2))^2 \right] \\ &= E \left[ (e_1^2 - e_2^2 - (\sigma_1^2 - \sigma_2^2))^2 \right] \end{aligned}$$

### 5.3 (c)

Since  $\hat{\theta} = \theta + \frac{\sqrt{n}(\hat{\theta} - \theta)}{\sqrt{n}}$ , and  $\sqrt{n}(\hat{\theta} - \theta) \xrightarrow{d} \mathcal{N}(0, V_\theta)$ ,

$$\hat{\theta} \xrightarrow{d} \mathcal{N}\left(\theta, \frac{V_\theta}{n}\right)$$

An estimator of the asymptotic variance of  $\hat{\theta}$  is

$$\hat{V}_{\hat{\theta}} = \frac{1}{n} \left( \frac{1}{n} \sum_{i=1}^n (e_{1i}^2 - e_{2i}^2 - (\sigma_1^2 - \sigma_2^2))^2 \right).$$

### 5.4 (d)

$H_0$  is equivalent to  $\theta = 0$ , and  $H_1$  is equivalent to  $\theta \neq 0$ . Under  $H_0$  the test statistic is

$$\begin{aligned} T &= \frac{\hat{\theta} - 0}{\sqrt{\frac{V_\theta}{n}}} \\ &\sim \mathcal{N}(0, 1). \end{aligned}$$

A test of asymptotic size  $\alpha$  can be, we reject the null hypothesis if  $T < a_1$  or  $T > a_2$ , where  $a_1$  is the  $\frac{\alpha}{2}$  quantile of  $\mathcal{N}(0, 1)$  and  $a_2$  is the  $1 - \frac{\alpha}{2}$  of  $\mathcal{N}(0, 1)$ .

### 5.5 (e)

Acceptance of the null hypothesis means these two models have same variance for the residuals, therefore they fit the data equally well.

## 6 Exercise 9.26

### 6.1 (a)

	Estimations	Standard errors HC1
$\hat{\beta}_1$	-3.527	1.7186
$\hat{\beta}_2$	0.720	0.0326
$\hat{\beta}_3$	0.436	0.2456
$\hat{\beta}_4$	-0.220	0.3238
$\hat{\beta}_5$	0.427	0.0755

### 6.2 (b)

$$\begin{aligned} \log C &= \beta_1 + \beta_2 \log Q + \beta_3 \log PL + \beta_4 \log PK + \beta_5 \log PF + e \\ &= \beta_1 + \log Q^{\beta_2} + \log PL^{\beta_3} + \log PK^{\beta_4} + \log PF^{\beta_5} + e \end{aligned}$$

Then we have

$$C = \exp(\beta_1 + e) \cdot Q^{\beta_2} \cdot PL^{\beta_3} \cdot PK^{\beta_4} \cdot PF^{\beta_5}$$

For  $\beta_3 + \beta_4 + \beta_5 = 1$ , this means the cost function has a constant return to scale for labor, capital and fuel.

### 6.3 (e)

Define a function  $r : \mathbb{R}^5 \rightarrow \mathbb{R}$ , such that  $r(\beta_1, \beta_2, \beta_3, \beta_4, \beta_5) = \beta_3 + \beta_4 + \beta_5$ . Then we have  $\nabla r' = [0, 0, 1, 1, 1]'$ . By Delta method and affine transform of normal distribution, we establish

$$\sqrt{n} \left( \hat{\beta}_3 + \hat{\beta}_4 + \hat{\beta}_5 - (\beta_3 + \beta_4 + \beta_5) \right) \xrightarrow{d} \mathcal{N}(0, V)$$

where  $V = \nabla r' \mathbf{V}_\beta \nabla r$ .  $\mathbf{V}_\beta$  is the asymptotic covariance matrix of  $\sqrt{n}(\hat{\beta} - \beta)$ . Then the Wald statistic is

$$\begin{aligned} \mathcal{W}_n &= \frac{\left( \hat{\beta}_3 + \hat{\beta}_4 + \hat{\beta}_5 - (\beta_3 + \beta_4 + \beta_5) \right)^2}{\hat{V}} \\ &= 0.645. \end{aligned}$$

Since  $\mathcal{W}_n \xrightarrow{d} \chi_1^2$ , the critical value  $c = 3.84$  for  $\alpha = 0.05$ . We reject the null hypothesis.