


Gap structure of 1D cut and project Hamiltonians

Nicolas Macé, Anuradha Jagannathan, Frédéric Piéchon

Laboratoire de Physique des Solides
Université Paris-Saclay

September 20, 2016



OUTLINE

1 The gap labeling theorem


2 The Fibonacci chain

3 General case

4 Conclusion

ELECTRONS ON QUASIPERIODIC CHAINS

Canonical cut-and-project method



Quasiperiodic chain:

$$\alpha \in \mathbb{R} \setminus \mathbb{Q}$$

Approximants:

$$\alpha_l = \frac{p_l}{q_l} \in \mathbb{Q}$$

$$\alpha_l \rightarrow \alpha$$


Hamiltonian: $H(\alpha) = \sum_{x \in \mathbb{Z}} t_{x,x+1} |x\rangle \langle x+1| + \text{h.c.}$

2 lengths \rightarrow 2 jump amplitudes $t_{x,x+1} = t_1$ or t_2 .

THE ELECTRONIC SPECTRUM

A convenient way to plot the spectrum: the integrated density of states (idos).

$\text{idos}(E)$ = fraction of states below energy E



idos of the Fibonacci Hamiltonian

- Electronic spectrum of quasiperiodic chains is hard to describe
- Rather: describe the idos in the gaps \rightarrow **gap labeling theorem**

$$\text{idos}(E \in \text{gap}) = n\tau^{-2} \mod 1$$

THE GAP LABELING THEOREM

The IDOS inside spectral gaps can be indexed using the irrational involved in the construction of the chain

$$\text{idos}(E \in \text{gap}) = \frac{n}{1 + \alpha} \mod 1$$

- Constrains the spectrum... but is not enough to reconstruct it
- Model independent! (while the spectrum is model dependent)
→ n is a topological invariant

THE GAP LABELING THEOREM

$$\text{idos}(E \in \text{gap}) = \frac{n}{1 + \alpha} \mod 1$$

- Has the gap label n a physical interpretation?
- Can the theorem be applied to approximants?
- Does it help understanding the quasiperiodic limit?

A SIMPLE, BUT INCORRECT PROOF

Let $\alpha_l = \frac{p_l}{q_l} \rightarrow \alpha$ be a sequence of approximants.

$N_l = p_l + q_l$ is the maximum number of energy bands.

$$\text{idos}(E \in \text{gap}) = \frac{j(E)}{N_l}$$

We can find integers n, k such that $j = nq_l + kN_l$.

$$\text{idos}(E \in \text{gap}) = \frac{nq_l}{p_l + q_l} \mod 1$$

Letting $l \rightarrow \infty$,

$$\text{idos}(E \in \text{gap}) = \frac{n}{1 + \alpha} \mod 1$$

A SIMPLE, BUT INCORRECT PROOF

Let $\alpha_l = \frac{p_l}{q_l} \rightarrow \alpha$ be a sequence of approximants.

$N_l = p_l + q_l$ is the maximum number of energy bands.

$$\text{id}os(E \in \text{gap}) = \frac{j(E)}{N_l}$$

We can find integers n, k such that $j = nq_l + kN_l$.

$$\text{id}os(E \in \text{gap}) = \frac{nq_l}{p_l + q_l} \mod 1$$

Letting $l \rightarrow \infty$,

$$\text{id}os(E \in \text{gap}) = \frac{n}{1 + \alpha} \mod 1$$

Problem

n may depend on l .

A CONCRETE EXAMPLE

Gaps of successive approximants of the Fibonacci chain.



$\langle E \rangle_l$: mean energy of a gap, $\Delta_l(\langle E \rangle)$: its width.

Identify two gaps if they overlap:

$$0.5\Delta_l(\langle E \rangle) > |\langle E \rangle_l - \langle E' \rangle_{l+1}|$$

A CONCRETE EXAMPLE

Gaps of successive approximants of the Fibonacci chain.



- Blue labeled gaps have a stable label gap label → stable gaps.
- Red labeled do not → transient gaps.

A CONCRETE EXAMPLE

Gaps of successive approximants of the Fibonacci chain.



- **Transient** gaps closes as $l \rightarrow \infty$
- **Stable** gaps stay open.

A CONCRETE EXAMPLE

Gaps of successive approximants of the Fibonacci chain.



- **Stable gaps:** n independent of $l \rightarrow$ naive proof works.
- **Transient gaps:** n depends on l , but these gaps disappear.

\rightarrow the naive proof works and correctly labels the gaps.

RECURSIVE GAP LABELING

...translates into recursive construction of the gaps!

Recursive construction of the spectrum...



$$G_l^{\text{left}} = M^{-2} G_{l-2}$$

$$G_l^0 = M^{-3} G_{l-3} + \mathbf{g}_1$$

$$G_l^{\text{right}} = M^{-2} G_{l-2} + \mathbf{g}_2$$

Where G_l is the set of labels:


$$G_l = \{(m, n) | \text{id}os = n / (1 + \alpha) + m\}$$

$$M = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$$

- **Stable gaps** are the iterates of the 2 main gaps
- **Transient gaps** are the iterates of the central gap.

GENERAL CASE

We plot the gapwidth Δ_l as a function of the label for various quasiperiodic chains:



- The width decreases as a power-law of the label
- Above a critical label, all gaps are transient
- Recursive gap labeling using the Hofstadter rules? [Rüdinger, Piéchon 98]

CONCLUSION AND PERSPECTIVES

- The gap labeling theorem can be extended to approximants
- The price to pay is the introduction of transient gaps, absent in the quasiperiodic case
- Gap labels have a physical meaning:
 - It orders gap by decreasing width
 - It separates stable from transient gaps
 - It can be interpreted as a Chern number [Levy *et al* 2015]

Perspectives:

- Understand the gap width behavior with the gap label
- Investigate to 2D quasicrystals, which also have gaps [Prunelé *et al* 2002]

THE GAP LABELING THEOREM: PRECISE STATEMENT

Let w be a cut-and-project word. Consider the Hamiltonian:

$$H(w) = \sum_{x,y} t(T^{-y}w, x - y) |x\rangle \langle y|$$

Interactions must be local:

$$\sup_{u \in \text{Hull}(w)} \sum_x |t(u, x)| < \infty$$

Gap labeling theorem:

The idos in a gap is a linear combination of frequencies of local environments of w .

taken from *The non-commutative geometry of aperiodic solids*, Bellissard
2003.

CASES WHERE THE NAIVE PROOF FAILS

Consider approximants to the Fibonacci chain. Gaps are labeled by

$$\text{idos}(n) = \frac{n}{1 + \alpha_l} \mod 1$$

Consider the sequence of gap labels

$$n_{l=3k} = \left\lfloor \frac{(2 + \sqrt{5})^k}{2\sqrt{5}} \right\rfloor$$


We have $\text{idos}(n_l) = 1/2$ (it labels the $E = 0$ gap). Taking $l \rightarrow \infty$, we could – incorrectly – conclude that $1/2$ is a gap.

However, there is no finite n such that

$$\frac{1}{2} = n\tau^{-2} \mod 1$$

CONUMBERING AND GAP LABELING

Conumbering: labeling of the atoms according to their internal space position.





see Mosseri & Sire 1990.

$$\text{idos} = \frac{j}{N_l}$$

← conumbering
normal numbering →

$$\text{idos} = \frac{n}{1 + \alpha_l} \mod 1$$

CONUMBERING AND GAP LABELING



Plotting the local density of states makes the symmetry between gap labels and atomic labels evident.