

Deep learning versus manual morphology-based embryo selection in IVF: a randomized, double-blind noninferiority trial

Received: 25 January 2024

Accepted: 29 June 2024

Published online: 9 August 2024

 Check for updates

A list of authors and their affiliations appears at the end of the paper

To assess the value of deep learning in selecting the optimal embryo for **in vitro fertilization, a multicenter, randomized, double-blind, noninferiority** parallel-group trial was conducted across 14 in vitro fertilization clinics in Australia and Europe. Women under 42 years of age with at least two early-stage blastocysts on day 5 were randomized to either the control arm, using standard morphological assessment, or the study arm, employing a deep learning algorithm, intelligent Data Analysis Score (iDAScore), for **embryo selection. The primary endpoint was a clinical pregnancy rate with a noninferiority margin of 5%. The trial included 1,066 patients (533 in the iDAScore group and 533 in the morphology group). The iDAScore group exhibited a clinical pregnancy rate of 46.5% (248 of 533 patients), compared to 48.2% (257 of 533 patients) in the morphology arm (risk difference -1.7%; 95% confidence interval -7.7, 4.3; $P = 0.62$). This study was not able to demonstrate noninferiority of deep learning for clinical pregnancy rate when compared to standard morphology and a predefined prioritization scheme. Australian New Zealand Clinical Trials Registry (ANZCTR) registration: [379161](#).**

Artificial intelligence (AI), specifically deep learning, has been heralded as a revolutionary technology, with its potential in healthcare, particularly in relation to the interpretation and reporting of medical images, widely acknowledged¹. Despite its potential, the clinical utility of AI in healthcare remains underexplored in randomized controlled trials (RCTs)².

During in vitro fertilization (IVF), a key challenge is selecting the embryo that will give the patient the best chance of a live birth from a group of usable embryos. Currently, during the culture period, the embryologist removes the embryos from the incubator, usually more than once, and performs a 'snapshot' morphological assessment under a microscope³. This approach is time consuming, subjective and has, in essence, changed little since the first IVF birth 45 years ago^{4,5}.

The advent of time-lapse incubators, capable of capturing frequent embryo images over five to six days of growth, has circumvented

the need to remove embryos from the incubator for assessment, minimizing disturbance⁶. Combining this with a validated automated selection algorithm has the additional potential to substantially improve workflow efficiency compared to standard incubation and to overcome subjectivity⁷. To date, several studies have employed deep learning algorithms for embryo quality grading or development stage classification based on static images from time-lapse incubators^{8–11}. Recently, the deep learning algorithm iDAScore was developed that uses spatial (morphological) and temporal (morphokinetic) patterns from time-lapse images across the first six days of embryological development (to the blastocyst stage) to predict, with high discriminative ability and reproducibility, the probability of a transferred embryo implanting and progressing to a clinical pregnancy^{12,13}.

Despite numerous proposed applications of AI in IVF, their integration into routine clinical practice hinges on robust evidence

✉ e-mail: peter.illingworth@virtushealth.com.au




Fig. 1 | Patient flow diagram. The number of patients randomized in the electronic case report form (eCRF) to the study group and the control group then, following exclusions for discontinued intervention, the number analyzed per protocol.

demonstrating their efficacy compared to traditional methods¹⁴. A number of studies of AI have suggested significant improvements in consistency of embryo scoring^{15,16}, but these evaluations have been retrospective. Two previous studies^{17,18} have prospectively examined the efficacy of traditional machine learning algorithms based on previous data and human input for embryo selection. By contrast, deep learning is a more advanced form of machine learning that enables computers to learn from experience and understand the world in terms of a hierarchy of concepts¹⁹. Because the computer gathers knowledge from experience, there is, unlike conventional machine learning, no need for a human operator to formally specify any of the knowledge needed by the computer. No RCTs of deep learning in embryo selection have previously been reported. Despite the potential utility of deep learning, there is a need to demonstrate noninferiority to standard embryologist assessment of the embryos.

We therefore undertook a prospective randomized noninferiority trial to determine whether selection of a single blastocyst for transfer by deep learning results in a clinical pregnancy rate comparable with a 5% inferiority margin to that achieved by trained embryologists using standard morphology criteria.

Results

From 16 January 2020 until 30 September 2022, a total of 1,751 participants were eligible, with 1,066 patients included in the trial (533 in the study arm and 533 in the control arm). Of these, 1,002 participants were included in the per-protocol (PP) analysis after the exclusion of 64 participants, owing to protocol violations (Patient flow diagram; Fig. 1). The demographics and clinical characteristics of the patients at the time of randomization were well balanced between the trial groups, except from the type of transfer, with an extra 7.5% of patients undergoing a frozen–thawed embryo transfer in the study group compared to the control group (Table 1).

Primary outcome

The primary outcome, clinical pregnancy, occurred in 248 of 533 patients (46.5%) in the study group and in 257 of 533 patients (48.2%) in the control group. Noninferiority of embryo selection using the deep learning algorithm was not shown, with an absolute risk difference of −1.7 percentage points (95% confidence interval (CI), −7.7, 4.3) and a rate ratio of 0.96 (95% CI, 0.85, 1.10) (Table 2 and Fig. 2a). The results were similar for the PP analysis, full analysis set (FAS) and the primary efficacy analysis with adjustment for center and selected allocation variables (Table 2 and Supplementary Tables 1 and 2).

Secondary outcomes

Results similar to the primary outcome were observed for the secondary outcomes of the positive human chorionic gonadotropin (hCG) rate, nonviable intrauterine pregnancies and ongoing pregnancy rate (Supplementary Tables 3 and 4). Live birth rate was similar between the two groups where the live birth rate for the iDAScore group was 39.8% (212 of 533 patients) and 43.5% (232 of 533 patients) in the standard morphology criteria group (risk difference −3.9%; 95% CI, −9.9, 2.2; $P = 0.24$) (Supplementary Tables 3 and 4).

The study group was found to have an almost 10-fold reduction in the time required for evaluation, with a mean standard deviation time of 21.3 ± 18.1 seconds in comparison to the control group, which took 208.3 ± 144.7 seconds ($P < 0.001$) regardless of the number of embryos available on day 5 (Fig. 3).

Safety

No medical adverse events were recorded, while adverse device effects were reviewed and recorded as protocol breaches.

Sensitivity analyses

A prespecified sensitivity analysis, by adjusting for each woman's age, the number of previous stimulated IVF cycles leading to oocyte pick-up,

Table 1 | Baseline and treatment characteristics (intention-to-treat population)

Variable	Study group (n=533)	Control (n=533)
Maternal age	33.7 (3.7) 34 (22; 43)	33.8 (3.8) 34 (20; 42)
Reason for infertility (couple)		
No clinical subfertility	44 (8.3%)	42 (7.9%)
Unexplained infertility	206 (38.6%)	202 (37.9%)
Known cause for infertility	283 (53.1%)	289 (54.2%)
Maternal body mass index	25.4 (4.9) 24.3 (15.1; 44.6)	25.5 (5.0) 24.4 (15.6; 48.9)
Number of previous stimulated IVF cycles leading to oocyte pick-up	0.533 (1.008) 0 (0; 9)	0.606 (1.103) 0 (0; 11)
Subjects with a previous pregnancy in current relationship	146 (27.4%)	155 (29.1%)
FSH treatment total dosage	2,426 (1206) 2,163 (56; 7,500)	2,393 (1231) 2,200 (77; 6,750)
GnRH downregulation		
Agonist	65 (12.2%)	65 (12.2%)
Antagonist	468 (87.8%)	468 (87.8%)
Number of oocytes	14.8 (8.0) 13 (2; 52)	14.7 (8.6) 13 (2; 76)
Method of fertilization		
Standard IVF	226 (42.4%)	227 (42.6%)
ICSI	278 (52.2%)	276 (51.8%)
Combined	29 (5.4%)	30 (5.6%)
Number of normally fertilized oocytes (2PN)	9.02 (5.20) 8 (1; 34)	9.01 (5.35) 8 (1; 44)
Number of blastocysts at Gardner grade 2 or beyond by day 5	5.26 (3.70) 4 (0; 28)	5.03 (3.53) 4 (0; 29)
Number of cryopreserved embryos on day 5 and 6	4.80 (3.86) 4 (0; 28)	4.08 (3.72) 3 (0; 31)
Gardner grade on day 5 of the transferred embryo ^a		
1 or 2	21 (4.0%)	15 (2.8%)
3AA, 4AA, 5AA, 6AA	320 (61.1%)	361 (68.5%)
Others	183 (34.9%)	151 (28.7%)
iDAScore of the transferred embryo	9.09 (0.68) 9.3 (4.7; 9.9)	
Type of transfer		
Fresh (current randomized cycle)	320 (60.3%)	353 (66.4%)
After cryopreservation or total freeze	211 (39.7%)	179 (33.6%)

For categorical variables *n* (%) is presented. For continuous variables mean (s.d.)/median (minimum; maximum) is presented. ^aFreeze-all embryos were selected on information up to day 6, which, in some cases, led to an embryo graded 1 on day 5 being prioritized. FSH, follicle-stimulating hormone; GnRH, gonadotropin-releasing hormone; ICSI, intracytoplasmic sperm injection.

the number of oocytes obtained, the fertilization method used, the number of normally fertilized oocytes and the center, did not materially change the results obtained (Table 2).

Assessment of the efficacy of the deep learning algorithm compared to standard morphology criteria in each of the 14 study centers showed broadly similar outcomes, with risk ratios (RR) for clinical pregnancy ranging from 0.23 to 2.08, with all centers having wide CIs that included unity (Fig. 2b).

In the prespecified subgroup analysis where the results in only women >35 years of age were compared, clinical pregnancy occurred

in 85 of 228 (37.3%) in the study group and in 89 of 228 (39.0%) in the control group, with a risk difference between the groups of −1.8 (95% CI, −11.1, 7.6). The positive hCG rate, the nonviable intrauterine pregnancy, ongoing pregnancy rate and live birth rate were also similar between the study and control groups (Supplementary Table 5). In a further prespecified interaction analysis, we observed differing performance between the study and control groups in fresh-embryo transfer and freeze-all cycles (RR, 1.08 versus RR, 0.81; *P* = 0.022).

To assess the moderating potential of variables including maternal age and the number of available blastocysts, prespecified interaction analyses were performed in the FAS. Neither maternal age nor the number of available blastocysts had any significant interaction with the clinical pregnancy effect sizes obtained in the groups compared (Supplementary Table 6). Clinical pregnancy rates were comparable in fresh cycles, with the study and control groups resulting in 48.1% (154 of 320) and 44.5% (157 of 353) (*P* = 0.35). By contrast, the freeze-all cycles showed a significant difference (*P* = 0.032) with the study and control groups resulting in 49.5% (94 of 190) and 61.3% (100 of 163) (Supplementary Table 7). The baseline demographics (Supplementary Table 8) within these subgroups revealed no significant imbalances.

Pregnancy follow-up (Supplementary Table 9) showed no statistically significant difference in pregnancy outcomes for the two groups. There was no significant difference in the male-to-female ratio in the study group compared to the control group (53.5% versus 58.6%, *P* = 0.32; Supplementary Table 9).

We investigated concordance of embryo selection and found that, of the 523 participants in the study group, the deep learning algorithm and standard morphology selected the same embryo for transfer in 344 participants (65.8%), while in 179 participants, different embryos were selected by the two techniques. The clinical pregnancy rates in these two groups were 48.3% and 44.7% (Supplementary Table 10). The Gardner grades for the embryos in each group are shown in Supplementary Table 11.

Post hoc analyses

Additional analysis was performed to investigate any difference in laboratory outcomes of participants who underwent a fresh transfer and those who underwent a frozen transfer after a freeze-all cycle (Supplementary Tables 12 and 13). The participants who underwent a frozen transfer had a significantly higher number of oocytes obtained, 20.4 (19.3–21.4) in the freeze-all group, compared to 11.7 (11.3–12.0) in the fresh-transfer group. This resulted in the freeze-all group having significantly more day 5 blastocysts: 7.0 (6.6–7.5) in the freeze-all group and 4.2 (4.0–4.4) in the fresh-transfer group. The Gardner grades for the embryos in each group are shown in Supplementary Table 13.

Discussion

This RCT rigorously evaluated deep learning in embryology laboratories. The principal finding was that this study was not able to demonstrate noninferiority of deep learning in terms of clinical pregnancy rates when compared to standard morphology and a predefined prioritization scheme. However, the study did demonstrate that deep learning, as exemplified by the iDAScore, significantly accelerates evaluation times compared to standard morphology-based embryo selection.

Before this study, the performance of AI algorithms for blastocyst transfer and their impact on clinical pregnancy outcomes had not been directly compared to standard morphological criteria used by embryologists in a prospective RCT setting. Most existing studies have primarily focused on retrospective analyses of AI's capability to objectively grade embryos and blastocysts. A recent systematic review⁷ only identified three studies that report the association with live birth rate^{20–22}. Each of these studies was considerably smaller than the current trial (175 to 458 patients), used locally derived datasets with internal validation and were not RCTs^{20–22}.

Table 2 | Primary and adjusted efficacy analysis on clinical pregnancy with fetal heartbeat after the first embryo transfer

Population	Study group ^a	Control ^a	P value ^b	Crude difference ^c (95% CI)	Crude risk ratio (95% CI)	Adjusted ^d difference (95% CI)	Adjusted ^d risk ratio (95% CI)
Intention to treat	248 of 533 (46.5%) (42.2%–50.9%)	257 of 533 (48.2%) (43.9%–52.6%)	0.62	–1.7 (–7.7; 4.3)	0.96 (0.85; 1.10)	–1.5 (–5.8; 2.8)	0.97 (0.88; 1.06)
Per protocol	237 of 500 (47.4%) (43.0%–51.9%)	245 of 502 (48.8%) (44.4%–53.3%)	0.66	–1.4 (–7.6; 4.8)	0.97 (0.85; 1.10)	–1.4 (–5.8; 2.9)	0.97 (0.88; 1.07)

^aFor categorical variables proportion, percentage and exact 95% CI is presented. ^bFor comparison between groups, Fisher’s exact test (2-sided) was used. ^cConfidence limits calculated using Farrington–Manning test. ^dAdjusted for each woman’s age, number of previous stimulated IVF cycles leading to oocyte pick-up, number of oocytes, fertilization method, number of 2PN oocytes and center.




Fig. 2 | Non-inferiority analysis and risk ratio per center. a, Crude and adjusted risk difference (red circle represents mean risk difference and blue error bars represent 95% CI of the risk difference) in clinical pregnancy rates between the iDAScore and the control groups (a negative risk difference indicates lower clinical pregnancy rate in the iDAScore group) per intention-to-treat (ITT) analysis ($n = 1,066$) and PP analysis ($n = 1,002$). **b**, Risk ratio (red circle represents mean risk ratio and blue error bars represent 95% CI of the risk ratio) of clinical pregnancy rate for each participating center (a ratio >1 indicates better results in the iDAScore group).

Previously, a machine learning algorithm, used adjunctively with morphology, trained to predict blastocyst development potential on day 3 of embryo development was tested prospectively in a previous multicenter study by Kieslinger et al.¹⁷. No difference in ongoing pregnancy rate was observed when using this algorithm compared to using standard morphology. The Kieslinger study highlights one of the challenges in performing clinical studies. The study was registered in 2015, but blastocyst stage transfer is now routinely performed by most clinics. Similarly, the known implantation data score (KIDScore), a morphokinetic algorithm requiring manual evaluation of embryos, has been prospectively evaluated¹⁸. No difference in ongoing pregnancy rates between KIDScore and standard morphology were reported, with no notable workflow efficiency due to the manual input requirement.

Our study, using a deep learning algorithm in combination with time-lapse, diverges from these approaches by assessing blastocyst development without the need for manual inputs, thus reducing evaluation time. In combination with the use of time-lapse incubation systems, deep learning embryo assessment offers the potential for minimizing time and risks associated with handling and moving embryos in the laboratory²³. However, potential laboratory efficiency gains from deep learning are only a component of the costs of IVF and have to be considered within the context of formal cost-effectiveness studies of the complex health economics of this emerging technology.

Although the pregnancy rates were clinically similar between the two groups, we could not conclude noninferiority because the lower bound of the CI surpassed our predetermined noninferiority margin of -5% . The study design of noninferiority was selected as the primary scientific objective of our study to evaluate whether the automated selection of a single blastocyst for transfer by the deep learning algorithm (iDAScore) yields a clinical pregnancy rate comparable to that

achieved by trained embryologists using standard morphology criteria and a predefined prioritization scheme.

An important deviation from the predefined hypothesis was the unexpectedly higher pregnancy rates (48.2%) in the control group, which significantly exceeded the anticipated rate of 35.4% , calculated from retrospective data from a population meeting the entry criteria to this study, used for the sample size calculation. This deviation negatively impacted on the power of this trial to conclude noninferiority. The higher pregnancy rates observed in both groups, surpassing typical rates reported in US, European and Australian national datasets²⁴, may be a result of the participation in an RCT environment (the Hawthorne effect²⁵). For example, a similar prospective trial assessing the efficacy of freezing all embryos²⁶ observed similar elevated pregnancy rates. The higher pregnancy rates observed could also be an outcome of the rigorous morphological assessment protocol employed. As part of our trial design, we standardized embryo selection across participating centers, using a study-specific prioritization scheme (detailed in the Supplementary Information), based on the Gardner grading scheme²⁷. This standardization, whether through AI or a uniform morphological assessment protocol, suggests potential for enhancing outcomes compared to current variable practices. This finding underscores the importance of consistency in embryo evaluation methodologies⁴, which has consistently been shown by AI on static images and time-lapse sequences^{8–13}, and hints at the potential benefits of integrating standardized approaches in IVF procedures.

Regardless of the cause of the higher pregnancy rates observed, future trials to assess an effect of this magnitude, assuming similar control group pregnancy rates and trial parameters (5% noninferiority margin, true difference of -1.7% , 90% power, $\alpha = 0.05$ and $\beta = 0.10$) would require an impractically larger sample size to demonstrate




Fig. 3 | Time use for day 5 evaluation in the iDAScore group and control group. Mean and standard deviation for all patients ($n = 38$) and for embryo cohort sizes from 2–5 ($n = 9$), 6–9 ($n = 16$) or 10 or more ($n = 13$) in the treatment.

noninferiority, estimated at around 7,800 participants²⁸. The inability of a practically sized trial to detect a small but clinically important effect of this sort sets a challenge for the future design of RCTs.

We observed an inconsistency in the performance of the deep learning model between fresh- and frozen-embryo transfers. In contrast to the fresh-embryo transfers, where the iDAScore group had a 3.7% higher clinical pregnancy rate, embryo selection by the deep learning model significantly underperformed compared to the control in the frozen-embryo group. This finding was surprising as previous studies based on retrospective data have found a significantly better iDAScore ranking in thawed-blastocyst data in older women²⁹ and thawed-euploid transfers³⁰. The reason for the disparity is unclear. In the freeze-all cases, there were more embryos to choose from, and this may be a factor in the difference or it may be speculated that elements of the basis of iDAScore analysis preferentially selected embryos with a predisposition to a poorer freeze–thaw performance. Finally, it is possible that the outcome observed in this trial for frozen embryos could be attributable to chance alone as this was an observational post hoc analysis. It should be noted that the clinical pregnancy rate in the fresh transfers in the control group was 44.5%, whereas the frozen-embryo transfers in the same group had a remarkably higher clinical pregnancy rate of 61.3%. Further investigation into the factors influencing outcomes in frozen-embryo transfer is warranted.

While live birth is normally perceived as the definitive outcome in studies of assisted reproduction, this study used clinical pregnancy as the primary outcome, while reporting live birth as a secondary outcome. This was on the basis that the deep learning system was specifically trained on clinical pregnancy^{12,13,29,31} and the aim of the trial was to test whether iDAScore achieves noninferiority in the endpoint on which it had been trained. However, analysis of the live birth data did not materially alter the conclusion reached by the trial.

Recently, several authors have expressed concerns about possible biases introduced by AI concerning sex ratios³². For example, Ueno et al.³¹ observed a nonsignificant increase in the male ratio with increasing iDAScore on a large retrospective live birth dataset. However, this was not confirmed in our prospective study, where no significant difference was found in the male-to-female ratio.

Another ethical concern when using deep learning for embryo selection is the black-box nature of such models³². Some studies have investigated explainability by introducing so-called heat maps to show where and when a deep learning network focuses when generating a score¹⁶. However, the clinical value of such approaches needs further

studies. Currently, most studies on explainability have investigated the correlation between well-established morphological and morphokinetic parameters and the output from deep learning models^{13,30}. These studies have found a strong correlation between iDAScore and manual embryo morphology and morphokinetics, suggesting that the deep learning models directly or indirectly focus on image features in a way similar to that done by embryologists. This study did not add to the understanding of how AI interprets embryogenesis. However, ongoing improvements in AI methodologies, coupled with interdisciplinary research efforts, will gradually enhance our collective knowledge of embryogenesis, ultimately contributing to the refinement of assisted reproductive technologies.

It is important to acknowledge several limitations in our trial. First, iDAScore was derived and tested solely within the context of the EmbryoScope incubator, limiting its generalizability to other time-lapse incubator systems. Second, the time-to-pregnancy was not assessed, as only the first embryo was prioritized for transfer, leaving an equivalent number of embryos available for future use in both groups. Similarly, we have not reported cumulative live birth rates because that would require transfer of all embryos, although we anticipate this to be similar as no embryos were deselected for use based on the iDAScore. As we had underestimated the time required for standard morphological criteria assessment, a smaller substudy than planned was required to show the observed time differences. Last, the continued evolution of deep learning algorithms³³ presents a challenge for ongoing evaluation via traditional RCTs, suggesting the necessity for alternative research methodologies in assessing future iterations³⁴.

The present randomized trial examined the efficacy of using a deep learning algorithm for the selection of which embryo to transfer for couples undertaking assisted conception. This study was unable to demonstrate noninferiority in clinical pregnancy rate to standard morphology. However, the deep learning approach studied did provide a consistent user-independent approach with a 10-fold reduction in assessment time.

Online content

Any methods, additional references, Nature Portfolio reporting summaries, source data, extended data, supplementary information, acknowledgements, peer review information; details of author contributions and competing interests; and statements of data and code availability are available at <https://doi.org/10.1038/s41591-024-03166-5>.

References

- Topol, E. J. High-performance medicine: the convergence of human and artificial intelligence. *Nat. Med.* **25**, 44–56 (2019).
- Lang, K. et al. Artificial intelligence-supported screen reading versus standard double reading in the mammography screening with Artificial Intelligence trial (MASAI): a clinical safety analysis of a randomized, controlled, non-inferiority, single-blinded, screening accuracy study. *Lancet Oncol.* **24**, 936–944 (2023).
- Alpha Scientists in Reproductive Medicine and ESHRE Special Interest Group of Embryology. The Istanbul consensus workshop on embryo assessment: proceedings of an expert meeting. *Hum. Reprod.* **26**, 1270–1283 (2011).
- Storr, A., Venetis, C. A., Cooke, S., Kilani, S. & Ledger, W. Inter-observer and intra-observer agreement between embryologists during selection of a single day 5 embryo for transfer: a multicentre study. *Hum. Reprod.* **32**, 307–314 (2017).
- Steptoe, P. C. & Edwards, R. G. Birth after reimplantation of a human embryo. *Lancet* **2**, 366 (1978).
- Armstrong, S., Bhide, P., Jordan, V., Pacey, A. & Farquhar, C. Time-lapse systems for embryo incubation and assessment in assisted reproduction. *Cochrane Database Syst. Rev.* **5**, CD011320 (2018).

7. Salih, M. et al. Embryo selection through artificial intelligence versus embryologists: a systematic review. *Hum. Reprod. Open* **3**, hoad031 (2023).
8. Dirvanauskas, D., Maskeliunas, R., Raudonis, V. & Damasevicius, R. Embryo development stage prediction algorithm for automated time lapse incubators. *Comput. Methods Prog. Biomed.* **177**, 161–174 (2019).
9. Kragh, M. F., Rimestad, J., Berntsen, J. & Karstoft, H. Automatic grading of human blastocysts from time-lapse imaging. *Comput. Biol. Med.* **115**, 103494 (2019).
10. Miyagi, Y., Habara, T., Hirata, R. & Hayashi, N. Feasibility of deep learning for predicting live birth from a blastocyst image in patients classified by age. *Reprod. Med. Biol.* **18**, 190–203 (2019).
11. VerMilyea, M. et al. Development of an artificial intelligence-based assessment model for prediction of embryo viability using static images captured by optical light microscopy during IVF. *Hum. Reprod.* **35**, 770–784 (2020).
12. Tran, A., Cooke, S., Illingworth, P. J. & Gardner, D. K. Deep learning as a predictive tool for fetal heart pregnancy following time-lapse incubation and blastocyst transfer. *Hum. Reprod.* **34**, 1011–1018 (2019).
13. Berntsen, J., Rimestad, J., Lassen, T. J., Tran, D. & Kragh, M. F. Robust and generalizable embryo selection based on artificial intelligence and time-lapse sequences. *PLoS ONE* **17**, e0262661 (2022).
14. Jiang, V. S. & Bormann, C. L. Artificial intelligence in the in vitro fertilization laboratory: a review of advancements over the last decade. *Fertil. Steril.* **120**, 17–23 (2023).
15. Bormann, C. L. et al. Consistency and objectivity of automated embryo assessments using deep neural networks. *Fertil. Steril.* **113**, 781–787 (2020).
16. Payá, E., Bori, L., Colomer, A., Meseguer, M. & Naranjo, V. Automatic characterization of human embryos at day 4 post-insemination from time-lapse imaging using supervised contrastive learning and inductive transfer learning techniques. *Comput. Methods Prog. Biomed.* **221**, 106895 (2022).
17. Kieslinger, D. C. et al. Clinical outcomes of uninterrupted embryo culture with or without time-lapse-based embryo selection versus interrupted standard culture (SelectIMO): a three-armed, multicentre, double-blind, randomised controlled trial. *Lancet* **401**, 1438–1446 (2023).
18. Ahlström, A., Lundin, K. & Lind, A. K. A double-blind randomized controlled trial investigating a time-lapse algorithm for selecting day 5 blastocysts for transfer. *Hum. Reprod.* **37**, 708–717 (2022).
19. Kim, K. G. Book review: deep learning. *Health. Inform. Res.* **22**, 351 (2016).
20. Sawada, Y. et al. Evaluation of artificial intelligence using time-lapse images of IVF embryos to predict live birth. *Reprod. Biomed. Online* **43**, 843–852 (2021).
21. Benchaib, M., Labrune, E., Giscard d'Estaing, S., Salle, B. & Lornage, J. Shallow artificial networks with morphokinetic time-lapse parameters coupled to ART data allow to predict live birth. *Reprod. Biomed. Online* **21**, e12486 (2022).
22. Wang, S., Chen, L. & Sun, H. Interpretable artificial intelligence-assisted embryo selection improved single-blastocyst transfer outcomes: a prospective cohort study. *Reprod. Biomed. Online* **47**, 103371 (2023).
23. Alikani, M., Go, K. J., McCaffrey, C. & McCulloh, D. H. Comprehensive evaluation of contemporary assisted reproduction technology laboratory operations to determine staffing levels that promote patient safety and quality care. *Fertil. Steril.* **102**, 1350–1356 (2014).
24. De Geyter, C. et al. Twenty years of the European IVF-monitoring Consortium registry: what have we learned? A comparison with registries from two other regions. *Hum. Reprod.* **35**, 2832–2849 (2020).
25. McCambridge, J., Witton, J. & Elbourne, D. R. Systematic review of the Hawthorne effect: New concepts are needed to study research participation effects. *J. Clin. Epidemiol.* **67**, 267–277 (2014).
26. Wei, D. et al. Frozen versus fresh single blastocyst transfer in ovulatory women: a multicentre randomised controlled trial. *Lancet* **393**, 1310–1318 (2019).
27. Gardner D. K. & Schoolcraft W. B. in *Towards Reproductive Certainty: Infertility and Genetics Beyond* (eds Jansen R. & Mortimer D.) 378–388 (Parthenon, 1999).
28. D'Agostino, R. B. Sr., Massaro, J. M. & Sullivan, L. M. Non-inferiority trials: design concepts and issues – the encounter of academic consultants in statistics. *Stat. Med.* **22**, 169–186 (2003).
29. Ueno, S. et al. Pregnancy prediction performance of an annotation-free embryo scoring system on the basis of deep learning after single vitrified-warmed blastocyst transfer: a single-center large cohort retrospective study. *Fertil. Steril.* **116**, 1172–1180 (2021).
30. Cimadomo, D. et al. Towards automation in IVF: pre-clinical validation of a deep learning-based embryo grading system during PGT-A cycles. *J. Clin. Med.* **12**, 1806 (2023).
31. Ueno, S., Berntsen, J., Ito, M., Okimura, T. & Kato, K. Correlation between an annotation-free embryo scoring system based on deep learning and live birth / neonatal outcomes after single vitrified-warmed blastocyst transfer: a single-centre, large-cohort retrospective study. *J. Assist. Reprod. Genet.* **39**, 2089–2099 (2022).
32. Afnan, M. A. M. et al. Interpretable, not black-box, artificial intelligence should be used for embryo selection. *Hum. Reprod. Open* **2021**, hoab040 (2021).
33. Lassen, J. T., Kragh, M. F., Rimestad, J., Johansen, M. N. & Berntsen, J. Development and validation of deep learning based embryo selection across multiple days of transfer. *Sci. Rep.* **13**, 4235 (2023).
34. Hernán, M. A. & Robins, J. M. Using big data to emulate a target trial when a randomized trial is not available. *Am. J. Epidemiol.* **183**, 758–764 (2016).

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2024

Peter J. Illingworth¹✉, Christos Venetis^{2,3,4}, David K. Gardner^{5,6}, Scott M. Nelson^{7,8}, Jørgen Berntsen⁹, Mark G. Larman¹⁰, Franca Agresta¹¹, Saran Ahitan¹², Aisling Ahlström¹³, Fleur Cattrall⁵, Simon Cooke², Kristy Demmers¹⁴, Anette Gabrielsen¹⁵, Johnny Hindkjær¹⁶, Rebecca L. Kelley⁵, Charlotte Knight², Lisa Lee¹⁵, Robert Lahoud², Manveen Mangat², Hannah Park¹⁷, Anthony Price¹⁸, Geoffrey Trew^{8,19}, Bettina Troest²⁰, Anna Vincent⁸, Susanne Wennerström¹³, Lyndsey Zujovic¹² & Thorir Hardarson¹⁰

¹Virtus Health, Sydney, New South Wales, Australia. ²IVFAustralia, Sydney, New South Wales, Australia. ³Unit for Human Reproduction, 1st Dept of Ob/Gyn, Medical School, Faculty of Health Sciences, Aristotle University of Thessaloniki, Thessaloniki, Greece. ⁴Centre for Big Data Research in Health, Faculty of Medicine and Health, University of New South Wales, Sydney, New South Wales, Australia. ⁵Melbourne IVF, Melbourne, Victoria, Australia. ⁶School of BioSciences, University of Melbourne, Parkville, Victoria, Australia. ⁷School of Medicine, Dentistry and Nursing, University of Glasgow, Glasgow, UK. ⁸TFP Fertility, Institute of Reproductive Sciences, Oxford, UK. ⁹Vitrolife, Viby J, Denmark. ¹⁰Vitrolife, Gothenburg, Sweden. ¹¹Virtus Health, Melbourne, Victoria, Australia. ¹²TFP Fertility, Nottingham, UK. ¹³IVIRMA Global Research Alliance, Livio Gothenburg, Gothenburg, Sweden. ¹⁴Queensland Fertility Group, Brisbane, Queensland, Australia. ¹⁵The Fertility Unit, Horsens Hospital, Horsens, Denmark. ¹⁶Aagaard, Aarhus, Denmark. ¹⁷Dept of Reproductive Medicine, Sahlgrenska University Hospital, Gothenburg, Sweden. ¹⁸TFP Fertility, Southampton, UK. ¹⁹Imperial College London, London, UK. ²⁰The Fertility Unit, Aalborg University Hospital, Aalborg, Denmark. ✉e-mail: peter.illingworth@virtushealth.com.au

Methods

Trial design and oversight

This randomized, noninferiority, double-blind parallel-group controlled trial was carried out at 14 IVF clinics in Australia and Europe. Written informed consent was obtained from all participants. Each participating clinic's respective Human Research Ethics Committee reviewed and approved the trial protocol, which had been previously registered with the Australian New Zealand Clinical Trials Registry (ANZCTR) registration: 379161. All the data in the study were collected using a commercial eCRF provider MariaDB v.10.6 and Database Platform VI v.7.1.6 by Stockholm Data Design, Sweden. The randomization was performed within the eCRF using web-based interactive response technology. The statistical analysis plan and detailed protocol are available in the Supplementary Information. A steering committee, responsible for trial design and the manuscript's first draft, supervised the trial. They affirm the data completeness and accuracy and the trial's fidelity to the protocol. An independent data safety and monitoring board provided consistent oversight for the trial's conduct. The trial was implemented using EmbryoScope incubators and received funding from the manufacturer, Vitrolife, Denmark.

Study population

The study was conducted between January 2020 and September 2022. We included women undergoing IVF or ICSI after ovarian stimulation with gonadotrophins and gonadotrophin releasing hormone analogs and with the intention to treat by either transfer of a single fresh embryo on day 5 or, in a freeze-all cycle, the first thawed embryo. Inclusion criteria required women to be before their 42nd completed birthday with at least two early blastocysts on day 5 of embryo culture, the day of randomization. We excluded participants if they were involved in treatment using donated eggs, intended to perform preimplantation genetic testing, used additional laboratory interventions such as intracytoplasmic morphologically selected sperm injection, polarized light microscopy, assisted hatching or had previously participated in the trial.

Ovarian stimulation and triggering of final oocyte maturation was performed under each site's standard protocol, supervised by the treating physician. After oocyte retrieval, all embryos were fertilized by either IVF or ICSI based on individual clinical decisions and incubated at 5% oxygen in the EmbryoScope+ time-lapse incubator system (Vitrolife) until day 5 or 6.

Provided at least two embryos reached the early blastocyst stage (grade 2 according to Gardner grading)¹⁸ or beyond at 114–118 hours post insemination (hpi) (day 5 of embryo culture), patients were eligible for randomization. The embryos were assessed using standard morphology criteria (Gardner grade)¹⁸ previously demonstrated in a similar population¹⁸. To ensure that the embryologist was blinded to the randomization at the time of morphology selection, the highest-grade blastocyst was selected for transfer, using the prioritization strategy detailed in the clinical investigation protocol in the Supplementary Information, before randomization. Once this selection had been performed and documented, the participant was then randomized through a web-based interactive response technology (MariaDB v.10.6 and Database Platform VI v.7.1.6 by Stockholm Data Design) to ensure allocation concealment.

If the patient was randomized into the control arm, the blastocyst selected on morphological grounds was transferred and iDAScore was not performed. If the patient was randomized into the study arm, iDAScore v.1.2 was performed to generate a score for each embryo and the blastocyst with the highest iDAScore was then selected for transfer.

The iDAScore v.1 algorithm was trained and evaluated based on a large dataset from 18 IVF centers consisting of 115,832 embryos, of which 14,644 were transferred embryos with known outcome¹³. The algorithm is based on an inflated three-dimensional convolutional neural network where the output is fed to a bidirectional long short-term

memory. The input to the model was 128 images of the central focal plane and a resolution of 256×256 pixels. The sequence was offset by 12 hpi, thus effectively covering 12 to 140 hpi. The v.1 iDAScore was used throughout the study period and no clinic-specific or other adjustments were applied during the course of the study.

The trial-group assignment was performed in an unblinded manner to the embryologist, but both the treating clinician and the patient remained blinded to the randomization outcome until after the first embryo transfer. Any remaining embryos were cryopreserved if they met the normal freezing criteria for the laboratory or, in addition, in the study arm if they had reached Gardner grade 3 or beyond and had an iDAScore of 5 or more. All remaining embryos were re-evaluated on day 6, rescored at 138–142 hpi and assessed using the Gardner grade and iDAScore. In both groups, the prioritization of frozen embryos for subsequent warming and transfer was based on all information up to day 6. All blastocysts were transferred in EmbryoGlue (Vitrolife). Embryo transfer and luteal support were performed and administered according to each clinic's standard protocol.

Time taken to evaluate the embryos for transfer

We compared the time taken to evaluate the same embryos across three laboratories using the two assessment techniques. We initiated a timer from the moment of opening the patient file in the EmbryoScope+ software until the selection of the embryo for transfer, using either iDAScore or standard morphology criteria.

Outcomes

The primary outcome for this study was the achievement of **clinical pregnancy after the first embryo transfer, whether fresh or frozen–thawed in freeze-all cycles**. We defined clinical pregnancy as an intrauterine gestation with a fetal heartbeat observed after 7–9 weeks gestation. Secondary outcomes included positive hCG rate, nonviable intrauterine pregnancies, ongoing pregnancy rate (at 12 weeks gestation) and live birth rate. Nonviable intrauterine pregnancy was defined as a pregnancy in which an intrauterine gestation was seen with no fetal heartbeat found. Live birth was defined as delivery of at least one liveborn child.

Statistical analyses

The noninferiority margin for this study was established at -5% for clinical pregnancy rate. On the basis of existing evidence from participating clinics, we expected the clinical pregnancy rate following the first embryo transfer to be 35.4% . To demonstrate with 90% power ($\alpha = 0.05$ and $\beta = 0.10$) that the lower limit of the two-sided 95% CI for the difference between the iDAScore and the standard morphology criteria group would not be less than -5% , with an expected increase in clinical pregnancy of 5% or more in the iDAScore group, we required 494 women per group. This figure was increased to 520 women per group to account for a potential 5% of loss to follow-up.

For the time taken to evaluate embryos for transfer substudy, the standard deviation for the difference between the morphological (Gardner grade) and iDAScore assessment was estimated as 60.6 seconds from a pilot study of 20 patients. To find a mean difference in times between Gardner grade and iDAScore of 45 seconds with a two-sided Fisher's nonparametric permutation test for paired observations, on significance level 0.05, with a power of 80% we required a minimum of 51 patients. The substudy was performed in three clinics and, to ensure a representation of clinical practice, different embryo cohort sizes, ranging from small (2–5 embryos), medium (6–10 embryos) to large (>10 embryos), were included.

The primary efficacy analysis was performed in the intention-to-treat population (denoted as the ITT in the clinical investigation protocol and statistical analysis plan in the Supplementary information), including all patients who underwent randomization and had an embryo transfer. All randomized subjects with measurement of

the primary efficacy value were included in the FAS. We conducted complementary analyses on the PP population. Planned and prespecified sensitivity analyses, adjusting for significant a priori specified covariates using multivariable logistic regression were performed. These covariates included each woman's age (<25, 25 to <30, 30 to <35, 35 to <40 and ≥40), number of previous stimulated IVF cycles leading to oocyte pick-up (0 or >1), number of oocytes (2 to 5, 6 to 10, 11 to 15, 16 to 20, 21 to 25 and >26), fertilization method (IVF or ICSI), number of two pronuclear oocytes (2 to 5, 6 to 10, 11 to 15, 16 to 20, 21 to 25 and >26) and center. We also undertook a prespecified subgroup analysis for the older female age groups (>35 years) and an interaction analysis according to each woman's age and the total number of embryos available for selection. Last, we compared the performance of the iDAScore algorithm at each individual center.

For the time taken to evaluate embryo substudy, we used Fisher's nonparametric permutation test for paired observations and estimated the differences between the morphological and iDAScore with varying embryo numbers. All the statistical analyses were executed using SAS v.9.4, with the support of N.-G. Pehrsson of Statistiska Konsultgruppen. Additional details are provided in the Methods section of the Supplementary Appendix.

Role of the funding source

The project was funded by a grant from Vitrolife. As sponsor, employees of Vitrolife Group participated in the study design and writing of the report. Data collection and analysis were performed by independent entities.

Reporting summary

Further information on research design is available in the Nature Portfolio Reporting Summary linked to this article.

Data availability

Data collected for the study, including deidentified participant data and the data dictionary, will be made available to others. Related documents, including the study protocol, the statistical analysis plan and the informed consent forms will be available following publication. The data and documents will be made available on request to academic researchers following review by the study's steering committee (P.J.I., C.V., D.K.G., S.M.N., J.B., M.G.L. and T.H.) and completion of a data sharing agreement. Requests for data sharing should be addressed to the corresponding author. Approval of requests for academic purposes will be provided within 3 weeks and the data supplied following approval. Approval will not be provided for commercial use of the data.

Code availability

The iDAScore algorithm is commercially available. It was developed by Vitrolife to be used within the EmbryoViewer workstation with EmbryoScope time-lapse incubators. The code is proprietary, but iDAScore and the necessary hardware can be obtained from Vitrolife (www.vitrolife.com).

Acknowledgements

The project was funded by a grant from Vitrolife. The funders had no role in study design. Three employees of the funder formed part of the steering panel overseeing data collection, analysis, decision to publish and preparation of the paper. We thank all the participants in this study and the following people for their contribution to the paper: S. Apter and C. Stakeberg (Vitrolife, Sweden) supported and coordinated the

clinical study; M. Montag contributed to the editing of this paper; N.-G. Pehrsson, P. Ekman, M. Molin and H. Imberg at Statistiska Konsultgruppen contributed to the statistical analysis of our data; and our many medical colleagues, nursing staff and embryologists who contributed to patient recruitment and data collection.

Author contributions

P.J.I., C.V., D.K.G., S.M.N. and T.H. contributed to original study design, were involved in patient recruitment, took part in data analysis and drafted the paper. J.B. and M.G.L. contributed to original study design, took part in data analysis and drafted the paper. F.A., S.A., A.A., F.C., S.C., K.D., A.G., J.H., R.L.K., C.K., L.L., R.L., M.M., H.P., A.P., G.T., B.T., A.V., S.W. and L.Z. were involved in patient recruitment and reviewed the content of the paper. All authors had full access to all the data in the study, all had approval of the version to submit for publication and all agree to be accountable for the content. P.J.I. (academic team), J.B. and T.H. had regular access to and were able to verify the underlying data.

Competing interests

P.J.I. has provided paid advice to Western Australia government on medical cases and has previously (until June 2022) held shares in Virtus Health, a commercial IVF company. C.V. has received consulting fees, lecture payments and travel expenses from pharmaceutical firms with a commercial portfolio in IVF, and has previously (until June 2022) held shares in Virtus Health, a commercial IVF company. D.K.G. has received a research grant from Vitrolife to develop culture medium. S.M.N. has received consulting fees, lecture payments and travel expenses from pharmaceutical firms with a commercial portfolio in IVF. J.B. and M.G.L. are employees of Vitrolife and own shares in Vitrolife, who have a pending iDAScore patent. R.L.K. has received a grant from a pharmaceutical firm with a commercial portfolio in IVF, has received speaking fees and travel expenses from Vitrolife, who have a pending iDAScore patent. R.L. has received lecture payments and travel expenses from pharmaceutical firms with a commercial portfolio in IVF. M.M. has received lecture payments and travel expenses from pharmaceutical firms with a commercial portfolio in IVF and has previously (until June 2022) held shares in Virtus Health, a commercial IVF company. H.P. and B.T. are employed by a company where Vitrolife funded research. T.H. was previously employed by Vitrolife, who have a pending iDAScore patent. The other authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41591-024-03166-5>.

Correspondence and requests for materials should be addressed to Peter J. Illingworth.

Peer review information *Nature Medicine* thanks Kelley M. Kidwell, Dorit Kieslinger, Hadi Shafiee and the other, anonymous, reviewer(s) for their contribution to the peer review of this work. Primary Handling Editor: Sonia Muliyil, in collaboration with the *Nature Medicine* team.

Reprints and permissions information is available at www.nature.com/reprints.

Reporting Summary

Nature Portfolio wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Portfolio policies, see our [Editorial Policies](#) and the [Editorial Policy Checklist](#).

Statistics

For all statistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.

- | | |
|-------------------------------------|--|
| n/a | Confirmed |
| ☐ | ☒ The exact sample size (<i>n</i>) for each experimental group/condition, given as a discrete number and unit of measurement |
| ☐ | ☒ A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly |
| ☐ | ☒ The statistical test(s) used AND whether they are one- or two-sided
<i>Only common tests should be described solely by name; describe more complex techniques in the Methods section.</i> |
| ☐ | ☒ A description of all covariates tested |
| ☐ | ☒ A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons |
| ☐ | ☒ A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals) |
| ☐ | ☒ For null hypothesis testing, the test statistic (e.g. <i>F</i> , <i>t</i> , <i>r</i>) with confidence intervals, effect sizes, degrees of freedom and <i>P</i> value noted
<i>Give P values as exact values whenever suitable.</i> |
| ☐ | ☒ For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings |
| ☐ | ☒ For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes |
| ☒ | ☐ Estimates of effect sizes (e.g. Cohen's <i>d</i> , Pearson's <i>r</i>), indicating how they were calculated |

Our web collection on [statistics for biologists](#) contains articles on many of the points above.

Software and code

Policy information about [availability of computer code](#)

Data collection	The image analysis was performed by iDAScore Version 1.2 (www.vitrolife.com)The iDAScore® algorithm is commercially available. It was developed by Vitrolife to be used within the EmbryoViewer workstation with EmbryoScope time-lapse incubators. The code is proprietary, but iDAScore and the necessary hardware can be obtained from Vitrolife (www.vitrolife.com). All data in the study was collected using a commercial eCRF provider MariaDB v10.6 and Database Platform VI v7.1.6 by Stockholm Data Design, Sweden. The randomisation was performed within the eCRF using Web-based interactive response technology.
Data analysis	Raw analysis data was stored as csv files and analyzed using SAS v9.4 as described in detail in the Statistical Analysis Plan

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Portfolio [guidelines for submitting code & software](#) for further information.

Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A description of any restrictions on data availability
- For clinical datasets or third party data, please ensure that the statement adheres to our [policy](#)

Data collected for the study, including deidentified participant data, and the data dictionary, will be made available to others. Related documents, including the study protocol, the statistical analysis plan and the informed consent forms will be available following publication. The data and documents will be made available on request to academic researchers following review by the Study steering committee (PI, CV, DG, SN, JB, ML, TH) and completion of a data sharing agreement. Requests for data sharing should be addressed to the corresponding author. Approval of requests for academic purposes will be provided within three weeks and the data supplied following approval. Approval will not be provided for commercial use of the data.

Human research participants

Policy information about [studies involving human research participants and Sex and Gender in Research](#).

Reporting on sex and gender

This is a study of embryos. All the participants giving consent are women from whose eggs, the resulting embryos are being studied in the project and their partners (whether male or female).

Population characteristics

The study was conducted between January 2020 and September 2022. We included women undergoing IVF or ICSI after ovarian stimulation with gonadotrophins and gonadotrophin releasing hormone analogues and with the intention to treat by either transfer of a single fresh embryo on day 5 or, in a freeze-all cycle, the first rewarmed embryo. Inclusion criteria required women to be before their 42nd completed birthday with at least two early blastocysts on day 5 of embryo culture, the day of randomisation. We excluded participants if they were involved in treatment using donated eggs, intended to perform preimplantation genetic testing, used additional laboratory interventions such as intracytoplasmic morphologically selected sperm injection (IMSI), polarised light microscopy, assisted hatching, or had previously participated in the trial.

Recruitment

The study was conducted between January 2020 and September 2022. Potential participants were identified by staff members and their eligibility reviewed by one of investigator. If eligible, informed consent was sought. The resulting population, was self-selected for willingness to participate in the trial. No self-selection biases in the overall population were apparent.

Ethics oversight

IVFAustralia HREC, an HREC that is accredited with the Australian NHMRC regulating Ethics Committee.

Note that full information on the approval of the study protocol must also be provided in the manuscript.

Field-specific reporting

Please select the one below that is the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

☒ Life sciences ☐ Behavioural & social sciences ☐ Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see [nature.com/documents/nr-reporting-summary-flat.pdf](https://www.nature.com/documents/nr-reporting-summary-flat.pdf)

Life sciences study design

All studies must disclose on these points even when the disclosure is negative.

Sample size

The non-inferiority margin for this study was established at -5% for clinical pregnancy rate. On the basis of existing evidence from participating clinics, we expected the clinical pregnancy rate following the first embryo transfer to be 35.4%. To demonstrate with 90% power ($\alpha=0.05$ & $\beta=0.10$) that the lower limit of the two-sided 95% confidence interval (CI) for the difference between the iDAScore and the standard morphology criteria group would not be less than -5%, with an expected increase in clinical pregnancy of 5% or more in the iDAScore group, we required 494 women per group. This figure was increased to 520 women per group to account for a potential 5% of loss to follow-up.

Data exclusions

No data were excluded

Replication

Not applicable to a randomised controlled trial

Randomization

The participant was allocated to treatment by randomisation through a Web-based interactive response technology (MariaDB v10.6 and Database Platform VI v7.1.6 by Stockholm Data Design, Sweden) to ensure allocation concealment.

Blinding

The embryologist was blinded to the randomisation at the time of morphology selection, by ensuring that the highest-grade morphology blastocyst was selected, using the prioritisation strategy detailed in the CIP Appendix, prior to randomisation. Both the treating clinician and

the patient remained blinded to the randomisation outcome until after the first embryo transfer.

Reporting for specific materials, systems and methods

We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.

Materials & experimental systems

n/a	Involved in the study
☒	☐ Antibodies
☒	☐ Eukaryotic cell lines
☒	☐ Palaeontology and archaeology
☒	☐ Animals and other organisms
☐	☒ Clinical data
☒	☐ Dual use research of concern

Methods

n/a	Involved in the study
☒	☐ ChIP-seq
☒	☐ Flow cytometry
☒	☐ MRI-based neuroimaging

Clinical data

Policy information about [clinical studies](#)

All manuscripts should comply with the ICMJE [guidelines for publication of clinical research](#) and a completed [CONSORT checklist](#) must be included with all submissions.

Clinical trial registration Australian New Zealand Clinical Trial Registry (ANZCTR #ID 379161)

Study protocol Attached

Data collection The study was conducted between January 2020 and September 2022 at 14 IVF clinics in Australia and Europe

Outcomes The primary outcome for this study was the achievement of clinical pregnancy after the first embryo transfer, whether fresh or frozen-thawed in freeze-all cycles. We defined clinical pregnancy as an intrauterine gestation with a fetal heartbeat observed after 7-9 weeks gestation. Secondary outcomes included positive hCG rate, non-viable intrauterine pregnancies, ongoing pregnancy rate (at 12 weeks gestation) and live birth rate. Non-viable intrauterine pregnancy was defined as a pregnancy where an intrauterine gestation was seen with no fetal heart found. Live birth was defined as delivery of at least one liveborn child. We compared the time taken to evaluate the same embryos across three laboratories using the two assessment techniques