

```

t1=0:0.001:2;
t2=0:0.001:1;
s1=(t1<1)&(t1~=0);
s2=(t2<=1/4)+(t2>=3/4)-1/2;
%Sketch the signal on the interval [0, 2].
figure;
plot(t1,s1);xlabel('t'),ylabel('s(t)');xlim([-0.2 2.2]);ylim([-0.2 1.2]);

plot(t2,s2);xlabel('t'),ylabel('s(t)');xlim([-0.05 1.05]);ylim([-0.55 0.55]);

```

### 3.2.1 Synthesis of Periodic Signals

1. Period  $T_0 = 2$ . For  $t \in [0, 2]$ :

$$s(t) = \text{rect}\left(t - \frac{1}{2}\right)$$

2. Period  $T_0 = 1$ . For  $t \in \left[-\frac{1}{2}, \frac{1}{2}\right]$ :

$$s(t) = \text{rect}(2t) - \frac{1}{2}$$

$$s(t) = a_0 + \sum_{k=1}^{\infty} A_k \sin(2\pi k f_0 t + \theta_k)$$

Signal 1:  $T_0 = 2$ ,  $s(t) = \text{rect}(t - 1/2)$

```

clear;
% Define signal parameters
T0_1 = 2; % Period
f0_1 = 1 / T0_1; % Fundamental frequency
t = linspace(0, T0_1, 1000); % Time vector for plotting

% Compute Fourier series coefficients
k_max = 10; % Number of terms in the series

s_t_1 = zeros(size(t)); % Initialize the signal

for k = 1:k_max

    % Construct the signal using the computed coefficients


```

```

s_t_1=s_t_1+2*sin((2*k-1)*pi/2)/((2*k-1)*pi)*sin((2*k-1)*pi.*(t-0.5)+pi/2);
end

% Plot the signal
figure;
subplot(2, 1, 1);
plot(t, s_t_1);
title('Signal  $\diamond(\diamond) = \text{rect}(\diamond - 1/2)$  for  $\diamond\theta = 2$ ');
xlabel('Time (t)');
ylabel('Amplitude');

```



Signal 2:  $\diamond\theta = 1$ ,  $\diamond(\diamond) = \text{rect}(2\diamond) - 1/2$

```

% Define signal parameters
T0_2 = 1; % Period
f0_2 = 1 / T0_2; % Fundamental frequency
t = linspace(-0.5, 0.5, 1000); % Time vector for plotting

% Compute Fourier series coefficients
k_max = 10; % Number of terms in the series

s_t_2 = zeros(size(t)); % Initialize the signal

for k = 1:k_max

```


```

% Construct the signal using the computed coefficients
s_t_2 = s_t_2 + 2*sin((2*k-1)*pi/2)/((2*k-1)*pi)*sin((2*k-1)*pi.*(t-0.5)+pi/2);
end

% Plot the signal
figure;

plot(t, s_t_2);
title('Signal  $s(t) = \text{rect}(2t) - 1/2$  for  $T_0 = 1$ ');
xlabel('Time (t)');
ylabel('Amplitude');

```



```

% Plot the Fourier series coefficients A

```

### 3.5.1 Discrete-Time Fourier Transform

$$X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n]e^{-j\omega n}$$

Since  $X(e^{j\omega})$  is a periodic function of  $\omega$  with a period of  $2\pi$ , we need only to compute  $X(e^{j\omega})$  for  $-\pi < \omega < \pi$ .

Here  $n_0$  is the time index corresponding to the 1st element of the  $\mathbf{x}$  vector, and  $dw$  is the spacing between the samples of the Matlab vector  $\mathbf{X}$ . For example, if  $\mathbf{x}$  is a vector of length  $N$ , then its DTFT is computed by

$$X(\omega) = \sum_{n=1}^N x[n]e^{-j\omega(n+n_0-1)}$$

Where  $\omega$  is a vector of values formed by  $\mathbf{w}=(-\pi:dw:\pi)$ .

```
% Signal 1: x(n) = δ(n)
n = -10:10;
x1 = (n == 0); % Impulse signal
n0 = -10; % Starting index
dw = 0.01; % Frequency spacing

% Compute DTFT
X1 = DTFT(x1, n0, dw);

% Signal 2: x(n) = δ(n - 5)
n2 = -10:10;
x2 = (n2 == 5); % Impulse signal shifted to n = 5
n0 = -10; % Starting index
dw = 0.01; % Frequency spacing

% Compute DTFT
X2 = DTFT(x2, n0, dw);

% Signal 3: x(n) = (0.5)^n * u(n)
n3 = 0:20;
x3 = (0.5).^n3; % Exponential signal
n0 = 0; % Starting index
dw = 0.01; % Frequency spacing

% Compute DTFT
X3 = DTFT(x3, n0, dw);

% Create a 3x2 subplot for all six plots
figure;

% Plot Signal 1
subplot(3, 2, 1);
```

```

plot(-pi:dw:pi, abs(X1));
title('Magnitude of  $X(e^{j\omega})$  for  $\delta(n)$ ');
xlabel('\omega (radians)');
ylabel('|X(e^{j\omega})|');

subplot(3, 2, 2);
plot(-pi:dw:pi, angle(X1));
title('Phase of  $X(e^{j\omega})$  for  $\delta(n)$ ');
xlabel('\omega (radians)');
ylabel('Phase');

% Plot Signal 2
subplot(3, 2, 3);
plot(-pi:dw:pi, abs(X2));
title('Magnitude of  $X(e^{j\omega})$  for  $\delta(n - 5)$ ');
xlabel('\omega (radians)');
ylabel('|X(e^{j\omega})|');

subplot(3, 2, 4);
plot(-pi:dw:pi, angle(X2));
title('Phase of  $X(e^{j\omega})$  for  $\delta(n - 5)$ ');
xlabel('\omega (radians)');
ylabel('Phase');

% Plot Signal 3
subplot(3, 2, 5);
plot(-pi:dw:pi, abs(X3));
title('Magnitude of  $X(e^{j\omega})$  for  $(0.5)^n \cdot u(n)$ ', 'Interpreter',
'latex');
xlabel('$\omega$ (radians)', 'Interpreter', 'latex');
ylabel('$|X(e^{j\omega})|$', 'Interpreter', 'latex');

subplot(3, 2, 6);
plot(-pi:dw:pi, angle(X3));
title('Phase of  $X(e^{j\omega})$  for  $(0.5)^n \cdot u(n)$ ', 'Interpreter', 'latex');
xlabel('$\omega$ (radians)', 'Interpreter', 'latex');
ylabel('Phase', 'Interpreter', 'latex');

```

