

4 clicker questions today



# PHYS 2211 K

Week 6, Lecture 2

2022/02/17

Dr Alicea ([eaalicea@gatech.edu](mailto:eaalicea@gatech.edu))

## On today's class...

1. Curving motion problems  
(in 2D and 3D)
2. Kinesthetic sensations  
(feeling weightless, feeling heavier)

# CLICKER 1: Favorite element-bending




A. Waterbending



B. Earthbending




C. Firebending



D. Airbending

# From Tuesday

- $\hat{p}$  axis is **parallel** to the direction of the motion
- $\hat{n}$  axis is **perpendicular** to the direction of the motion and positive towards the center of the turning circle
- The coordinates move and change with the object's motion!



# Also from Tuesday

Changes the  
**speed** of the  
object

$$\vec{F}_{\text{net}\parallel} = \left( \frac{d\vec{p}}{dt} \right)_{\parallel} = \frac{d|\vec{p}|}{dt} \hat{p} = \left( \frac{|\vec{p}_f| - |\vec{p}_i|}{\Delta t} \right) \hat{p}$$

Changes the  
**direction** of  
motion



$$\vec{F}_{\text{net}\perp} = \left( \frac{d\vec{p}}{dt} \right)_{\perp} = |\vec{p}| \frac{d\hat{p}}{dt} = \frac{mv^2}{R} \hat{n}$$

$$\vec{F}_{\text{net}} = \vec{F}_{\text{net}\parallel} + \vec{F}_{\text{net}\perp}$$

# Solving curving motion problems


- Similar to equilibrium problems, but  $F_{net}$  is not equal to zero
- Components of  $F_{net}$  are not  $F_{net,x}$  and  $F_{net,y}$ , but rather  $F_{net\_parallel}$  and  $F_{net\_perp}$
- Draw FBD
  - $\hat{p}$  axis points in direction of motion
  - $\hat{n}$  axis points towards center of turning circle
  - Draw forces as arrows and place angles as needed
- Find the components of  $F_{net}$ 
  - $F_{net\_parallel} = \text{sum of all parallel components}$
  - $F_{net\_perp} = \text{sum of all perpendicular components}$
  - Remember that  $F_{net\_perp} = mv^2/R (\hat{n})$
- Solve for the unknowns

**Example:** A man of **mass  $m$**  swings from a rope of **length  $L$** . At one particular moment, the rope makes an angle **theta** with the vertical and the man moves with **speed  $v$** . What is the **tension** in the rope?




**CLICKER 2:** The orbit (**radius  $R = 3.8\text{e}8\text{ m}$** ) of the Moon (**mass  $m = 7.3\text{e}22\text{ kg}$** ) is in **uniform circular motion** (constant speed) around the Earth (**mass  $M = 6\text{e}24\text{ kg}$** ). What is the orbital speed of the moon?

- A. 1 km/s
- B. 5 km/s
- C. 10 km/s
- D. 20 km/s
- E. 30 km/s




**SOLUTION:** The orbit (radius  $R = 3.8\text{e}8 \text{ m}$ ) of the Moon (mass  $m = 7.3\text{e}22 \text{ kg}$ ) is in uniform circular motion around the Earth (mass  $M = 6\text{e}24 \text{ kg}$ ). What is the orbital speed of the moon?








Example: A Ferris wheel of **radius  $R$**  rotates **clockwise** at **constant speed  $v$** . You have **mass  $m$**  and are currently in the location shown, at an angle  $\theta$  from the horizontal. Find the parallel and perpendicular components of the **contact force** you feel from the seat.




**Example:** A Ferris wheel of radius  $R$  rotates clockwise at constant speed  $v$ . You have mass  $m$  and are currently in the location shown, at an angle  $\theta$  from the horizontal. Find the parallel and perpendicular components of the contact force you feel from the seat.




CLICKER 3: What does your **FBD** look like when you're at the **TOP** of the Ferris wheel? (ignore the size of the arrows)




**A**



**B**





**C**




**D**


CLICKER 4: What does your **FBD** look like when you're at the **BOTTOM** of the Ferris wheel? (ignore the size of the arrows)




**A**



**B**



**C**




**D**

# Weightlessness



- Your **weight** is caused by Earth's gravity,  $\text{weight} = |F_g| = mg$
- You **FEEL** your weight when you're **in contact** with the ground,  $|F_c| = |F_g|$
- If  $|F_c| < |F_g|$ , then you feel **lighter**
- Similarly, if  $|F_c| > |F_g|$ , then you feel **heavier**
- If there's **no contact force**,  $|F_c| = 0$ , pushing back on you, then you'll **feel weightless**, even though you still have weight!



Example: Two planes fly **parallel to each other** with **speed  $v$**  when they see a blip in the radar up ahead. They split and fly around the whatever-it-is in **vertical half-circles**. **Which pilot feels heavier** in the middle of their half-circle - meaning, at the top for the top plane, or at the bottom for the bottom plane?




**Example:** Two planes fly parallel to each other with speed  $v$  when they see a blip in the radar up ahead. They split and fly around the whatever-it-is in vertical half-circles. Which pilot feels heavier in the middle of their half-circle - meaning, at the top for the top plane, or at the bottom for the bottom plane?



# Curving motion in 3D

- Sometimes you need more than just  $\hat{p}$  and  $\hat{n}$  to describe the motion of an object
- In these situations, you need to use xyz coordinates **in addition to**  $\hat{p}$   $\hat{n}$  coordinates



**Example (“conical pendulum”)** A ball of mass  $m$  is attached to a string and moves in a horizontal circular path. The string has length  $L$  and makes an angle  $\theta$  with the vertical. The ball moves with constant speed, but you don’t know what this speed is. **What is the tension in the string? What is the speed of the ball?**







**Example ("conical pendulum")** A ball of mass  $m$  is attached to a string and moves in a horizontal circular path. The string has length  $L$  and makes an angle  $\theta$  with the vertical. The ball moves with constant speed, but you don't know what this speed is.

What is the tension in the string?



**Example ("conical pendulum")** A ball of mass  $m$  is attached to a string and moves in a horizontal circular path. The string has length  $L$  and makes an angle  $\theta$  with the vertical. The ball moves with constant speed, but you don't know what this speed is.

What is the speed of the ball?




**Example:** An amusement park ride is shaped like a **cylinder of radius  $R$**  that spins. People stand against the inner wall of the cylinder, which has **coefficient of friction  $\mu$** . When the cylinder spins, the floor drops, but yet the people are still “stuck” to the wall. **How fast does the cylinder have to spin to avoid having people slipping and falling off the wall?**




Things we will assume here:



- $\mu$  is the same for every person and corresponds to  $f_{s,\max}$
- The cylinder stays horizontal (no tilt) and spins at a constant rate



**Example:** An amusement park ride is shaped like a cylinder of radius  $R$  that spins. People stand against the inner wall of the cylinder, which has coefficient of friction  $\mu$ . When the cylinder spins, the floor drops, but yet the people are still “stuck” to the wall. How fast does the cylinder have to spin to avoid having people slipping and falling off the wall?



What about when the cylinder is **tilted**?



# Old problems with video solutions

- From Dr Greco!
  - An object moving in a circle along an angled surface
    - <https://vimeo.com/208202487>
  - Riding a sleigh over a hill
    - <https://vimeo.com/158651602>
  - A bug slides off a sphere
    - <https://vimeo.com/158393644>
  - A student riding a ferris wheel
    - <https://vimeo.com/30277476>
  - The three body problem
    - <https://vimeo.com/30277550>

