


## Physics 2211 GPS Week 2

### Problem #1

The position of an object moving along the y-axis, as a function of time, is plotted in the adjacent diagram. For region D, answer the following series of questions. Once you have finished, answer these questions again for A, then C, and then B.



Note: Your instructor may ask you to describe the motion for each region. While doing this, use your hand to represent the object and move it according to the motion information given in that region. Be sure to define where, relative to your hand, the origin is located. The instructor may also ask you to imagine that your camera is taking snapshots of your hand (in equal intervals of time). If you place a dot at the center of your hand in snapshot, what would the pattern of dots together look like.

The object's motion is	D	A	C	B
in the $+y$ direction				
in the $-y$ direction				
changing directions				
stationary				
unable to be determined with the given information				

The object's speed is	D	A	C	B
increasing				
decreasing				
constant				
sometimes increasing and sometimes decreasing				
zero				
unable to be determined with the given information				

The object's displacement within a given region is	D	A	C	B
in the $+y$ direction				
in the $-y$ direction				
changing directions				
zero				
unable to be determined with the given information				

The object's velocity is	D	A	C	B
in the $+y$ direction				
in the $-y$ direction				
changing directions				
zero				
unable to be determined with the given information				

The object's change in velocity is	D	A	C	B
in the $+y$ direction				
in the $-y$ direction				
changing directions				
zero				
unable to be determined with the given information				

## Problem #2

Part 1: A proton traveling with a velocity of  $\langle 700, 700, -700 \rangle$  m/s and is located at  $\langle 13, -4, 17 \rangle$  m at a time  $t = 0.8$  s. Assume the net force on the proton is zero.

(a) Find the velocity of the proton at time  $t = 0.836$  s. (Hint: Use Newton's 2nd Law)

(b) Find the position of the proton at time  $t = 0.836$  s.

(c) How can the proton be moving if the net force on the proton is zero? Explain briefly.

Part 2: A ball of mass  $M = 1$  kg is located at  $\langle 24, -40, 12 \rangle$  m at time  $t = 14.9$  s. Assume the net force on the ball is zero. Find the velocity of the ball and the position of the ball at  $t = 15.1$  s.

Part 3: A ball of mass  $M = 1$  kg is located at  $\langle 24, -40, 12 \rangle$  m at time  $t = 14.9$  s. The ball is located at  $\langle 27, -30, 20 \rangle$  at  $t = 15.0$  s. Assume the net force on the ball is zero. Find the velocity and position of the ball at  $t = 15.1$  s.


### Problem #3

An object's initial position at  $\vec{r}_i$  and final position at  $\vec{r}_f$  are shown on the figure below. Using the numbered direction arrows shown, indicate (by number) which arrow best represents the direction of the quantities listed below. If the quantity has zero magnitude or if more information is needed to determine the direction, indicate using the corresponding number listed below.

The average velocity  $\vec{v}_{avg}$  \_\_\_\_\_

The displacement  $\Delta\vec{r}$  \_\_\_\_\_

The change in velocity  $\Delta\vec{v}$  \_\_\_\_\_



9 zero magnitude  
10 more info needed

An object's initial velocity  $\vec{v}_i$  and final velocity  $\vec{v}_f$  are shown below. Using the numbered direction arrows shown, indicate (by number) which arrow best represents the direction of the quantities listed below. If the quantity has zero magnitude or cannot be determined, indicate using the corresponding number listed below.


The initial position vector  $\vec{r}_i$  \_\_\_\_\_

The change in velocity  $\Delta\vec{v}$  \_\_\_\_\_

The change in momentum  $\Delta\vec{p}$  \_\_\_\_\_

The acceleration  $\vec{a}$  \_\_\_\_\_

The net force  $\vec{F}_{net}$  \_\_\_\_\_




**9 zero magnitude**  
**10 more info needed**

#### Problem #4

To complete this series of questions you will need the Lab 1 starter code. You can find the link to the starter code in the **Lab 1 Instructions** page in Canvas. Copy the contents of the starter code into a new program in your own GlowScript account.

(a) An object moves away from the origin at constant speed. Sketch, by hand, the position vs. time.

(b) For the following position vs. time graph, create a computational model based on Newton's 2nd Law (or, equivalently, the Momentum Principle) that correctly captures the described motion. Use your model to plot both the position vs. time and the velocity vs. time.



(c) An object moves away from the origin at a steady speed for 10s, reverses direction and moves back toward the origin at the same speed for 10s. Sketch, by hand, the position vs. time.