


Physics 2211 GPS Week 9

Problem #1

The diagram shows the path of a spacecraft orbiting a star. You will be asked to rank order various quantities in terms of their values at the locations marked on the path, with the largest first. You can use the symbols “<” and “=”. For example, if you were asked to rank order the locations in terms of their distance from the star: “A < B < C < D < E”



(a) Rank order the locations on the path in terms of the spacecraft's **kinetic energy** at each location, starting with the location where the kinetic energy is the largest.

(b) Consider the system of the spacecraft plus the star. **Which of the following statements are correct?**

- A. As the kinetic energy of the system increases, the gravitational potential energy of the system decreases.
- B. As the spacecraft slows down, the kinetic energy of the system decreases.
- C. As the spacecraft slows down, energy is lost from the system.
- D. External work must be done on the system to speed up the spacecraft.
- E. As the spacecraft's kinetic energy increases, the gravitational potential energy of the system also increases.
- F. Along this path the gravitational potential energy of the system is never zero.
- G. The sum of the kinetic energy of the system plus the gravitational potential energy of the system is a positive number.
- H. The gravitational potential energy of the system is inversely proportional to the square of the distance between the spacecraft and star.
- I. The sum of the kinetic energy of the system plus the gravitational potential energy of the system is the same at every location along this path.
- J. At every location along the spacecraft's path the gravitational potential energy of the system is negative.

Problem #2


In the rough approximation that the density of a planet is uniform throughout its interior, the gravitational field strength (force per unit mass) inside the planet at a distance r from the center is $\frac{GM}{R^3}r$, where M is the mass of the planet and R is the radius of the planet.

- A. Using the uniform-density approximation, calculate the amount of energy required to move an object of mass m from the center of a planet to the surface.
- B. For comparison, how much energy would be required to move the mass from the surface of the planet to a very large distance away?


- C. Imagine that a small hole is drilled through the center of the Earth from one side to the other. Determine the speed of an object of mass m , dropped into this hole, when it reaches the center of the planet. The mass of the Earth is $M = 6 \times 10^{24}$ kg and its radius is $R = 6.4 \times 10^6$ m.

Problem #3


Here is a portion of the elliptical orbit of an asteroid moving around the sun from A to B to C to D .




(a) For the system consisting of the Sun plus the asteroid, graph the gravitational potential energy U , the kinetic energy K , and the sum $K + U$ (total energy), as a function of the separation distance between Sun and asteroid. **Label each curve.** Along the r axis are shown the various distances between Sun and asteroid.




(b) A spacecraft leaves the surface of a planet at exactly the escape speed. For the system consisting of a planet and a spacecraft, graph the gravitational potential energy U , the kinetic energy K , and the sum $K + U$, as a function of the separation distance between planet and the spacecraft. ***Label each curve.***



(c) A spacecraft leaves the surface of a planet with a velocity that is twice the escape speed. For the system consisting of a planet and a spacecraft, graph the gravitational potential energy U , the kinetic energy K , and the sum $K + U$, as a function of the separation distance between planet and the spacecraft. ***Label each curve.***



(d) Two charged particle with opposite charge and identical mass are released from rest a distance R from each other. For the system consisting of the two charges, graph the electric potential energy U , the kinetic energy K , and the sum $K + U$, as a function of the separation distance between the two charges. ***Label each curve.***



(e) Two charged particle with identical charge and mass are released from rest a distance R from each other. For the system consisting of the two charges, graph the electric potential energy U , the kinetic energy K , and the sum $K + U$, as a function of the separation distance between the two charges. ***Label each curve.***

