KADUNA POULTRY DISEASE DIAGNOSTIC LABORATORY
NO. 1, KUBANI CRESCENT, BARNAWA, KADUNA
kadunapoultrylab@gmail.com



CLIENT: A.S. FARMS T/WADA – KADUNA **November 14, 2025**

Case No: KD25/4539

Dear Sir/Ma,
Find below, the result of Necropsy and antibiotic sensitivity test carried out on sample from **bird** submitted for analysis.

History:
Age 24 weeks **Breed/Species**: Commercial Layers **Source of Birds:** **Feed:** Chikun layer pellet **Population:** 300 **Water Source:** Borehole **Housing:** Battery Cage **Major Complaint:** Mortality, drop in production. **Mortality pattern:** 3, 0, 2. **Medication and Vaccination History: Drugs:** Mentoforte + Sawke + Kenflox **Vaccination:** Lasota 10-11-2025.

Necropsy Findings: Empty crops, cachectic bird, hepatic congestion, dehydrated carcass, congested lungs, swollen and congested kidneys, regressed ovarian follicle, necrotic caecal tonsils, mucoid enteritis with watery greenish colour digesta, erosion of the proventricular mucosa, mucoid exudates in the trachea, sinusitis.

Clinical Diagnosis: Systemic Bacterial Infection.
Laboratory Diagnosis: Systemic Bacterial Infection.
Isolate*:* *Citrobacter spp*

SENSITIVE	INTERMEDIATE	RESISTANT
Ciprofloxacin	Penstrept	Tilmicosin
Florfenicol	Gentamicin	
Enrofloxacin	Amoxicillin + Colistin	

Comments and Recommendations:

Based on the PM lesions, culture, and AST, we suggest the following:

1. Administer Enrofloxacin **(Enrobest®)** @ 1ml to 2litres combine with liver tonic **(Hepaklin®)** @ 7ml to 10 litres of drinking water for **5 - 7 consecutive days.**
2. The use of liver tonic should be extended for **14 days.**
3. Ensure that all forms of stress (especially heat stress) are avoided as much as possible.
4. It is of importance to ensure supply of clean and wholesome water by using water sanitizer daily except 2 days before, during and a day after vaccination day(s).

Adhere strictly to other biosecurity measures.

Dr. Temi Ibrahim
08067853699.