

KADUNA POULTRY DISEASE DIAGNOSTIC LABORATORY

NO. 1, KUBANI CRESCENT, BARNAWA, KADUNA kadunapoultrylab@gmail.com

CLIENT: HADIRAF FARMS KADUNA September 8th, 2025

Case No: KD25/3176

Dear Sir/Ma,

Find below, the result of Necropsy and antibiotic sensitivity test carried out on sample from **birds** submitted for analysis.

History:

Age: 10weeks **Breed/Species**: Commercial Layer pullets **Source of Birds:** Agrited **Feed:** Hadiraf feed **Population:** 5,000 **Water Source:** Bore hole **Housing:** Deep litter (saw dust) **Major Complaint:** Mortality, discharge in the eyes, swollen eyes. **Mortality pattern:** 20,18,24. **Medication and Vaccination History: Drugs:** Floricol **Vaccination:** ND + IB @ 4 weeks.

Necropsy Findings: Pasty vents, (5//5), empty crops (5/5), partly autolysed carcasses (2/5), swollen eyes (2/5) with caseous materials (3/5), unilateral femoral head necrosis (1/5), cachectic birds (5/5), hepatic congestion (5/5), congested lungs (5/5), swollen and congested kidneys with mild urates deposits in the ureter (5/5), haemorrhagic ceca tonsils (3/5), mucoid enteritis (5/5), erosion of the proventricular mucosa (3/5), mucoid exudates in the trachea (3/5), caseous material in the sinus (4/5), sinusitis (1/5).

Clinical Diagnosis: Systemic Bacterial Infection, Coccidiosis.
Laboratory Diagnosis: Systemic Bacterial Infection, Coccidiosis.
Microscopy: *Eimeria oocyst* (++)
Isolate: *Citrobacter spp*

SENSITIVE	INTERMEDIATE	RESISTANT
Enrofloxacin	Amoxicillin+colistin	Tilmicosin phosphate
Ciprofloxacin		
Penstrep		
Florfenicol		
Gentamicin		

Comments and Recommendations:

Based on the PM lesions, culture, and AST, we suggest the following:

1. **Day 1-3**; Administer Oregano essential oil or 2.5% Toltrazuril (**Kepcox or Oregostim®**) to bird @ 1ml to 2 litres of drinking water. Ensure to change litter material on the 3rd day of medication use.
2. Administer **20% Enrofloxacin** (e.g **Enrocare-BH/Kattle-care®**) to bird @ 1ml to 2 litres from day 4 to day 8.
3. Ensure water spillage is avoided to prevent coccidiosis.
4. We encourage the use of wood shavings in place of sawdust to avoid fungal infection.
5. It is of importance to ensure daily water sanitization except 2 days before, during and a day after vaccination day(s).

Adhere strictly to other biosecurity measures.

Dr. Muhammad Jibril
07034773648.