

SQUAREFREE SEMIPRIME UNIT FRACTIONS: A CHARACTERIZATION AND A LOCAL LIMIT THEOREM

YUREN TANG

ΤΟΙΣ ΕΜΕ ΦΙΛΟΥΣΙΝ
ΚΑΙ ΟΙΣ ΦΙΛΩ

ABSTRACT. We characterize the positive rationals that are finite sums of reciprocals of distinct squarefree semiprimes, that is, products of two distinct primes. A reduced rational $a/b > 0$ admits such a representation if and only if b is squarefree; moreover, the representation may avoid any prescribed finite set of denominators. This answers Erdős Problem 306 affirmatively.

More quantitatively, fix a reduced $t = a/b \in (0, 1)$ with squarefree b and a fixed $\gamma > 1$ satisfying $t < (\log \gamma)^2/2 < 1$. For a target-adapted semiprime family with natural common denominator L_X , reciprocal-sum variance σ_X^2 , and exact Bernoulli mass W_X , we prove, for every fixed $U > 0$, uniformly for $|j| \leq UL_X\sigma_X$,

$$L_X\sigma_X W_X\left(t + \frac{j}{L_X}\right) = \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{j^2}{2L_X^2\sigma_X^2}\right) + o_U(1).$$

Thus the exact coefficients have a local Gaussian profile. The central coefficient gives the proper-target theorem, and prime dilution yields the full characterization. The same family realizes a growing consecutive block of nearby lattice targets and has exponentially many exact representations at the ambient entropy rate in a mesoscopic typical-cardinality window.

1. INTRODUCTION

An Egyptian-fraction representation of a positive rational number is an expression

$$q = \frac{1}{n_1} + \cdots + \frac{1}{n_k}, \quad 1 < n_1 < \cdots < n_k,$$

and a classical line of questions asks what remains possible when the denominators are arithmetically restricted; for a broad survey of Egyptian-fraction questions and recent developments, see Bloom and Elsholtz [2]. Erdős and Graham asked whether every positive rational a/b with squarefree reduced denominator can be represented using only products of two distinct primes [5, p. 38]. This is now commonly listed as Erdős Problem 306 [1].

The analogous problem with three distinct prime factors has a longer history. Butler, Erdős and Graham proved that every natural number is a sum of distinct

2020 *Mathematics Subject Classification.* Primary 11D68; Secondary 11B34, 60F05.

Key words and phrases. Egyptian fractions, squarefree semiprimes, representation functions, local limit theorem, finite Fourier analysis, Erdős Problem 306.

ORCID: 0009-0006-0847-3330.

unit fractions whose denominators have three distinct prime divisors [3]. For two prime factors, the integer case remained conjectural there. Recently Li proved that every natural number is a sum of distinct unit fractions whose denominators are products of two distinct primes and, for each squarefree b , extended the result to all a/b above an explicit threshold depending on b [7]. The lower range left by this result is the remaining case for a full characterization.

Our first result gives the exact answer.

Theorem 1.1 (Squarefree-denominator characterization). *Let $q = a/b > 0$ be in lowest terms. Then q is a finite sum of reciprocals of pairwise distinct squarefree semiprimes if and only if b is squarefree. More strongly, if $T \subset \mathbb{N}$ is finite, then the representing denominators may be chosen outside T .*

The necessity is immediate: the least common multiple of finitely many squarefree integers is squarefree, and the reduced denominator of their reciprocal sum divides that least common multiple. The content is the converse, including the finite-avoidance statement.

An earlier Lean formalization of the characterization, without the finite-avoidance clause, is available separately [8]. The present paper grew out of that development; finite avoidance and the local-limit and fibre results are additional.

The converse follows from a more quantitative statement. Fix a reduced proper rational

$$t = \frac{a}{b} \in (0, 1)$$

with squarefree denominator b , and fix $\gamma > 1$ such that

$$t < \lambda_\gamma := \frac{(\log \gamma)^2}{2} < 1. \quad (1.1)$$

Such a γ exists for every $0 < t < 1$. Put

$$\begin{aligned} Z &= X^\gamma, \\ \mathcal{P}_X &= \{p \text{ prime} : X \leq p < Z\}, \\ \mathcal{B}_X &= \{q \text{ prime} : Z/2 \leq q < Z\}. \end{aligned} \quad (1.2)$$

Let

$$K_X^* = \lceil (\log Z)^2 \rceil$$

and choose any $\mathcal{B}_X^* \subset \mathcal{B}_X$ with $|\mathcal{B}_X^*| = K_X^*$ once X is large. If S_b is the set of prime divisors of b , define

$$\mathcal{E}_X = \{pq : p, q \in \mathcal{P}_X, p < q\} \sqcup \{rq : r \in S_b, q \in \mathcal{B}_X^*\}. \quad (1.3)$$

For large X , every element of $\mathcal{E}_X$ is a distinct squarefree semiprime and any fixed finite forbidden set is automatically avoided. Write

$$\Lambda_X = \sum_{e \in \mathcal{E}_X} \frac{1}{e}, \quad L_X = b \prod_{p \in \mathcal{P}_X} p, \quad \theta_X = \frac{t}{\Lambda_X}. \quad (1.4)$$

Then

$$\Lambda_X \longrightarrow \lambda_\gamma, \quad \theta_X \longrightarrow \alpha := \frac{2t}{(\log \gamma)^2} \in (0, 1). \quad (1.5)$$

Select each $e \in \mathcal{E}_X$ independently with probability θ_X and set

$$S_X = \sum_{e \in \mathcal{E}_X} \frac{\xi_e}{e}, \quad \sigma_X^2 = \text{Var}(S_X). \quad (1.6)$$

Then $\mathbb{E}S_X = t$. The sparse target rows contribute only $O_b((\log Z)^2/Z^2)$ to the reciprocal square mass, so for every fixed $\gamma > 1$ the complete prime-pair block supplies the leading variance:

$$\sigma_X^2 \sim \frac{\alpha(1-\alpha)}{2X^2 \log^2 X}. \quad (1.7)$$

For lattice targets $s \in L_X^{-1}\mathbb{Z}$, let

$$W_X(s) = \Pr(S_X = s).$$

The Bernoulli-weighted exact coefficients have a local Gaussian profile.

Theorem 1.2 (Local Gaussian profile). *Fix a reduced $t = a/b \in (0, 1)$ with square-free b , and fix $\gamma > 1$ satisfying (1.1). For every fixed $U > 0$, uniformly for integers*

$$|j| \leq UL_X \sigma_X,$$

one has

$$L_X \sigma_X W_X\left(t + \frac{j}{L_X}\right) = \Phi\left(\frac{j}{L_X \sigma_X}\right) + o_U(1), \quad (1.8)$$

where

$$\Phi(u) = \frac{1}{\sqrt{2\pi}} e^{-u^2/2}.$$

In particular,

$$L_X W_X(t) \sim \frac{1}{\sqrt{2\pi} \sigma_X}. \quad (1.9)$$

Thus the exact target coefficient has the size suggested by a lattice local central-limit heuristic, even though the Fourier side lives on the highly composite group $\mathbb{Z}/L_X\mathbb{Z}$ and contains an enormous collection of noncoherent Chinese-remainder configurations.

The central value already gives an avoiding representation of every proper squarefree-denominator target. Theorem 1.1 then follows by a short prime-dilution argument: an arbitrary positive a/b is written as a sum of finitely many identical proper targets with a fresh squarefree denominator.

The same local limit gives a large exact fibre. Put

$$M_X = |\mathcal{E}_X|, \quad \mathcal{R}_X(t) = \left\{ A \subseteq \mathcal{E}_X : \sum_{e \in A} \frac{1}{e} = t \right\}, \quad (1.10)$$

and let $H(u) = -u \log u - (1-u) \log(1-u)$. Set

$$w_X = \left\lceil 2\sqrt{M_X \log L_X} \right\rceil. \quad (1.11)$$

Theorem 1.3 (Multiplicity and typical cardinality). *With t, γ fixed as above,*

$$\#\{A \in \mathcal{R}_X(t) : |A| - \theta_X M_X \leq w_X\} \geq \exp([H(\alpha) - o(1)]M_X). \quad (1.12)$$

Consequently

$$\log |\mathcal{R}_X(t)| \geq [H(\alpha) - o(1)]M_X. \quad (1.13)$$

Moreover, for some integer

$$k_X = \theta_X M_X + O(w_X),$$

one has

$$\#\{A \in \mathcal{R}_X(t) : |A| = k_X\} \geq \exp([H(\alpha) - o(1)]M_X). \quad (1.14)$$

The exponent $H(\alpha)$ is the ambient binomial entropy exponent near the typical Bernoulli cardinality. At exponential scale, the exact reciprocal-sum constraint therefore costs only the subexponential lattice factor already visible in the local limit, within this mesoscopic cardinality window. For $0 < t < 1/2$, one may choose $\gamma = e^{2\sqrt{t}}$ so that $\alpha = 1/2$ and the rate is the full binary entropy $\log 2$; Section 7 records this as a short corollary.

The moving part of Theorem 1.2 gives a second form of abundance. Since $L_X \sigma_X \rightarrow \infty$ while $\sigma_X \rightarrow 0$, one fixed family $\mathcal{E}_X$ realizes a consecutive block of

$$2\lfloor UL_X \sigma_X \rfloor + 1$$

nearby lattice targets, whose number tends to infinity although the corresponding real interval has radius $U\sigma_X \rightarrow 0$.

1.1. Fourier architecture. Character orthogonality gives

$$L_X W_X(t) = \sum_{h \bmod L_X} F_X(h), \quad (1.15)$$

where

$$F_X(h) = e(-th) \prod_{e \in \mathcal{E}_X} ((1 - \theta_X) + \theta_X e(h/e)), \quad e(u) = e^{2\pi i u}. \quad (1.16)$$

For $h = m \pmod{L_X}$, Taylor expansion gives

$$F_X(m) = \exp(-2\pi^2 m^2 \sigma_X^2 + o(1))$$

uniformly on each fixed standardized interval $|m| \leq C/\sigma_X$. The exact centring identity $\theta_X \Lambda_X = t$ cancels the linear phase. If only these frequencies mattered, Theorem 1.2 would be a routine Gaussian Riemann-sum calculation.

The difficulty is that L_X contains every prime in $\mathcal{P}_X$. A general frequency has one CRT coordinate modulo each of these primes, and many noncoherent assignments can make large subsets of the Bernoulli factors nearly undamped. The proof uses the geometry of the denominator family to compress this space.

The top block $\mathcal{B}_X$ acts as an *anchor*. Denominators qq' with $q, q' \in \mathcal{B}_X$ measure an internal quadratic energy. We prove that sufficiently low anchor energy forces every anchor coordinate to be the reduction of one and the same integer m , with no exceptional coordinates. Above the forcing threshold, a fingerprint argument controls the complete weighted population of anchor assignments.

Once the anchor label m is known, the complete cross factors rq with $q \in \mathcal{B}_X$ and $r \in \mathcal{P}_X \setminus \mathcal{B}_X$ observe every remaining prime coordinate against the common anchor. The fixed coordinates $r \mid b$ need much less: any K_X^* top primes already separate their residue states, so only the sparse rows rq with $q \in \mathcal{B}_X^*$ are retained. Factors not used for decoding are kept and later suppress decoded labels outside the Gaussian range.

The sparse target rows illustrate a useful separation of roles. Their reciprocal mass, square mass, and cardinality are all negligible compared with the complete pair family, yet because b is fixed they give superpolynomially small row tails and exact residue identification on the major arc. A vanishing-mass correction can therefore restore arithmetic observability without changing the limiting Gaussian law.

The Fourier sum therefore has three conceptual pieces:

- (i) a Gaussian core $h = m \pmod{L_X}$ with $|m| = O(\sigma_X^{-1})$;

- (ii) decoded-label tails, suppressed either by the complete-pair variance or by retained complete-pair damping;
- (iii) an energetic/off-minimizer remainder, whose total weighted mass is exponentially small.

The technical proof subdivides the decoded-label tails at two transition scales, but these are subordinate to the three-part structure above.

The moving-target statement requires no new minor-frequency analysis. Replacing t by $t + j/L_X$ simply multiplies (1.16) by the unit character

$$e(-hj/L_X).$$

Every modulus estimate is unchanged. On the integer frequencies this factor becomes the Fourier variable in the Gaussian Riemann sum and yields the profile Φ .

1.2. Exact equality and finite avoidance. The Fourier calculation first detects a subset whose integer weights sum to $L_X t$ modulo L_X . Exact equality is then deterministic: every subset reciprocal sum lies in $[0, \Lambda_X]$ with $\Lambda_X < 1$, and so does the proper target t . Two such numbers congruent modulo one must be equal.

The analytic argument therefore gives every proper squarefree-denominator target directly. To reach an arbitrary positive rational a/b , choose a fresh prime $p > a$ with $p \nmid b$ and apply the proper-target theorem p times to $a/(bp)$ on pairwise disjoint supports. Their union represents a/b . The quantitative theorems are deliberately not transferred through this closure: Theorems 1.2 and 1.3 concern one target-adapted direct family.

1.3. Related restricted-denominator work. The present problem belongs to a wider literature on Egyptian fractions with restricted prime support. For explicit short representations of 1 by semiprime denominators, see Watanabe [9]. Different restrictions, in which all prime factors are drawn from a fixed finite set, are studied by Czenky, McGovern, Plavnik, Rowell and Watkins [4]. These directions are complementary to the expanding squarefree-semiprime family used here.

1.4. Organization. Section 2 constructs the direct family and records its reciprocal mass, variance, and lattice scale. Section 3 gives the finite Fourier formula and separates the no-wrap exactness step. Section 4 proves one-anchor rigidity, including the zero-exception and fingerprint estimates. Section 5 establishes row observability, decoder identification, weighted row compression, and retained damping. Section 6 evaluates the Fourier coefficient on the local-limit scale. Section 7 derives multiplicity and entropy consequences, Section 8 proves the moving-target profile, Section 9 completes the characterization and finite-avoidance consequences, and Section 10 records the exact scope and limitations of the method.

2. THE DIRECT SEMIPRIME FAMILY

Throughout Sections 2–8, fix a reduced rational

$$t = \frac{a}{b} \in (0, 1),$$

with squarefree denominator b , a finite set $T \subset \mathbb{N}$, and a real number γ satisfying

$$\gamma > 1, \quad t < \lambda_\gamma := \frac{(\log \gamma)^2}{2} < 1. \quad (2.1)$$

All implicit constants may depend on the fixed data t, γ, b , unless explicitly stated otherwise, and T is fixed before $X \rightarrow \infty$.

Set

$$Z = X^\gamma, \quad \mathcal{P} = \{p \text{ prime} : X \leq p < Z\}, \quad \mathcal{B} = \{q \text{ prime} : Z/2 \leq q < Z\}. \quad (2.2)$$

Since $\gamma > 1$, one has $\mathcal{B} \subset \mathcal{P}$ for all sufficiently large X . Let S_b denote the set of prime divisors of b . Put

$$K_* = \lceil (\log Z)^2 \rceil,$$

and, for all sufficiently large X , choose any subset

$$\mathcal{B}_* \subset \mathcal{B}, \quad |\mathcal{B}_*| = K_*.$$

No distributional property of $\mathcal{B}_*$ will be used beyond its cardinality. Define

$$\mathcal{E}_{\text{pair}} = \{pq : p, q \in \mathcal{P}, p < q\}, \quad \mathcal{E}_b = \{rq : r \in S_b, q \in \mathcal{B}_*\}, \quad (2.3)$$

$$\mathcal{E} = \mathcal{E}_{\text{pair}} \sqcup \mathcal{E}_b, \quad L = b \prod_{p \in \mathcal{P}} p. \quad (2.4)$$

For sufficiently large X , every prime in $\mathcal{P}$ exceeds every prime divisor of b , and every denominator in (2.3) lies outside T .

Lemma 2.1 (Arithmetic shape). *For all sufficiently large X :*

- (i) every $e \in \mathcal{E}$ is the product of two distinct primes;
- (ii) the elements of $\mathcal{E}$ are pairwise distinct and lie outside T ;
- (iii) every $e \in \mathcal{E}$ divides L ;
- (iv) L is squarefree.

Proof. The two factors of every denominator in (2.3) are distinct once X exceeds the prime divisors of b . Unique factorization separates products of two primes from $\mathcal{P}$ from products rq with $r \mid b$ and $q \in \mathcal{B}_*$, and also proves distinctness inside each family. The choice of X gives avoidance of T . Since b is squarefree and disjoint in prime support from $\mathcal{P}$, the integer L is squarefree and every denominator in (2.3) divides it. $\square$

2.1. Reciprocal load and variance. Write

$$H_{\mathcal{P}} = \sum_{p \in \mathcal{P}} \frac{1}{p}, \quad U_{\mathcal{P}} = \sum_{p \in \mathcal{P}} \frac{1}{p^2}.$$

The prime number theorem and partial summation give

$$H_{\mathcal{P}} = \log \gamma + o(1), \quad U_{\mathcal{P}} \sim \frac{1}{X \log X}. \quad (2.5)$$

Consequently

$$\sum_{e \in \mathcal{E}_{\text{pair}}} \frac{1}{e} = \frac{1}{2}(H_{\mathcal{P}}^2 - U_{\mathcal{P}}) \longrightarrow \lambda_\gamma = \frac{(\log \gamma)^2}{2}. \quad (2.6)$$

Since every $q \in \mathcal{B}_*$ is comparable with Z , the sparse target rows satisfy

$$\sum_{e \in \mathcal{E}_b} \frac{1}{e} = O_b\left(\frac{K_*}{Z}\right) = o(1). \quad (2.7)$$

Thus, with

$$\Lambda = \sum_{e \in \mathcal{E}} \frac{1}{e}, \quad (2.8)$$

one has

$$\Lambda \longrightarrow \lambda_\gamma. \quad (2.9)$$

Fix constants

$$t < \lambda_- < \lambda_\gamma < \lambda_+ < 1. \quad (2.10)$$

For all sufficiently large X , $\Lambda \in (\lambda_-, \lambda_+)$. Define

$$\theta = \frac{t}{\Lambda}. \quad (2.11)$$

Then θ remains in a fixed compact subinterval of $(0, 1)$ and

$$\theta \Lambda = t. \quad (2.12)$$

The complete-pair identity gives

$$\sum_{e \in \mathcal{E}_{\text{pair}}} \frac{1}{e^2} = \frac{1}{2} \left(U_{\mathcal{P}}^2 - \sum_{p \in \mathcal{P}} \frac{1}{p^4} \right) \sim \frac{1}{2X^2 \log^2 X}. \quad (2.13)$$

The sparse target-row square mass is

$$\sum_{e \in \mathcal{E}_b} \frac{1}{e^2} = O_b\left(\frac{K_*}{Z^2}\right). \quad (2.14)$$

Because $Z = X^\gamma$, $K_* = O((\log Z)^2)$, and $\gamma > 1$, the ratio of (2.14) to the pair square mass is

$$O_b\left(K_* \frac{X^2 \log^2 X}{Z^2}\right) = X^{2-2\gamma} \text{polylog } X \longrightarrow 0.$$

Thus the complete prime-pair block supplies the leading variance throughout the present regime.

Let $(\xi_e)_{e \in \mathcal{E}}$ be independent Bernoulli variables with parameter θ , and set

$$S = \sum_{e \in \mathcal{E}} \frac{\xi_e}{e}. \quad (2.15)$$

Then, by (2.12),

$$\mathbb{E}S = t, \quad (2.16)$$

and

$$\sigma^2 := \text{Var}(S) = \theta(1 - \theta) \sum_{e \in \mathcal{E}} \frac{1}{e^2}. \quad (2.17)$$

Since

$$\theta \longrightarrow \alpha := \alpha_{t,\gamma} := \frac{t}{\lambda_\gamma} = \frac{2t}{(\log \gamma)^2} \in (0, 1), \quad (2.18)$$

we obtain

$$\sigma^2 \sim \frac{\alpha(1 - \alpha)}{2X^2 \log^2 X}, \quad \sigma^{-1} \asymp_{t,\gamma} X \log X. \quad (2.19)$$

2.2. Family size and lattice scale.

Lemma 2.2. *As $X \rightarrow \infty$,*

$$M := |\mathcal{E}| = (1 + o(1)) \frac{Z^2}{2 \log^2 Z}, \quad (2.20)$$

$$\log L = (1 + o(1))Z = o(M), \quad (2.21)$$

and

$$L\sigma \longrightarrow \infty. \quad (2.22)$$

Proof. The prime number theorem gives

$$|\mathcal{P}| = (1 + o(1)) \frac{Z}{\log Z},$$

so

$$|\mathcal{E}_{\text{pair}}| = \binom{|\mathcal{P}|}{2} = (1 + o(1)) \frac{Z^2}{2 \log^2 Z}.$$

The sparse target rows have only $O_b(K_*) = O_b((\log Z)^2)$ elements, which is negligible at this scale. This proves (2.20).

By the prime number theorem in the form $\vartheta(x) \sim x$,

$$\log L = O_b(1) + \sum_{X \leq p < Z} \log p = (1 + o(1))Z,$$

which gives (2.21). Finally (2.19) and the exponential size of L imply (2.22). $\square$

Because every $e \in \mathcal{E}$ divides L and $Lt = aL/b$ is an integer, the random variable LS is integer-valued and the exact event

$$S = t \quad (2.23)$$

is a single lattice atom. Sections 3–6 determine its asymptotic mass.

3. FOURIER INVERSION AND EXACTNESS

For a subset $A \subseteq \mathcal{E}$, give A the Bernoulli weight

$$w(A) = \theta^{|A|} (1 - \theta)^{M - |A|}. \quad (3.1)$$

Let P be the total weight of exact representations of the fixed target $t = a/b$. We first work in the finite group $\mathbb{Z}/L\mathbb{Z}$.

3.1. The target coefficient. Let P^{quot} be the probability of the congruence

$$\sum_{e \in \mathcal{E}} \xi_e \frac{L}{e} \equiv Lt = \frac{aL}{b} \pmod{L}. \quad (3.2)$$

Character orthogonality gives

$$LP^{\text{quot}} = \sum_{h \bmod L} F(h), \quad (3.3)$$

where

$$F(h) = e(-th) \prod_{e \in \mathcal{E}} ((1 - \theta) + \theta e(h/e)), \quad e(u) = e^{2\pi i u}. \quad (3.4)$$

We will use the following modulus estimate repeatedly.

Lemma 3.1 (Bernoulli modulus bound). *There is $\kappa > 0$, depending only on the fixed parameters t and γ , such that for every real u ,*

$$|(1 - \theta) + \theta e(u)| \leq \exp(-\kappa \|u\|^2), \quad (3.5)$$

where $\|u\|$ denotes the distance from u to the nearest integer.

Proof. We have

$$|(1 - \theta) + \theta e(u)|^2 = 1 - 4\theta(1 - \theta) \sin^2(\pi u).$$

By Section 2, θ lies in a fixed compact subinterval of $(0, 1)$, while $\sin(\pi x) \geq 2x$ for $0 \leq x \leq 1/2$. Thus

$$4\theta(1 - \theta) \sin^2(\pi u) \geq c \|u\|^2.$$

Using $1 - v \leq e^{-v}$ and adjusting the constant after taking square roots proves (3.5). $\square$

In particular,

$$|F(h)| \leq \exp\left(-\kappa \sum_{e \in \mathcal{E}} \|h/e\|^2\right). \quad (3.6)$$

The target character has modulus one and therefore plays no role in the modulus estimates. We must understand which CRT configurations make a large collection of phases h/e simultaneously close to integers.

3.2. CRT coordinates and the factor partition. Since L is squarefree, CRT identifies a frequency $h \bmod L$ with one residue coordinate modulo every prime dividing L . The denominator family naturally separates these coordinates.

The primes in $\mathcal{B}$ are the *anchor coordinates*. Denominators qq' with $q, q' \in \mathcal{B}$ measure their internal coherence. Once the anchor is coherent, denominators rq with $q \in \mathcal{B}$ and $r \in \mathcal{P} \setminus \mathcal{B}$ observe the remaining prime rows against the common anchor, while the sparse denominators rq with $q \in \mathcal{B}_*$ and $r \in S_b$ observe the fixed target coordinates. Denominators involving no anchor prime are deliberately left unused by the decoder and will later provide extra damping.

Every denominator in $\mathcal{E}$ belongs to exactly one of the classes

- (i) qq' with $q, q' \in \mathcal{B}$: anchor–anchor factors;
- (ii) rq with $q \in \mathcal{B}, r \in \mathcal{P} \setminus \mathcal{B}$: lower rows;
- (iii) rq with $q \in \mathcal{B}_*, r \in S_b$: sparse target rows;
- (iv) pp' with $p, p' \in \mathcal{P} \setminus \mathcal{B}$: retained lower–lower factors.

This exact partition is used in Section 5: classes (i)–(iii) recover the coherent CRT state where exact identification is needed, while class (iv) suppresses decoded labels after the decoder has done its work.

3.3. Integer frequency classes. Every integer m determines the CRT assignment obtained by reducing m modulo each prime dividing L . When $|m| < e/2$ for every $e \in \mathcal{E}$,

$$\left\| \frac{m}{e} \right\| = \frac{|m|}{e}. \quad (3.7)$$

Moreover the linear phase in the logarithm of (3.4) cancels exactly because

$$\theta \sum_{e \in \mathcal{E}} \frac{1}{e} = t. \quad (3.8)$$

This produces the Gaussian main term in Section 6. Before the Taylor expansion can control the full Fourier sum, however, we must show that every sufficiently low-energy CRT configuration is not merely close to an integer assignment but is exactly one on the coordinates required by the major arc. Section 4 proves exact coherence for the anchor block and Section 5 identifies the remaining coordinates on their respective decoder ranges.

3.4. No-wrap as a separate exactness step. The Fourier calculation takes place in $\mathbb{Z}/L\mathbb{Z}$, whereas the theorem is an equality in $\mathbb{Q}$. We keep these two assertions logically separate.

Lemma 3.2 (No-wrap). *For all sufficiently large X , so that $\Lambda < 1$, let $s \in L^{-1}\mathbb{Z} \cap [0, \Lambda]$. If $A \subseteq \mathcal{E}$ satisfies*

$$\sum_{e \in A} \frac{L}{e} \equiv Ls \pmod{L}, \quad (3.9)$$

then

$$\sum_{e \in A} \frac{1}{e} = s. \quad (3.10)$$

Proof. After division by L ,

$$D = \sum_{e \in A} \frac{1}{e} - s$$

is an integer. Both terms lie in $[0, \Lambda]$, and $\Lambda < 1$, so $-1 < D < 1$. Hence $D = 0$. $\square$

Taking $s = t$ gives

$$P^{\text{quot}} = P. \quad (3.11)$$

We nevertheless retain the quotient notation during the Fourier estimates to emphasize that positivity in the finite group and exactness in the fundamental interval are different inputs.

The same lemma applies verbatim to moving targets. If

$$t_j = t + \frac{j}{L}$$

lies in $[0, \Lambda]$, then a subset sum congruent to t_j modulo one is equal to t_j . For every fixed standardized radius U , this condition holds for all $|j| \leq UL\sigma$ once X is sufficiently large, because $\sigma \rightarrow 0$ and t lies strictly inside the limiting interval $[0, \lambda_\gamma]$.

4. ONE-ANCHOR RIGIDITY

We isolate the combinatorial statement used on the top block. Let $Y \rightarrow \infty$, and let Q be a set of primes in $[Y, 2Y)$ satisfying

$$c \frac{Y}{\log Y} \leq N := |Q| \leq C \frac{Y}{\log Y}. \quad (4.1)$$

An assignment is a tuple $a = (a_p)_{p \in Q}$ with $a_p \in \mathbb{Z}/p\mathbb{Z}$. For distinct $p, q \in Q$, let $H_{pq}(a)$ be the centred simultaneous CRT lift

$$H_{pq} \equiv a_p \pmod{p}, \quad H_{pq} \equiv a_q \pmod{q}, \quad -\frac{pq}{2} < H_{pq} \leq \frac{pq}{2}. \quad (4.2)$$

Define

$$Q(a) = \sum_{p < q \in Q} \left(\frac{H_{pq}(a)}{pq} \right)^2, \quad \sigma_Q^2 = \sum_{p < q \in Q} \frac{1}{p^2 q^2}. \quad (4.3)$$

Then

$$\sigma_Q^2 \asymp \frac{1}{Y^2 \log^2 Y}. \quad (4.4)$$

The main conclusion of this section is that energy below the scale $Y/\log^3 Y$ forces an exact integer state on every coordinate. The logic has two complementary parts. Low energy first forces one dominant integer label, and reciprocal dispersion then eliminates every exceptional coordinate. Above the rigidity threshold, a fingerprint determines almost all coordinates; the entropy of its completions is small enough to be absorbed by the Bernoulli damping. Thus the first mechanism identifies every low-energy top assignment by an exact integer label, while the second makes the energetic remainder negligible in weighted sum.

4.1. Reciprocal dispersion.

Lemma 4.1 (Reciprocal dispersion). *Let q be prime with $Y \leq q < 4Y$, let F be a set of $s \geq 16$ primes in $[Y, 2Y)$, none equal to q , and let $d \not\equiv 0 \pmod{q}$. Then at least $s/2$ primes $p \in F$ satisfy*

$$\left\| \frac{dp^{-1}}{q} \right\| \geq \frac{s}{64Y}.$$

Consequently

$$\sum_{p \in F} \left\| \frac{dp^{-1}}{q} \right\|^2 \geq c_D \frac{s^3}{Y^2} \quad (4.5)$$

for an absolute constant $c_D > 0$.

Proof. Put $\delta = s/(64Y)$. If

$$\left\| \frac{dp^{-1}}{q} \right\| \leq \delta,$$

choose the nearest integer ℓ and write

$$v = dp^{-1} - \ell q.$$

Then

$$|v| \leq \delta q \leq s/16, \quad vp \equiv d \pmod{q}.$$

For a fixed nonzero integer v , the last congruence places p in one residue class modulo q . Since $[Y, 2Y)$ has length at most q , such a class contains at most two integers. There are at most $2(s/16) + 1$ possible values of v . Hence fewer than $s/2$ primes can satisfy the displayed small-distance condition. At least $s/2$ primes therefore have distance at least δ , and summing their squares proves (4.5). $\square$

The hypothesis $s \geq 16$ will be retained explicitly whenever Lemma 4.1 is used.

4.2. Repulsion between label classes. Fix $\rho \in (0, 1/4)$. An integer m is called a *dominant label* for a if $|m| \leq Y^2/2$ and

$$a_p \equiv m \pmod{p}$$

for at least $(1 - \rho)N$ primes $p \in Q$. Two dominant labels are equal: their classes overlap in at least two primes, whose product divides their difference, while the difference has absolute value at most Y^2 .

Lemma 4.2 (Cross-label energy). *Let $C_m, C_{m'} \subseteq Q$ be disjoint label classes with $m \neq m'$, and suppose*

$$\begin{aligned} |m|, |m'| &\leq B_0 < Y^2/4, \\ |C_m| &\geq \max\left(16, \frac{128B_0}{Y}\right), \quad |C_{m'}| \geq 2. \end{aligned} \quad (4.6)$$

Then

$$\sum_{p \in C_m} \sum_{q \in C_{m'}} \left(\frac{H_{pq}}{pq} \right)^2 \geq c_E \frac{|C_m|^3 (|C_{m'}| - 1)}{Y^2}. \quad (4.7)$$

Proof. Put $d = m' - m$. Since $0 < |d| < Y^2/2$, at most one prime in $C_{m'}$ can divide d . Fix any other $q \in C_{m'}$. For $p \in C_m$, write

$$H_{pq} = m + jp.$$

Reducing modulo q gives $jp \equiv d \pmod{q}$, and therefore

$$\left\| \frac{dp^{-1}}{q} \right\| \leq \frac{|H_{pq}|}{pq} + \frac{|m|}{pq}. \quad (4.8)$$

By Lemma 4.1, for at least half the primes $p \in C_m$ the left side is at least $|C_m|/(64Y)$. The size condition in (4.6) gives

$$\frac{|m|}{pq} \leq \frac{B_0}{Y^2} \leq \frac{|C_m|}{128Y}.$$

Thus at least half the p satisfy

$$\frac{|H_{pq}|}{pq} \geq \frac{|C_m|}{128Y}.$$

The energy contributed by this fixed q is $\gg |C_m|^3/Y^2$. Summing over all but at most one $q \in C_{m'}$ gives (4.7). $\square$

The factor $|C_{m'}| - 1$ in (4.7) is important and will be kept when the class energies are summed.

4.3. Low energy forces a dominant label.

Proposition 4.3 (Dominant-label forcing). *There is $c_w > 0$ such that, for all sufficiently large Y ,*

$$Q(a) < c_w \frac{Y}{\log^3 Y} \quad (4.9)$$

implies that a has a unique dominant label.

Proof. If $Q(a) = 0$, every centred pair lift vanishes, so every coordinate is the residue of the integer 0. We may therefore assume

$$0 < R := Q(a) < c \frac{Y}{\log^3 Y}, \quad (4.10)$$

with c to be chosen sufficiently small.

Set

$$B_0 = A_\rho \frac{\sqrt{R} Y^2}{N}, \quad (4.11)$$

where A_ρ is a large fixed constant. Call an edge $\{p, q\}$ bad if $|H_{pq}| > B_0$. Since $pq < 4Y^2$, every bad edge contributes more than $B_0^2/(16Y^4)$ to R , so

$$\#\{\text{bad edges}\} \leq \frac{16RY^4}{B_0^2}. \quad (4.12)$$

Some base prime p_0 has bad degree at most

$$\frac{32RY^4}{B_0^2 N} = \frac{32N}{A_\rho^2}. \quad (4.13)$$

Choose A_ρ so that this is at most $\rho N/8$.

For every nonbad neighbour q of p_0 , set

$$\ell(q) = H_{p_0 q}.$$

Then $|\ell(q)| \leq B_0$, $\ell(q) \equiv a_q \pmod{q}$, and all $\ell(q)$ lie in one residue class modulo p_0 . For every integer label m that occurs, define its label class by

$$C_m = \{q : \{p_0, q\} \text{ is nonbad and } \ell(q) = m\}.$$

These classes partition the nonbad neighbours of p_0 , and every $q \in C_m$ satisfies $a_q \equiv m \pmod{q}$. The number M_0 of possible integer labels therefore satisfies

$$M_0 \leq \frac{2B_0}{Y} + 2. \quad (4.14)$$

For sufficiently small c and large Y , (4.10)–(4.11) also give $B_0 < Y^2/4$.

Assume for contradiction that no class is dominant. Call a label class small if its size is less than

$$s_0 = 256 \left(\frac{B_0}{Y} + 1 \right). \quad (4.15)$$

Then $s_0 \geq 16$ and $s_0 \geq 128B_0/Y$, so Lemma 4.2 applies to every substantial class.

Suppose first that the small classes contain at least $\rho N/4$ vertices. Then

$$\frac{\rho N}{4} \leq M_0 s_0 \leq C_\rho \left(\frac{B_0}{Y} + 1 \right)^2. \quad (4.16)$$

If $B_0/Y < 1$, the right side is bounded while $N \rightarrow \infty$, a contradiction. Otherwise (4.11) and (4.16) imply

$$N \leq C_\rho \frac{B_0^2}{Y^2} = C_\rho A_\rho^2 \frac{RY^2}{N^2},$$

and hence

$$R \gg \frac{N^3}{Y^2} \gg \frac{Y}{\log^3 Y},$$

contradicting (4.10) for sufficiently small c .

We may therefore assume that the small classes contain fewer than $\rho N/4$ vertices. After removing p_0 and its bad neighbours, the substantial classes have total size

$$S \geq (1 - \rho/2)N \quad (4.17)$$

for large Y . Let their sizes be $n_1, \dots, n_J$. Each $n_i \geq s_0$, while no n_i exceeds $(1 - \rho)N$.

Apply Lemma 4.2 to ordered pairs of distinct substantial classes. Every undirected cross edge is counted twice, so, after absorbing the factor $1/2$ into the constant,

$$R \geq \frac{c}{Y^2} \sum_i n_i^3 \sum_{j \neq i} (n_j - 1). \quad (4.18)$$

The inner sum is exactly

$$\sum_{j \neq i} (n_j - 1) = (S - n_i) - (J - 1). \quad (4.19)$$

Because every other substantial class has size at least s_0 ,

$$S - n_i \geq (J - 1)s_0,$$

and hence

$$(S - n_i) - (J - 1) \geq \left(1 - \frac{1}{s_0}\right) (S - n_i) \geq \frac{255}{256} (S - n_i). \quad (4.20)$$

Thus

$$R \geq \frac{c}{Y^2} \sum_i n_i^3 (S - n_i). \quad (4.21)$$

If the largest class has size at least $S/2$, then it contains $\gg N$ vertices while (4.17) and non-dominance leave $\gg_\rho N$ substantial vertices outside it. Its single term in (4.21) is therefore $\gg_\rho N^4/Y^2$, too large for (4.10).

If every class has size less than $S/2$, then $S - n_i \geq S/2$, and power mean gives

$$\sum_i n_i^3 \geq \frac{S^3}{J^2} \geq \frac{S^3}{M_0^2}.$$

Hence

$$R \geq c_\rho \frac{N^4}{M_0^2 Y^2}. \quad (4.22)$$

If $B_0/Y < 1$, this again exceeds the scale in (4.10). Otherwise, by (4.14),

$$M_0^2 \leq C \frac{B_0^2}{Y^2} = C A_\rho^2 \frac{R Y^2}{N^2}.$$

Substitution into (4.22) yields

$$R^2 \geq c \frac{N^6}{Y^4}, \quad R \geq c \frac{N^3}{Y^2} \gg \frac{Y}{\log^3 Y},$$

again a contradiction. Choosing c_w below the finitely many constants above proves the proposition. $\square$

4.4. No exceptional coordinates.

Proposition 4.4 (Exact zero-exception rigidity). *After decreasing c_w if necessary, every assignment satisfying (4.9) is represented by one integer m :*

$$a_p \equiv m \pmod{p} \quad (p \in Q). \quad (4.23)$$

Moreover

$$|m| \leq C_\rho \frac{\sqrt{Q(a)}}{\sigma_Q}. \quad (4.24)$$

Proof. By Proposition 4.3, there is a dominant class $C \subseteq Q$ with label m . For $p, q \in C$, the centred lift is exactly m . Hence

$$Q(a) \geq m^2 \sum_{p < q \in C} \frac{1}{p^2 q^2} \geq c_\rho m^2 \sigma_Q^2,$$

which proves (4.24).

Let $E_0 = Q \setminus C$. The label bound and the low-energy hypothesis imply, for large Y ,

$$|m| \leq \frac{|C|Y}{128}. \quad (4.25)$$

For each $q \in E_0$, the residue difference $d = a_q - m$ is nonzero modulo q . The same CRT subtraction as in Lemma 4.2, followed by Lemma 4.1, shows that the edges from q into C contribute at least

$$c \frac{|C|^3}{Y^2}$$

to $Q(a)$. These edge sets are disjoint as q varies. Therefore

$$|E_0| c \frac{|C|^3}{Y^2} \leq Q(a). \quad (4.26)$$

Since $|C| \gg Y/\log Y$,

$$|E_0| \leq C \frac{Q(a)Y^2}{N^3} < C' c_w. \quad (4.27)$$

Choose c_w so that $C' c_w < 1$. The quantity $|E_0|$ is an integer, hence $E_0 = \emptyset$. $\square$

The last integer step is the reason the argument gives exact coherence rather than merely a density-one label class.

4.5. Fingerprint entropy. Above the forcing threshold we need a weighted count of anchor assignments. A small fingerprint controls the possible completions.

Lemma 4.5 (Fingerprint rigidity). *Fix $F \subset Q$ with $|F| = s \geq 16$ and fix the residues on F . For $q \in Q \setminus F$ and $w \pmod{q}$, put*

$$t_q(w) = \sum_{p \in F} \left(\frac{H_{pq}(a_p, w)}{pq} \right)^2. \quad (4.28)$$

For a sufficiently small absolute $g > 0$, at most one residue $w \pmod{q}$ satisfies

$$t_q(w) < g \frac{s^3}{Y^2}. \quad (4.29)$$

Proof. If $w \neq w'$, the difference of the corresponding CRT lifts is divisible by p and reduces modulo q to $w - w'$. Hence

$$\left\| \frac{(w - w')p^{-1}}{q} \right\| \leq \frac{|H_{pq}(a_p, w)|}{pq} + \frac{|H_{pq}(a_p, w')|}{pq}.$$

Squaring, using $(u + v)^2 \leq 2u^2 + 2v^2$, and summing over $p \in F$ gives

$$\sum_{p \in F} \left\| \frac{(w - w')p^{-1}}{q} \right\|^2 \leq 2t_q(w) + 2t_q(w').$$

On the other hand, Lemma 4.1 gives the lower bound $c_D s^3 / Y^2$. If both residues satisfied (4.29), we would obtain

$$c_D \frac{s^3}{Y^2} < 4g \frac{s^3}{Y^2},$$

which is impossible once $4g < c_D$. Thus at most one such residue exists. $\square$

Proposition 4.6 (Fingerprint entropy). *For every $\varepsilon > 0$, all sufficiently large blocks satisfy*

$$\#\{a : Q(a) \leq R\} \leq e^{\varepsilon R} \quad (4.30)$$

uniformly for

$$R \geq c_w \frac{Y}{\log^3 Y}. \quad (4.31)$$

Proof. First suppose

$$R \leq c_* \frac{N^4}{Y^2}, \quad (4.32)$$

where $c_* > 0$ is sufficiently small. Choose

$$s = \lceil A(RY^2)^{1/4} \rceil \quad (4.33)$$

with A a fixed large constant, and use a predetermined fingerprint set F of size s . The lower bound (4.31) gives $s \geq 16$ for large Y , while (4.32) gives $s \leq N/2$ after c_* is fixed small enough.

There are at most $(2Y)^s$ assignments on F . For a complete assignment,

$$\sum_{q \notin F} t_q(a_q) \leq R.$$

By Lemma 4.5, all but at most

$$t \leq \frac{RY^2}{gs^3} \leq C_A s \quad (4.34)$$

coordinates outside F are forced to a unique residue. Choosing the exceptional coordinates and their residues contributes at most

$$\sum_{j \leq t} \binom{N}{j} (2Y)^j \leq \exp(C(s + t) \log Y).$$

Thus the logarithm of the total count is

$$O((RY^2)^{1/4} \log Y). \quad (4.35)$$

After division by R , the worst case occurs at the forcing floor (4.31), where (4.35) gives

$$O(Y^{-1/4} \log^{13/4} Y) \longrightarrow 0. \quad (4.36)$$

This is uniform in the range (4.32).

If instead $R > c_* N^4 / Y^2$, the trivial number of assignments is at most $(2Y)^N$, whose logarithm is $O(Y)$, while

$$R \gg \frac{N^4}{Y^2} \gg \frac{Y^2}{\log^4 Y}.$$

Thus again $O(Y) \leq \varepsilon R$ for large Y . $\square$

4.6. The weighted top partition. We apply the preceding statements to the top block $\mathcal{B} \subset [Z/2, Z)$, taking $Y = Z/2$. Define

$$Q_{\mathcal{B}}(a_{\mathcal{B}}) = \sum_{q < q' \in \mathcal{B}} \left(\frac{H_{qq'}(a_{\mathcal{B}})}{qq'} \right)^2, \quad \sigma_{\mathcal{B},0}^2 = \sum_{q < q' \in \mathcal{B}} \frac{1}{q^2 q'^2}. \quad (4.37)$$

Then

$$\sigma_{\mathcal{B},0}^2 \asymp \frac{1}{Z^2 \log^2 Z}. \quad (4.38)$$

There is a forcing floor

$$F_{\mathcal{B}} = c_{\mathcal{B}} \frac{Z}{\log^3 Z} \quad (4.39)$$

such that every assignment with $Q_{\mathcal{B}} < F_{\mathcal{B}}$ has a unique exact integer label m . Proposition 4.4 gives

$$|m| \ll \frac{\sqrt{F_{\mathcal{B}}}}{\sigma_{\mathcal{B},0}} = O\left(\frac{Z^{3/2}}{\sqrt{\log Z}}\right) = o(Z^2). \quad (4.40)$$

Thus m is the centred lift on every internal top pair and

$$Q_{\mathcal{B}} = m^2 \sigma_{\mathcal{B},0}^2. \quad (4.41)$$

Let

$$T_{\mathcal{B}}(a_{\mathcal{B}}) = \prod_{q < q' \in \mathcal{B}} |(1 - \theta) + \theta e(H_{qq'}/(qq'))|. \quad (4.42)$$

Lemma 3.1 gives

$$T_{\mathcal{B}}(a_{\mathcal{B}}) \leq \exp(-\kappa Q_{\mathcal{B}}(a_{\mathcal{B}})). \quad (4.43)$$

Choose $\varepsilon < \kappa/2$ in Proposition 4.6. For $k \geq \lfloor F_{\mathcal{B}} \rfloor$, the shell $k \leq Q_{\mathcal{B}} < k + 1$ contains at most $e^{\varepsilon(k+1)}$ assignments, while (4.43) bounds the weight of each by $e^{-\kappa k}$. Hence the geometric sum over the energetic shells gives

$$\sum_{Q_{\mathcal{B}} \geq F_{\mathcal{B}}} T_{\mathcal{B}}(a_{\mathcal{B}}) \ll \sum_{k \geq \lfloor F_{\mathcal{B}} \rfloor} e^{\varepsilon(k+1) - \kappa k} \ll \exp(-c F_{\mathcal{B}}). \quad (4.44)$$

We call assignments with $Q_{\mathcal{B}} < F_{\mathcal{B}}$ *low-energy decoded*; Proposition 4.4 supplies their exact integer labels. Assignments with $Q_{\mathcal{B}} \geq F_{\mathcal{B}}$ will simply be called *energetic*. No converse between integer coherence and this energy split is asserted.

Below the floor, the exact label determines the assignment, since $a_q \equiv m \pmod{q}$ for every $q \in \mathcal{B}$. Therefore

$$\sum_{Q_{\mathcal{B}} < F_{\mathcal{B}}} T_{\mathcal{B}}(a_{\mathcal{B}}) \leq \sum_{m \in \mathbb{Z}} e^{-\kappa m^2 \sigma_{\mathcal{B},0}^2} \ll \sigma_{\mathcal{B},0}^{-1} \ll Z \log Z. \quad (4.45)$$

Hence the complete weighted top partition satisfies

$$P_{\text{top}} := \sum_{a_{\mathcal{B}}} T_{\mathcal{B}}(a_{\mathcal{B}}) \ll Z \log Z, \quad (4.46)$$

while the contribution of assignments with $Q_{\mathcal{B}} \geq F_{\mathcal{B}}$ is exponentially small by (4.44).

The distinction between raw cardinality and weighted mass in (4.46) is essential. Section 5 will sum row fibres only after multiplying by this anchor weight.

5. ROW OBSERVABILITY AND DECODED-SKELETON COMPRESSION

Let

$$\mathcal{R} = (\mathcal{P} \setminus \mathcal{B}) \cup S_b. \quad (5.1)$$

For each row prime put

$$\mathcal{B}_r = \begin{cases} \mathcal{B}, & r \in \mathcal{P} \setminus \mathcal{B}, \\ \mathcal{B}_*, & r \in S_b. \end{cases}$$

Thus the lower prime rows retain the complete anchor block, whereas the finitely many target coordinates use only the sparse sensor set. Fix an anchor assignment on $\mathcal{B}$. For $r \in \mathcal{R}$, a row residue $x \in \mathbb{Z}/r\mathbb{Z}$, and $q \in \mathcal{B}_r$, let

$$u_{rq}(x) \in \mathbb{R}/\mathbb{Z}$$

be the phase $h/(rq)$ determined by $x \pmod{r}$ and the fixed anchor residue modulo q . If the row residue is changed by d , CRT gives

$$u_{rq}(x+d) - u_{rq}(x) \equiv \frac{dq^{-1}}{r} \pmod{1}. \quad (5.2)$$

Thus row observability is controlled by inverse residues on the circle.

5.1. Row code distance.

Lemma 5.1 (Cyclic energy with multiplicity). *Let r be prime and let $c_1, \dots, c_M$ be nonzero residues modulo r , each occurring at most μ times. If M/μ is sufficiently large, then for every $d \not\equiv 0 \pmod{r}$,*

$$\sum_{j=1}^M \left\| \frac{dc_j}{r} \right\|^2 \geq c \frac{M^3}{\mu^2 r^2}. \quad (5.3)$$

Proof. Multiplication by d permutes the nonzero residue classes. Put

$$s = \left\lfloor \frac{M}{4\mu} \right\rfloor.$$

For large M/μ , $s \geq M/(8\mu) \geq 1$. Fewer than $2\mu s \leq M/2$ entries can lie among the $2s$ nonzero classes nearest zero. Hence at least $M/2$ terms have circle distance at least $M/(8\mu r)$, proving (5.3). $\square$

Let $M_{\mathcal{B}} = |\mathcal{B}|$. Then

$$M_{\mathcal{B}} \asymp \frac{Z}{\log Z}. \quad (5.4)$$

Proposition 5.2 (Lower-prime row distance). *For every prime $r \in \mathcal{P} \setminus \mathcal{B}$,*

$$D_r := \min_{d \not\equiv 0 \pmod r} \sum_{q \in \mathcal{B}} \left\| \frac{dq^{-1}}{r} \right\|^2 \gg \frac{Z}{\log^3 Z}, \quad (5.5)$$

uniformly in r .

Proof. Inversion permutes the nonzero residue classes modulo r . Since $\mathcal{B} \subset [Z/2, Z)$, one residue class modulo r is represented at most

$$\mu_r \leq \frac{Z}{2r} + 1 \leq \frac{2Z}{r}$$

times. Lemma 5.1 and (5.4) give

$$D_r \gg \frac{M_{\mathcal{B}}^3}{\mu_r^2 r^2} \gg \frac{Z^3 / \log^3 Z}{Z^2} = \frac{Z}{\log^3 Z}.$$

Moreover $M_{\mathcal{B}} / \mu_r \gg r / \log Z \geq X / \log Z \rightarrow \infty$, so the largeness condition is uniform. $\square$

Proposition 5.3 (Sparse target-row distance). *For every prime $r \mid b$,*

$$D_r := \min_{d \not\equiv 0 \pmod r} \sum_{q \in \mathcal{B}_*} \left\| \frac{dq^{-1}}{r} \right\|^2 \geq \frac{K_*}{r^2} \gg_b (\log Z)^2. \quad (5.6)$$

Proof. Every $q \in \mathcal{B}_*$ is coprime to r . If $d \not\equiv 0 \pmod r$, then $dq^{-1} \not\equiv 0 \pmod r$, so

$$\left\| \frac{dq^{-1}}{r} \right\| \geq \frac{1}{r}.$$

Summing over the K_* retained sensors gives the result. No distribution of the primes in $\mathcal{B}_*$ is used. $\square$

The target rows are therefore sparse in mass but strong in observability. Their only required population property is $K_* \rightarrow \infty$ fast enough that the row-tail loss remains superpolynomially small.

5.2. Row concentration. For $r \in \mathcal{R}$, define the complex row product

$$B_r(x) = \prod_{q \in \mathcal{B}_r} ((1 - \theta) + \theta e(u_{rq}(x))),$$

the row energy

$$E_r(x) = \sum_{q \in \mathcal{B}_r} \|u_{rq}(x)\|^2, \quad (5.7)$$

and the row modulus

$$A_r(x) = |B_r(x)| = \prod_{q \in \mathcal{B}_r} |(1 - \theta) + \theta e(u_{rq}(x))|. \quad (5.8)$$

Choose a minimizer x_r^* of E_r .

Lemma 5.4 (Row tail). *There is $c > 0$, depending only on the fixed data, such that*

$$\delta_r := \sum_{x \neq x_r^*} A_r(x) \leq r \exp(-c D_r). \quad (5.9)$$

Proof. Lemma 3.1 gives

$$A_r(x) \leq e^{-\kappa E_r(x)}.$$

For $x \neq x_r^*$, the squared distance between the two phase vectors is at least D_r . The circle triangle inequality and $(u+v)^2 \leq 2u^2 + 2v^2$ give

$$D_r \leq 2E_r(x) + 2E_r(x_r^*) \leq 4E_r(x),$$

since x_r^* minimizes the energy. Thus $E_r(x) \geq D_r/4$, and summing over at most r nonminimal states proves (5.9). $\square$

Propositions 5.2 and 5.3 imply

$$\Delta := \sum_{r \in \mathcal{R}} \delta_r \leq Z^2 \exp\left(-c \frac{Z}{\log^3 Z}\right) + O_b\left(e^{-c_b (\log Z)^2}\right) = o(1). \quad (5.10)$$

In particular $Z^A \Delta \rightarrow 0$ for every fixed $A > 0$.

5.3. Weighted row compression. Fix an anchor assignment. Let $C_{\mathcal{B}}$ be the complex product of all anchor–anchor Bernoulli factors, so $|C_{\mathcal{B}}| = T_{\mathcal{B}}$. Let $G_{\text{ret}}(x_{\mathcal{R}})$ be the complex product of all retained lower–lower factors. For every row state set

$$\omega_r(x) = \begin{cases} B_r(x)/A_r(x), & A_r(x) > 0, \\ 0, & A_r(x) = 0. \end{cases}$$

Then $B_r(x) = A_r(x)\omega_r(x)$ and $|\omega_r(x)| \leq 1$. For the fixed anchor assignment and a fixed row tuple $x_{\mathcal{R}}$, CRT determines a unique frequency $h \pmod{L}$. Hence the target character $e(-th)$ is a function of this tuple and may be absorbed, together with the row phases and retained product, into

$$G(x_{\mathcal{R}}) = e(-th) G_{\text{ret}}(x_{\mathcal{R}}) \prod_{r \in \mathcal{R}} \omega_r(x_r).$$

Thus

$$|G(x_{\mathcal{R}})| \leq 1. \quad (5.11)$$

By the exact factor partition of Section 3, the full complex Fourier contribution of the lower coordinates above this fixed anchor assignment is exactly

$$C_{\mathcal{B}} \sum_{x_{\mathcal{R}}} \prod_{r \in \mathcal{R}} A_r(x_r) G(x_{\mathcal{R}}).$$

No denominator is omitted or counted twice, and no complex row phase has been discarded.

Lemma 5.5 (Weighted row compression). *Suppose finitely many nonnegative row kernels have distinguished states x_r^* and satisfy*

$$\sum_{x_r \neq x_r^*} A_r(x_r) \leq \delta_r, \quad |G| \leq 1.$$

Then

$$\sum_{x_{\mathcal{R}}} \prod_r A_r(x_r) G(x_{\mathcal{R}}) = \prod_r A_r(x_r^*) G(x^*) + E_{\text{fib}}, \quad (5.12)$$

where

$$|E_{\text{fib}}| \leq \prod_r (A_r(x_r^*) + \delta_r) - \prod_r A_r(x_r^*) \leq \prod_r (1 + \delta_r) - 1 \leq e^{\Delta} - 1. \quad (5.13)$$

The same statement holds for an empty or singleton row set under the usual empty-product convention.

Proof. Separate the distinguished point $x^* = (x_r^*)_r$. Off that point, discard G in absolute value. The pure product sum factors:

$$\sum_{x_R} \prod_r A_r(x_r) = \prod_r \sum_{x_r} A_r(x_r) \leq \prod_r (A_r(x_r^*) + \delta_r).$$

Subtract the distinguished-point pure product and add back its actual value with $G(x^*)$. This gives the first inequality in (5.13).

Writing $a_r = A_r(x_r^*) \in [0, 1]$, the function

$$H((a_r)) = \prod_r (a_r + \delta_r) - \prod_r a_r$$

is nondecreasing in every a_r . Therefore

$$H((a_r)) \leq H((1)) = \prod_r (1 + \delta_r) - 1,$$

and $1 + u \leq e^u$ gives the final bound. $\square$

Multiplying the error in (5.13) by $|C_{\mathcal{B}}| = T_{\mathcal{B}}$ and summing over anchor assignments, the total off-minimizer contribution is

$$\leq P_{\text{top}}(e^{\Delta} - 1) \ll Z \log Z (e^{\Delta} - 1) = o(1). \quad (5.14)$$

The raw number of anchor assignments never enters the estimate.

5.4. Decoder identification. Suppose the anchor assignment has $Q_{\mathcal{B}} < F_{\mathcal{B}}$, and let m be its exact integer label supplied by Proposition 4.4. The candidate row state $m \pmod{r}$ makes every cross phase congruent to $m/(rq)$, and hence

$$E_r(m \pmod{r}) \leq \frac{m^2}{r^2} \sum_{q \in \mathcal{B}_r} \frac{1}{q^2}. \quad (5.15)$$

If this is less than $D_r/8$, then it is the unique minimizer: for any other residue x ,

$$D_r \leq 2E_r(x) + 2E_r(m),$$

so

$$E_r(x) \geq \frac{D_r}{2} - E_r(m) > \frac{3D_r}{8} > E_r(m). \quad (5.16)$$

Put

$$M_{\text{dec}} = \frac{XZ}{(\log Z)^2}. \quad (5.17)$$

Proposition 5.6 (Lower-prime decoder). *For every $r \in \mathcal{P} \setminus \mathcal{B}$ and every low-energy anchor label with $|m| \leq M_{\text{dec}}$,*

$$x_r^* = m \pmod{r}. \quad (5.18)$$

Proof. For a lower prime row, (5.15) gives

$$E_r(m) \ll \frac{m^2/r^2}{Z \log Z}.$$

This is largest at $r = X$. At $|m| = M_{\text{dec}}$,

$$E_r(m) \ll \frac{M_{\text{dec}}^2/X^2}{Z \log Z} = \frac{Z}{\log^5 Z}.$$

By Proposition 5.2, $D_r \gg Z/\log^3 Z$, so the ratio tends uniformly to zero. Apply (5.16). $\square$

Put $M_{\text{tar}} = Z/\log Z$.

Proposition 5.7 (Target-coordinate decoder). *For every prime $r \mid b$ and every low-energy anchor label with $|m| \leq M_{\text{tar}}$,*

$$x_r^* = m \pmod{r}. \quad (5.19)$$

Proof. For a target coordinate, (5.15) and $q \asymp Z$ on $\mathcal{B}_*$ give

$$E_r(m) \ll_b \frac{m^2 K_*}{Z^2}.$$

Proposition 5.3 gives $D_r \geq K_*/r^2 \gg_b K_*$. Hence

$$\frac{E_r(m)}{D_r} \ll_b \frac{m^2}{Z^2} \leq \frac{1}{\log^2 Z} \rightarrow 0.$$

Apply (5.16). $\square$

Since $\gamma > 1$, (2.19) implies

$$\sigma^{-1} \asymp X \log X = o(M_{\text{tar}}). \quad (5.20)$$

Since b is squarefree, agreement modulo every prime divisor of b is agreement modulo b . Thus every low-energy label in any fixed standardized Gaussian range $|m| \leq C/\sigma$ produces the frequency class $h = m \pmod{L}$ on the decoded skeleton for all sufficiently large X .

For target coordinates no decoder identification is asserted beyond M_{tar} . By contrast, Proposition 5.6 still identifies every prime coordinate in $\mathcal{P}$ throughout

$$X^2/4 < |m| \leq M_{\text{dec}}. \quad (5.21)$$

This weaker conclusion is enough for the retained damping below.

5.5. Retained lower–lower damping.

Proposition 5.8 (Adaptive retained damping). *For every integer m satisfying (5.21), the interval*

$$I_m = [2\sqrt{|m|}, 3\sqrt{|m|}] \quad (5.22)$$

lies in $[X, Z/2)$, contains

$$K_m \gg \frac{\sqrt{|m|}}{\log |m|} \quad (5.23)$$

primes, and the retained lower–lower factors on the distinguished skeleton contribute energy

$$Q_{\text{ret}}(m) \gg \frac{|m|}{(\log |m|)^2}. \quad (5.24)$$

Proof. Write $y = \sqrt{|m|}$. From $|m| > X^2/4$ we have $2y > X$. From $|m| \leq M_{\text{dec}}$,

$$3y \leq \frac{3\sqrt{XZ}}{\log Z} = o(Z),$$

because $Z = X^\gamma$ with $\gamma > 1$. Hence $3y < Z/2$ for large X , and $I_m \subset \mathcal{P} \setminus \mathcal{B}$. The prime number theorem in fixed-ratio intervals gives (5.23).

For distinct primes $p, q \in I_m$,

$$4|m| \leq pq \leq 9|m|,$$

so

$$\frac{1}{9} \leq \frac{|m|}{pq} \leq \frac{1}{4}.$$

Hence the circle distance of $m/(pq)$ is at least $1/9$. There are $\gg K_m^2$ retained pairs, all with prime coordinates decoded to m . Summing their squared distances proves (5.24). $\square$

By Lemma 3.1, if h_m^* denotes the distinguished row-minimizer state above a low-energy label m in the range (5.21), then

$$|F(h_m^*)| \leq \exp\left(-c \frac{|m|}{(\log |m|)^2}\right). \quad (5.25)$$

Only the prime-coordinate decoder is used here; the fixed coordinates of b may already have ceased to decode. Beyond M_{dec} no decoder identification is asserted. The anchor has converted low internal energy into one exact label, the row sensors identify the coordinates on the ranges where they are needed, Lemma 5.5 compresses the off-minimizer states without a cardinality loss, and the unused complete-pair factors suppress decoded labels after full target decoding has ceased.

6. LOCAL GAUSSIAN ANALYSIS

Recall from (3.3) that

$$LP^{\text{quot}} = \sum_{h \bmod L} F(h). \quad (6.1)$$

We now evaluate this coefficient on its natural scale $1/\sigma$.

Fix $C \geq 1$. The limit $X \rightarrow \infty$ will be taken with C fixed, and only afterwards will $C \rightarrow \infty$. Put

$$N = \left\lfloor \frac{C}{\sigma} \right\rfloor. \quad (6.2)$$

Since $\sigma^{-1} \asymp_{t,\gamma} X \log X$ and $\gamma > 1$, for fixed C and large X ,

$$N < \min\left(M_{\text{tar}}, \frac{X^2}{4}\right), \quad \frac{X^2}{4} < M_{\text{dec}}. \quad (6.3)$$

For orientation, the invariant scale relations used below are

$$\sigma^{-1} \ll X^2, \quad \sigma^{-1} \ll M_{\text{tar}}, \quad X^2 \ll M_{\text{dec}}, \quad \frac{Z^{3/2}}{\sqrt{\log Z}} = o(Z^2).$$

No fixed ordering between M_{dec} and the low-energy anchor-label ceiling $Z^{3/2}/\sqrt{\log Z}$ is needed. If the former exceeds the latter, the large decoded-label range below is simply empty.

By the terminology of Section 4, a top assignment is either low-energy decoded or energetic. Each low-energy decoded assignment has a unique exact

integer label m , lying in exactly one of the ranges

$$\begin{aligned} \text{I.} \quad & |m| \leq N, \\ \text{II.} \quad & N < |m| \leq X^2/4, \\ \text{III.} \quad & X^2/4 < |m| \leq M_{\text{dec}}, \\ \text{IV.} \quad & |m| > M_{\text{dec}}. \end{aligned} \tag{6.4}$$

Let $\mathcal{M}_<$ be the set of labels arising from low-energy decoded top assignments. By (4.45), the label determines the top assignment. For $m \in \mathcal{M}_<$ let h_m^* be its distinguished row-minimizer state; for an energetic anchor assignment $a_{\mathcal{B}}$ let $h^*(a_{\mathcal{B}})$ be the corresponding distinguished state. Finally let $\mathcal{E}_{\text{off}}$ be the exact sum of all Fourier terms whose row tuple is not distinguished. Lemma 5.5 then gives the exact disjoint decomposition

$$\begin{aligned} \sum_{h \bmod L} F(h) &= \sum_{\substack{m \in \mathcal{M}_< \\ |m| \leq N}} F(h_m^*) + \sum_{\substack{m \in \mathcal{M}_< \\ N < |m| \leq X^2/4}} F(h_m^*) \\ &+ \sum_{\substack{m \in \mathcal{M}_< \\ X^2/4 < |m| \leq M_{\text{dec}}}} F(h_m^*) + \sum_{\substack{m \in \mathcal{M}_< \\ |m| > M_{\text{dec}}}} F(h_m^*) \\ &+ \sum_{a_{\mathcal{B}}: Q_{\mathcal{B}}(a_{\mathcal{B}}) \geq F_{\mathcal{B}}} F(h^*(a_{\mathcal{B}})) + \mathcal{E}_{\text{off}}. \end{aligned}$$

Sections 6.1–6.4 treat the first four terms, while Section 6.5 treats the last two. In Sections 6.1–6.4, sums over m are understood to run over $m \in \mathcal{M}_<$. In particular, no classification of energetic assignments by integer coherence is required.

6.1. The Gaussian core. Every integer m with $|m| \leq N$ belongs to $\mathcal{M}_<$ for all sufficiently large X . Indeed, $|m| = O_C(X \log X) = o(Z^2)$, so m is the centred lift on every internal top pair, and by (2.19), (4.38), and (4.39),

$$Q_{\mathcal{B}} = m^2 \sigma_{\mathcal{B},0}^2 \ll_C \frac{X^2}{Z^2} = o(F_{\mathcal{B}}).$$

Propositions 5.6 and 5.7 therefore identify its distinguished state with the frequency $h = m \pmod{L}$.

Choosing $\eta_0 > 0$ sufficiently small, use the analytic branch of $\log$ near 1, normalized by $\log 1 = 0$. Since θ stays in a compact subinterval of $(0, 1)$, there is a constant $C_3 < \infty$ such that, uniformly for $|z| \leq \eta_0$,

$$\log((1 - \theta) + \theta e(z)) = 2\pi i \theta z - 2\pi^2 \theta (1 - \theta) z^2 + R_{\theta}(z), \tag{6.5}$$

with

$$|R_{\theta}(z)| \leq C_3 |z|^3. \tag{6.6}$$

The pair denominators are at least X^2 , while the sparse target denominators are comparable with Z up to constants depending on b . Uniformly for $|m| \leq C/\sigma$,

$$\max_{e \in \mathcal{E}_{\text{pair}}} \frac{|m|}{e} \ll_C \frac{\log X}{X} \rightarrow 0, \quad \max_{e \in \mathcal{E}_b} \frac{|m|}{e} \ll_{C,b} \frac{X \log X}{Z} \rightarrow 0. \tag{6.7}$$

For the aggregate cubic remainder, the pair family gives

$$\sum_{e \in \mathcal{E}_{\text{pair}}} \left| \frac{m}{e} \right|^3 \ll_C \frac{\log X}{X},$$

while the sparse target family gives

$$\sum_{e \in \mathcal{E}_b} \left| \frac{m}{e} \right|^3 \ll_{C,b} K_* \frac{X^3 \log^3 X}{Z^3} = o(1). \quad (6.8)$$

The last limit uses $K_* = O((\log Z)^2)$ and fixed $\gamma > 1$. Thus the aggregate remainder in (6.5) is $o_C(1)$ uniformly throughout the major range.

The linear term cancels exactly by (2.12):

$$2\pi i m \theta \sum_{e \in \mathcal{E}} \frac{1}{e} - 2\pi i m t = 0. \quad (6.9)$$

The quadratic term is $-2\pi^2 m^2 \sigma^2$. Therefore

$$F(m) = \exp(-2\pi^2 m^2 \sigma^2 + \varepsilon_{X,m}), \quad \max_{|m| \leq N} |\varepsilon_{X,m}| \longrightarrow 0 \quad (6.10)$$

for every fixed C .

Proposition 6.1 (Normalized major arc). *For every fixed $C > 0$,*

$$\sigma \sum_{|m| \leq C/\sigma} F(m) \longrightarrow \int_{-C}^C e^{-2\pi^2 u^2} du. \quad (6.11)$$

Proof. Put $u_m = m\sigma$. The mesh is $\sigma \rightarrow 0$, and (6.10) is uniform on $[-C, C]$. Thus the left side is a Riemann sum for the integral in (6.11). $\square$

The Gaussian core is therefore of order $1/\sigma$. We compare every remaining contribution with this same scale.

6.2. The first decoded-label tail. For

$$N < |m| \leq X^2/4, \quad (6.12)$$

Proposition 5.6 identifies every prime coordinate in $\mathcal{P}$ with m ; the fixed coordinates of b need not decode in this whole range. Let h_m^* be the distinguished row-minimizer state above the low-energy anchor label m . On every complete-pair factor its phase is $m/(pq)$, and since $pq \geq X^2$,

$$\left\| \frac{m}{pq} \right\| = \frac{|m|}{pq} \leq \frac{1}{4}.$$

Discarding the sparse target factors in modulus, using that the pair square mass is asymptotic to the full square mass, and recalling that $\theta(1 - \theta) \asymp 1$, Lemma 3.1 gives

$$|F(h_m^*)| \leq e^{-cm^2 \sigma^2}. \quad (6.13)$$

A Gaussian integral comparison yields

$$\sum_{|m| > C/\sigma} e^{-cm^2 \sigma^2} \leq K \sigma^{-1} e^{-cC^2/2}. \quad (6.14)$$

Hence

$$\limsup_{X \rightarrow \infty} \sigma \sum_{N < |m| \leq X^2/4} |F(h_m^*)| \leq K e^{-cC^2/2}. \quad (6.15)$$

This is the only complementary range whose normalized bound need not tend to zero at fixed C ; it disappears when $C \rightarrow \infty$. Full target-coordinate decoding is not used in this modulus-only tail.

6.3. The intermediate decoded-label tail. Suppose

$$X^2/4 < |m| \leq M_{\text{dec}}. \quad (6.16)$$

All prime coordinates still decode to m . By (5.25),

$$|F(h_m^*)| \leq \exp\left(-c \frac{|m|}{(\log |m|)^2}\right). \quad (6.17)$$

Throughout (6.16), $\log |m| \asymp \log X$. There are at most $2M_{\text{dec}} + 1 = O(XZ)$ labels, so

$$\sum_{X^2/4 < |m| \leq M_{\text{dec}}} |F(h_m^*)| \ll XZ \exp\left(-c \frac{X^2}{\log^2 X}\right) = o(\sigma^{-1}). \quad (6.18)$$

Here and below fixed powers of X are harmless because γ is fixed.

6.4. Large decoded anchor labels. For a low-energy top label m with $|m| > M_{\text{dec}}$, let h_m^* denote the single distinguished row-minimizer state above that anchor assignment. No decoder identification is asserted in this range. The internal anchor factor alone gives

$$|F(h_m^*)| \leq \exp(-\kappa m^2 \sigma_{\mathcal{B},0}^2),$$

while every row and residual factor has modulus at most one. Therefore

$$\sum_{|m| > M_{\text{dec}}} |F(h_m^*)| \ll \sigma_{\mathcal{B},0}^{-1} \exp\left(-c M_{\text{dec}}^2 \sigma_{\mathcal{B},0}^2\right). \quad (6.19)$$

If no low-energy anchor label reaches this range, the sum is empty and the same bound is trivial. Otherwise, from (4.38) and (5.17),

$$M_{\text{dec}}^2 \sigma_{\mathcal{B},0}^2 \gg \frac{X^2}{\log^6 Z}, \quad (6.20)$$

while $\sigma_{\mathcal{B},0}^{-1} = O(Z \log Z)$. Hence

$$\sum_{|m| > M_{\text{dec}}} |F(h_m^*)| = o(\sigma^{-1}). \quad (6.21)$$

6.5. Energetic anchors and off-minimizer row states. By (4.44), and since all factors outside the anchor have modulus at most one,

$$\left| \sum_{a_{\mathcal{B}}: Q_{\mathcal{B}}(a_{\mathcal{B}}) \geq F_{\mathcal{B}}} F(h^*(a_{\mathcal{B}})) \right| = O(e^{-cF_{\mathcal{B}}}). \quad (6.22)$$

This is exactly the energetic distinguished-state term in the decomposition above. Independently, (5.14) controls all off-minimizer row states over every anchor assignment:

$$|\mathcal{E}_{\text{off}}| = O\left(Z \log Z \left[Z^2 e^{-cZ/\log^3 Z} + e^{-c_b(\log Z)^2} \right]\right). \quad (6.23)$$

Thus no row fibre is omitted from the assembly. Both quantities are $o(\sigma^{-1})$, because $\sigma^{-1} \asymp X \log X$ and the displayed errors decay faster than every fixed power of Z .

6.6. Assembly and the local limit. Let $\mathfrak{R}_{X,C}$ be the Fourier contribution remaining after removing the frequencies $h = m \pmod{L}$ with $|m| \leq C/\sigma$. Equations (6.15), (6.18), (6.21), (6.22), and (6.23) give

$$\limsup_{X \rightarrow \infty} \sigma |\mathfrak{R}_{X,C}| \leq K e^{-cC^2/2}. \quad (6.24)$$

Combining this with Proposition 6.1,

$$\limsup_{X \rightarrow \infty} \left| \sigma LP^{\text{quot}} - \int_{-C}^C e^{-2\pi^2 u^2} du \right| \leq K e^{-cC^2/2}. \quad (6.25)$$

Now let $C \rightarrow \infty$. Since

$$\int_{\mathbb{R}} e^{-2\pi^2 u^2} du = \frac{1}{\sqrt{2\pi}}, \quad (6.26)$$

we obtain

$$\sigma LP^{\text{quot}} \longrightarrow \frac{1}{\sqrt{2\pi}}. \quad (6.27)$$

By Lemma 3.2, $P^{\text{quot}} = P$. Therefore

$$LP \sim \frac{1}{\sqrt{2\pi} \sigma}. \quad (6.28)$$

This is the central value of Theorem 1.2.

The order of limits is part of the argument:

$$X \rightarrow \infty \quad \text{at fixed } C, \quad \text{then} \quad C \rightarrow \infty. \quad (6.29)$$

No diagonal choice $C = C(X) \rightarrow \infty$ is asserted.

Corollary 6.2 (Quantitative positivity). *There is $c > 0$ such that, for all sufficiently large X ,*

$$LP \geq \frac{c}{\sigma}. \quad (6.30)$$

In particular $P > 0$.

The asymptotic (6.28), rather than mere positivity, drives the entropy-scale conclusions of Section 7. The full moving-target form of Theorem 1.2 is proved in Section 8 by inserting one unit character into the same analysis.

7. MULTIPLICITY AND TYPICAL CARDINALITY

Retain the fixed proper target $t = a/b$, exponent γ , and direct family $\mathcal{E} = \mathcal{E}_X$. Put

$$M = |\mathcal{E}|, \quad \mathcal{R}_X(t) = \left\{ A \subseteq \mathcal{E} : \sum_{e \in A} \frac{1}{e} = t \right\}. \quad (7.1)$$

By Lemma 2.2 and (2.19),

$$\log L = o(M), \quad |\log \sigma| = o(M). \quad (7.2)$$

7.1. The typical-cardinality window. Let

$$K = \sum_{e \in \mathcal{E}} \xi_e$$

be the number of selected denominators under the ambient Bernoulli law, and set

$$w = \left\lceil 2\sqrt{M \log L} \right\rceil. \quad (7.3)$$

Since $\log L = o(M)$,

$$\sqrt{M} \ll w = o(M). \quad (7.4)$$

Hoeffding's inequality [6] gives

$$\Pr(|K - \theta M| > w) \leq 2 \exp\left(-\frac{2w^2}{M}\right) \leq 2L^{-8}. \quad (7.5)$$

The local limit (6.28) implies

$$2L^{-8} = o(P), \quad (7.6)$$

where $P = W_X(t)$ is the exact-target Bernoulli mass. Consequently the exact target mass in the window satisfies

$$\sum_{\substack{A \in \mathcal{R}_X(t) \\ ||A| - \theta M| \leq w}} \theta^{|A|} (1 - \theta)^{M - |A|} = (1 - o(1))P. \quad (7.7)$$

If $|k - \theta M| \leq w$, write $k = \theta M + \delta$. Then

$$\log(\theta^k (1 - \theta)^{M - k}) = -MH(\theta) + \delta \log \frac{\theta}{1 - \theta}, \quad (7.8)$$

where

$$H(u) = -u \log u - (1 - u) \log(1 - u).$$

Since θ remains in a compact subinterval of $(0, 1)$,

$$\theta^k (1 - \theta)^{M - k} \leq \exp(-MH(\theta) + O(w)) \quad (7.9)$$

uniformly in the window. Dividing (7.7) by this uniform upper bound for an atom in the window and using

$$\log P = -\log L - \log \sigma + O(1) = o(M), \quad w = o(M), \quad H(\theta) \rightarrow H(\alpha),$$

gives

$$\#\{A \in \mathcal{R}_X(t) : ||A| - \theta M| \leq w\} \geq \exp([H(\alpha) - o(1)]M), \quad (7.10)$$

where

$$\alpha = \frac{2t}{(\log \gamma)^2}. \quad (7.11)$$

In particular,

$$\log |\mathcal{R}_X(t)| \geq [H(\alpha) - o(1)]M. \quad (7.12)$$

The exponent in (7.10) is optimal at the ambient exponential scale inside the same cardinality window: the standard entropy estimate for binomial coefficients gives

$$\#\{A \subseteq \mathcal{E} : ||A| - \theta M| \leq w\} \leq \exp([H(\alpha) + o(1)]M). \quad (7.13)$$

Thus imposing the exact reciprocal-sum constraint costs only a subexponential factor within this mesoscopic window. We do not claim a matching upper bound for the total exact fibre outside the window.

7.2. One exact cardinality. The interval $|k - \theta M| \leq w$ contains at most $2w + 1 = \exp(o(M))$ integers. By pigeonhole applied to (7.10), there is an integer

$$k_X = \theta M + O(w) \quad (7.14)$$

such that

$$\#\{A \in \mathcal{R}_X(t) : |A| = k_X\} \geq \exp([H(\alpha) - o(1)]M). \quad (7.15)$$

This completes Theorem 1.3.

Corollary 7.1 (Entropy saturation for small proper targets). *Fix a reduced target $0 < t < 1/2$ with squarefree denominator and choose*

$$\gamma = e^{2\sqrt{t}}.$$

Then the direct family is admissible, $\alpha = 1/2$, and

$$\#\{A \in \mathcal{R}_X(t) : ||A| - \theta M| \leq w\} \geq 2^{(1-o(1))M}.$$

Moreover some exact cardinality $k_X = \theta M + O(w)$ supports at least $2^{(1-o(1))M}$ exact representations.

Proof. Here $(\log \gamma)^2 = 4t$, so $\lambda_\gamma = 2t \in (t, 1)$ and $\gamma > 1$. Hence $\alpha = t/\lambda_\gamma = 1/2$ and $H(\alpha) = \log 2$. Apply (7.10) and (7.15). $\square$

The corollary optimizes the entropy rate per available denominator within this construction. It is not an absolute optimization of the number of representations across different values of γ , because the ambient family size M itself depends strongly on γ .

The conclusion is deliberately mesoscopic. We do not prove a prescribed-cardinality asymptotic, a conditional central limit theorem for $|A|$ on the natural $\sqrt{M}$ scale, or a bivariate local limit for reciprocal sum and cardinality.

8. MOVING-TARGET LOCAL SATURATION

Keep the fixed reduced target $t = a/b \in (0, 1)$, exponent γ , direct family $\mathcal{E} = \mathcal{E}_X$, common denominator $L = L_X$, and Bernoulli parameter $\theta = t/\Lambda$. For a lattice target $s \in L^{-1}\mathbb{Z}$, write

$$W_X(s) = \Pr\left(\sum_{e \in \mathcal{E}} \frac{\xi_e}{e} = s\right). \quad (8.1)$$

We prove the full statement of Theorem 1.2.

8.1. The target shift is a unit character. Let

$$t_j = t + \frac{j}{L}, \quad j \in \mathbb{Z}. \quad (8.2)$$

The quotient coefficient at t_j is

$$LW_X^{\text{quot}}(t_j) = \sum_{h \bmod L} F_j(h), \quad (8.3)$$

where

$$F_j(h) = e(-ht_j) \prod_{e \in \mathcal{E}} ((1 - \theta) + \theta e(h/e)).$$

Comparing with the central target $t_0 = t$ gives the exact identity

$$F_j(h) = e(-hj/L) F_0(h). \quad (8.4)$$

Hence

$$|F_j(h)| = |F_0(h)| \quad (8.5)$$

for every h and j . All anchor, row, decoder, compression, and decoded-label tail estimates in Sections 4–6 are therefore unchanged.

8.2. The Gaussian phase. Fix $U > 0$ and consider integers satisfying

$$|j| \leq UL\sigma. \quad (8.6)$$

Put

$$u_j = \frac{j}{L\sigma}, \quad |u_j| \leq U. \quad (8.7)$$

On the frequency $h = m \pmod{L}$, equations (6.10) and (8.4) give, uniformly for $|m| \leq C/\sigma$,

$$F_j(m) = \exp\left(-2\pi^2 m^2 \sigma^2 - \frac{2\pi i m j}{L} + \varepsilon_{X,m}\right). \quad (8.8)$$

With $x_m = m\sigma$, the new phase is

$$\exp(-2\pi i x_m u_j). \quad (8.9)$$

Proposition 8.1 (Compact-uniform major limit). *For fixed $C, U > 0$, uniformly for integers j satisfying (8.6),*

$$\sigma \sum_{|m| \leq C/\sigma} F_j(m) - \int_{-C}^C e^{-2\pi^2 x^2} e(-u_j x) dx \longrightarrow 0. \quad (8.10)$$

Proof. The Taylor remainder is the same uniform error as in (6.10). For $|u| \leq U$, the functions

$$x \mapsto e^{-2\pi^2 x^2} e(-ux)$$

have uniformly bounded derivative on $[-C, C]$. The Riemann-sum approximation is therefore uniform in $u \in [-U, U]$. $\square$

The full Fourier transform is

$$\int_{\mathbb{R}} e^{-2\pi^2 x^2} e(-ux) dx = \Phi(u) := \frac{1}{\sqrt{2\pi}} e^{-u^2/2}. \quad (8.11)$$

The truncation tail outside $[-C, C]$ is bounded by the absolute Gaussian tail, uniformly in u .

8.3. Uniform complementary estimates. By (8.5), every modulus estimate in Sections 6.2–6.5 is unchanged. If $\mathfrak{R}_{X,C}(j)$ is the contribution remaining after removing the frequencies $h = m \pmod{L}$ with $|m| \leq C/\sigma$, then

$$\limsup_{X \rightarrow \infty} \sup_{|j| \leq UL\sigma} \sigma |\mathfrak{R}_{X,C}(j)| \leq K e^{-cC^2/2}. \quad (8.12)$$

No factor counting the number of targets appears; the estimate is already uniform in the inserted unit character.

Combining Proposition 8.1 with (8.12), first taking $X \rightarrow \infty$ with C, U fixed and then $C \rightarrow \infty$, gives

$$L\sigma W_X^{\text{quot}}(t_j) = \Phi(u_j) + o_U(1) \quad (8.13)$$

uniformly for (8.6).

8.4. No-wrap and simultaneous exact realization. By (2.9) and (2.10), the fixed centre t lies a positive distance from both endpoints of the eventual interval $[0, \Lambda]$. Since $\sigma \rightarrow 0$, for each fixed U all targets t_j with (8.6) lie in $[0, \Lambda]$ for sufficiently large X . Lemma 3.2 therefore gives

$$W_X^{\text{quot}}(t_j) = W_X(t_j). \quad (8.14)$$

Together with (8.13), this proves

$$L\sigma W_X\left(t + \frac{j}{L}\right) = \Phi\left(\frac{j}{L\sigma}\right) + o_U(1) \quad (8.15)$$

uniformly for $|j| \leq UL\sigma$, which is Theorem 1.2.

Since Φ has a positive minimum on $[-U, U]$, every coefficient in this window is eventually positive simultaneously. Hence the integer subset-sum image

$$S_X = \left\{ \sum_{e \in A} \frac{L}{e} : A \subseteq \mathcal{E} \right\} \quad (8.16)$$

contains the entire block

$$\{Lt + j : |j| \leq \lfloor UL\sigma \rfloor\}. \quad (8.17)$$

By (2.19) and Lemma 2.2,

$$\sigma \rightarrow 0, \quad L\sigma \rightarrow \infty. \quad (8.18)$$

Thus the number of represented lattice points tends to infinity while their radius after division by L is at most $U\sigma \rightarrow 0$.

Dividing (8.15) by the central local limit also gives

$$\frac{W_X(t + j/L)}{W_X(t)} = \exp\left(-\frac{j^2}{2L^2\sigma^2}\right) + o_U(1) \quad (8.19)$$

uniformly on the same window.

Remark 8.2. There is no largest finite admissible U : every fixed U works after a threshold depending on U . The proof does not give a growing standardized radius $U = U_X \rightarrow \infty$; such a statement would require quantitative errors small enough to compete with $e^{-U_X^2/2}$.

9. CHARACTERIZATION AND FINITE-AVOIDANCE CONSEQUENCES

The analytic work is complete. We now pass from a direct proper target to arbitrary positive rationals and record the qualitative consequences of finite avoidance.

9.1. The direct proper-target theorem.

Theorem 9.1 (Avoiding representation of a proper target). *Let $t = a/b \in (0, 1)$ be reduced with squarefree b , let $T \subset \mathbb{N}$ be finite, and fix $\gamma > 1$ with*

$$t < \frac{(\log \gamma)^2}{2} < 1.$$

Then, for all sufficiently large X , there is a finite set $A \subseteq \mathcal{E}_X$ of pairwise distinct squarefree semiprimes, disjoint from T , such that

$$\sum_{e \in A} \frac{1}{e} = t. \quad (9.1)$$

Proof. By Lemma 2.1, for sufficiently large X the direct family consists of pairwise distinct squarefree semiprimes outside T . The central local limit (6.28) gives $P = W_X(t) > 0$ for all sufficiently large X , so some subset represents t exactly. $\square$

Every proper rational target is covered by some exponent in this regime. Indeed, for $0 < t < 1$ the interval

$$e^{\sqrt{2t}} < \gamma < e^{\sqrt{2}} \quad (9.2)$$

is nonempty.

9.2. The squarefree-denominator characterization. We can now prove Theorem 1.1.

Proof of Theorem 1.1. For necessity, suppose

$$q = \sum_{e \in A} \frac{1}{e}$$

with every e squarefree, and put $D = \text{lcm}(A)$. Then

$$q = \frac{\sum_{e \in A} D/e}{D}.$$

The reduced denominator of q divides the squarefree integer D , and is therefore squarefree.

Conversely, let

$$q = \frac{a}{b} > 0$$

be reduced with squarefree b , and let T be a prescribed finite forbidden set. Choose a prime $p > a$ with $p \nmid b$ and set

$$t = \frac{a}{bp}. \quad (9.3)$$

Then $0 < t < 1$. Since $p > a$, one has $p \nmid a$; together with $(a, b) = 1$ and $p \nmid b$, this shows that (9.3) is reduced and has squarefree denominator bp .

Choose once and for all a γ satisfying (9.2) for this t . Construct supports $A_1, \dots, A_p$ successively. At stage j , apply Theorem 9.1 to t , forbidding

$$T \cup A_1 \cup \dots \cup A_{j-1}.$$

The supports are pairwise disjoint, each avoids T , and each has reciprocal sum t . Their union therefore represents

$$pt = \frac{a}{b} = q.$$

This proves sufficiency and arbitrary finite avoidance for every squarefree reduced denominator. $\square$

9.3. Thresholds, disjoint supports, and many terms.

Corollary 9.2. *Let $q > 0$ have squarefree reduced denominator.*

- (i) *For every N , q has a representation all of whose denominators exceed N .*
- (ii) *q has infinitely many representations with pairwise disjoint supports.*
- (iii) *q has representations with arbitrarily many terms.*

Proof. For (i), forbid $\{1, \dots, N\}$. For (ii), construct representations recursively, at each stage forbidding every denominator used earlier. For (iii), use (i): if every denominator exceeds N , then every unit fraction is less than $1/N$, so any representation of the fixed positive number q has more than qN terms. $\square$

9.4. Finite prescription.

Proposition 9.3 (Extension of a partial support). *Let $q > 0$ have squarefree reduced denominator. Let S be a finite set of distinct squarefree semiprimes such that*

$$\sum_{e \in S} \frac{1}{e} \leq q, \tag{9.4}$$

and let T be a finite set disjoint from S . Then S extends to a representation of q by distinct squarefree semiprimes which avoids T .

Proof. If equality holds in (9.4), there is nothing to add. Otherwise set

$$r = q - \sum_{e \in S} \frac{1}{e} > 0.$$

Let b be the reduced denominator of q and put

$$D := \text{lcm}(\{b\} \cup S).$$

Since b and every $e \in S$ are squarefree, D is squarefree. Writing the terms over the common denominator D gives $r = N/D$ for some positive integer N , and the reduced denominator of r divides D . Theorem 1.1 therefore represents r using denominators outside $S \cup T$. Adjoining that support to S gives the result. $\square$

Remark 9.4 (Common proper refinements). Finite avoidance also gives common proper refinements. Here a refinement replaces each original term by a disjoint block with the same reciprocal sum; it is proper if none of the original denominators is reused. For two representations of the same q , take the squarefree least common multiple D of all their denominators. The resulting positive integer weights have equal total sum, so the elementary greedy algorithm gives a nonnegative integral transportation matrix with them as row and column sums.

Each positive cell value x_{ij}/D has reduced denominator dividing the squarefree integer D . Represent it successively while forbidding all denominators from the two original representations, any externally prescribed finite set, and every denominator introduced in earlier cells. The resulting pairwise disjoint cell supports form a proper refinement of both representations; iteration handles any finite family.

These structural consequences are not used elsewhere; they show that finite avoidance is substantially stronger than bare existence.

10. FURTHER QUESTIONS AND PERSPECTIVE

The characterization in Theorem 1.1 applies to every positive rational with squarefree reduced denominator. The quantitative theorems concern one target-adapted family for a fixed reduced proper target $t = a/b \in (0, 1)$ and fixed $\gamma > 1$ satisfying

$$t < \frac{(\log \gamma)^2}{2} < 1.$$

Every proper target admits such a γ . The complete prime-pair block supplies the leading variance, while each fixed coordinate dividing b is observed by only $K_* = \lceil (\log Z)^2 \rceil$ target sensors.

10.1. Quantitative boundaries. The direct local-limit theorem concerns proper targets. Arbitrary positive rationals, including integers, are reached by prime dilution in Section 9; no analogous local-limit claim is made for one family centred at $t = 1$.

The assumption $\gamma > 1$ gives the scale separations used by the sparse target decoder on the Gaussian range and by the adaptive retained interval below the top block. The constraint $t < (\log \gamma)^2/2 < 1$ keeps the Bernoulli parameter in $(0, 1)$ and the reciprocal load inside one no-wrap interval.

The moving-target theorem is uniform for every fixed $|j|/(L_X \sigma_X) \leq U$, with an X -threshold depending on U . It gives neither a growing radius $U_X \rightarrow \infty$ nor a gap-free theorem on a fixed target-independent primordial family. Theorem 1.3 gives the ambient entropy exponent in a mesoscopic cardinality window and at one exact cardinality; Corollary 7.1 attains the binary rate $\log 2$ for $0 < t < 1/2$. No prescribed-cardinality asymptotic, conditional central limit theorem, bivariate local limit, or matching global upper bound is asserted.

There is nevertheless a first-order indication of asymptotic decorrelation. If $K_X = \sum_{e \in \mathcal{E}_X} \xi_e$, then

$$\text{Cov}(S_X, K_X) = \theta_X(1 - \theta_X)\Lambda_X \asymp 1, \quad \text{Corr}(S_X, K_X) \asymp X^{1-\gamma}(\log X)^2 \longrightarrow 0,$$

using $\sigma_X \asymp (X \log X)^{-1}$ and $\sqrt{\text{Var } K_X} \asymp Z/\log Z$. This motivates, but does not prove, a joint local law.

10.2. Finite avoidance is not deletion resilience. If $b > 1$, fix a prime $r \mid b$ and delete its sparse target row

$$\mathcal{D}_{r,X} = \{rq : q \in \mathcal{B}_*\}. \quad (10.1)$$

Although $|\mathcal{D}_{r,X}| = K_* = O((\log Z)^2) = o(M_X)$, every remaining weight L_X/e is divisible by r , whereas $L_X t = aL_X/b$ is nonzero modulo r . Thus finite avoidance

does not imply resilience of one constructed family under even this vanishing-cardinality deletion. The sensors are negligible in mass, variance, and cardinality but indispensable for observability.

10.3. A reusable finite-Fourier pattern. The proof isolates four reusable ingredients: (i) a weighted anchor partition turning low internal energy into coherent labels; (ii) row observability on the ranges where exact identification is needed; (iii) weighted row compression without a raw fibre-cardinality loss; and (iv) retained factors suppressing decoded labels after decoding. Together with a positive Gaussian major term and no-wrap, these form a finite-Fourier pattern in which observability need not track probabilistic mass.

Natural next questions concern structured positive-density prime sets and finer statistics of the exact fibre: growing target windows, prescribed-cardinality local asymptotics, and joint reciprocal-sum/cardinality laws.

10.4. Final perspective. The squarefree denominator is both the elementary obstruction and the exact one. Proper targets are solved directly and arbitrary positive rationals then follow by prime dilution. On the same direct family the exact coefficients have a local Gaussian profile, the typical exact fibre has the ambient entropy rate, and a growing consecutive block of nearby lattice targets is realized. The complete pair family carries the mass and variance while only polylogarithmically many target sensors restore the missing CRT coordinates.

ACKNOWLEDGEMENTS

Generative AI systems assisted mathematical exploration, proof development, source and citation checking, and editorial revision. The author takes full responsibility for the mathematics, citations, and text.

REFERENCES

- [1] T. F. Bloom. *Erdős Problem #306*. Erdős Problems.
URL: <https://www.erdosproblems.com/306> (visited on 2026-08-10).
- [2] T. F. Bloom and C. Elsholtz. ‘Egyptian Fractions’. In: *Nieuw Archief voor Wiskunde* 23.4 (2022), pp. 237–245. arXiv: 2210.04496 [math.NT].
- [3] S. Butler, P. Erdős and R. Graham. ‘Egyptian Fractions with Each Denominator Having Three Distinct Prime Divisors’. In: *Integers* 15 (2015), A51. DOI: 10.5281/zenodo.10456687.
- [4] A. Czenky, E. McGovern, J. Plavnik, E. Rowell and A. Watkins. *Egyptian fractions for few primes*. 2025. arXiv: 2507.03727 [math.NT].
- [5] P. Erdős and R. L. Graham. *Old and New Problems and Results in Combinatorial Number Theory*. Vol. 28. Monographies de L’Enseignement Mathématique. Geneva: L’Enseignement Mathématique, Université de Genève, 1980.
URL: https://fanchung.ucsd.edu/ron/papers/80_11_number_theory.pdf (visited on 2026-08-10).
- [6] W. Hoeffding. ‘Probability Inequalities for Sums of Bounded Random Variables’. In: *Journal of the American Statistical Association* 58.301 (1963), pp. 13–30. DOI: 10.1080/01621459.1963.10500830.
- [7] S. Li. *Every natural number is a sum of distinct semiprime unit fractions*. 2026. arXiv: 2606.15159 [math.NT].
- [8] Y. Tang. *A machine-checked proof of Erdős Problem 306 in Lean 4*. Version 0.0.3. 2026. DOI: 10.5281/zenodo.20767390.
- [9] T. Watanabe. *New examples of the representation of 1 by the sum of reciprocals of semiprime numbers*. 2020. arXiv: 2009.03275 [math.NT].