

Building and evaluation of a PBPK model for Warfarin in healthy adults

Version	main-OSP12.1
based on <i>Model Snapshot</i> and <i>Evaluation Plan</i>	https://github.com/Open-Systems-Pharmacology/Warfarin-Model/releases/tag/vmain
OSP Version	12.1
Qualification Framework Version	3.4

This evaluation report and the corresponding PK-Sim project file are filed at:

<https://github.com/Open-Systems-Pharmacology/OSP-PBPK-Model-Library/>

Table of Contents

- [1 Introduction](#)
- [2 Methods](#)
 - [2.1 Modeling Strategy](#)
 - [2.2 Data](#)
 - [2.2.1 In vitro / physico-chemical Data](#)
 - [2.2.2 Clinical Data](#)
 - [2.2.2.1 Model Building S-Warfarin](#)
 - [2.2.2.2 Model Verification S- Warfarin](#)
 - [2.2.2.3 Model Verification R- Warfarin](#)
 - [2.3 Model Parameters and Assumptions](#)
 - [2.3.1 Absorption](#)
 - [2.3.2 Distribution](#)
 - [2.3.3 Metabolism and Elimination](#)
 - [2.3.4 Automated Parameter Identification](#)
- [3 Results and Discussion](#)
 - [3.1 Final input parameters](#)
 - [3.2 Diagnostics Plots S-warfarin](#)
 - [3.3 Diagnostics Plots R-warfarin](#)
 - [3.4 Concentration-Time Profiles](#)
- [4 Conclusion](#)
- [5 References](#)

1 Introduction

The presented model building and evaluation report evaluates the performance of a PBPK model for Warfarin in healthy adults.

Warfarin is a widely used oral anticoagulant that exerts its therapeutic effect by inhibiting the hepatic synthesis of vitamin K–dependent clotting factors. Warfarin therapy requires careful monitoring because of its narrow therapeutic index and the risk of bleeding [Uyungül 2014](#).

Warfarin is classified as a BCS Class I drug, which means it has high solubility and high permeability. This classification indicates that warfarin is expected to dissolve rapidly and be well absorbed in the gastrointestinal tract [Kim 2019](#).

Warfarin is administered as a racemic mixture of two enantiomers, S-warfarin and R-warfarin, which differ in both potency and metabolism. The S-enantiomer is approximately two- to five-fold more potent as an anticoagulant compared to the R-enantiomer, but it is cleared more rapidly [Maddison 2013](#), [Geng 2024](#). The two enantiomers are metabolized by different cytochrome P450 isoforms. S-warfarin is primarily cleared by CYP2C9 with additional contribution from CYP3A4 and reduction to alcohols, while R-warfarin is metabolized mainly by CYP1A2, CYP2C19, and CYP3A4, as well as reduction to alcohol. Renal elimination of the unchanged parent drug is small for both enantiomers [Wittkowsky 2003](#).

The herein presented model building and evaluation report evaluates the performance of the PBPK model for S-Warfarin and R-Warfarin in healthy adults. Both enantiomers are modeled separately. These models integrate physicochemical properties, binding characteristics, distribution, and the known stereospecific clearance pathways of each enantiomer. The performance of the models is evaluated against published clinical data for different dosing regimens under fasted and fed conditions.

The presented Warfarin PBPK model as well as the respective evaluation plan and evaluation report are provided open-source (<https://github.com/Open-Systems-Pharmacology/Warfarin-model>).

2 Methods

2.1 Modeling Strategy

The general concept of building a PBPK model has previously been described by Kuepfer et al. ([Kuepfer 2016](#)) The relevant anthropometric (height, weight) and physiological parameters (e.g. blood flows, organ volumes, binding protein concentrations, hematocrit, cardiac output) in adults was gathered from the literature and has been previously published ([PK-Sim Ontogeny Database Version 7.3](#)). The information was incorporated into PK-Sim® and was used as default values for the simulations in adults.

The applied activity and variability of plasma proteins and active processes that are integrated into PK-Sim® are described in the publicly available PK-Sim® Ontogeny Database Version 7.3 ([Schlender 2016](#)) or otherwise referenced for the specific process.

First, a base mean model was built using clinical Phase I data including selected single dose studies with intravenous and oral applications of Warfarin to find an appropriate structure to describe the pharmacokinetics in plasma. The mean PBPK model was developed using a typical European individual.

Unknown parameters were identified using the Parameter Identification module provided in PK-Sim®. Structural model selection was mainly guided by visual inspection of the resulting description of data and biological plausibility.

Once the appropriate structural model was identified, additional parameters for tablet formulations were identified.

The model was then verified by simulating:

- Oral administration of Warfarin in fasted state
- Oral administration of Warfarin in fed state
- Intravenous administration of Warfarin

Details about input data (physicochemical, *in vitro* and clinical) can be found in [Section 2.2](#).

Details about the structural model and its parameters can be found in [Section 2.3](#).

2.2 Data

2.2.1 In vitro / physico-chemical Data

A literature search was performed to collect available information on physicochemical properties of Warfarin . The obtained information from literature is summarized in the table below.

Physicochemical and binding properties

Parameter	Unit	Value	Source	Description
MW	g/mol	308.33	Drugbank	Molecular weight
pK _a	–	5.03 (acid)	Drugbank	Acid dissociation constant
Solubility	mg/L	17 (at pH 7)	Drugbank	Aqueous solubility
f _u _{plasma}		0.009	Takahashi 2003	Fraction unbound in plasma (albumin binding)
LogP	–	3.46	Avdeef 1998	Liposomal membrane-water partition coefficient
B/P	–	0.55	—	Blood-to-plasma ratio (used in reduction pathway)

S-warfarin - Metabolic and clearance parameters

Pathway / Enzyme	Parameter	Unit	Value	Source / Note
CYP2C9	Km	μM	4.4	Experimental Km within range 3.5 to 5.54 μM Shaik 2016 , Rettie 1992 , Rettie 1992 , Yong 2009 , Crewe 2011
CYP3A4	Km	μM	14.9	Experimental Km within range 14 to 27 Shaik 2016 , Jones 2010
Reduction to alcohols	Plasma CL	ml/min/kg	0.14	Metabolism reduction mediated by aldo-keto reductases (i.e., fraction metabolized ~ 10%) Wittkowsky 2003
	Specific CL	1/min	0.02	
Renal clearance	Plasma CL	l/h/kg	0.07	Consistent with fraction excreted in urine ~ 1% FDA 2010
	Specific CL	1/min	0.02	

R-warfarin - Metabolic and clearance parameters

Pathway / Enzyme	Parameter	Unit	Value	Source / Note
CYP1A2	Km	μM	648	Adopted from previous PK-Sim model
CYP2C19	Km	μM	391	Adopted from previous PK-Sim model
CYP3A4	Km	μM	586	Adopted from previous PK-Sim model
Reduction to alcohols	Plasma CL Specific CL	ml/h/kg 1/min	0.27 0.02	
Renal clearance	Plasma CL Specific CL	ml/h/kg 1/min	0.07 0.02	

2.2.2 Clinical Data

A literature search was performed to collect available clinical data on Warfarin in healthy adults.

2.2.2.1 Model Building S-Warfarin

The following studies were used for model building (training data):

Dose [mg]	Dosing	PK data	Dataset	Reference
15	po, sd	plasma	training	Soon 2006
25	po, sd	plasma	training	Schwartz 2008

iv = intravenous; po = oral administration; tab = tablet administration; sd = single dose; qd = once daily

2.2.2.2 Model Verification S- Warfarin

The following studies were used for model verification:

Dose [mg]	Dosing	PK data	Dataset	Reference
0.375	iv, tab, fasted, sd	plasma	verification	O'Sullivan 1993
7.5	po, tab, fasted with meal administration 3h post-dose, sd	plasma	verification	Frymoyer 2010
10	po, tab, fasted, sd	plasma	verification	Kim 2001
10	po, tab, fasted, sd	plasma	verification	Lilja 2007
10	po, tab, fasted, sd	plasma	verification	Ngo 2010
10	po, tab, fasted, sd	plasma	verification	Turpeinen 2013
13	po, tab, fed, sd	plasma	verification	Weber 1999
15	po, tab, fasted, sd	plasma	verification	Schwartz 2009
20	po, tab, fasted, sd	plasma	verification	Ouellet 2006
25	po, tab, fasted, sd	plasma	verification	Almeida 2008
25	po, tab, fasted, sd	plasma	verification	Dingemanse 2013
25	po, tab, fasted, sd	plasma	verification	He 2007
25	po, tab, fasted, sd	plasma	verification	Macha 2013
25	po, tab, fasted, sd	plasma	verification	Malhotra 2011
25	po, tab, fasted, sd	plasma	verification	Rahimy 2002
25	po, tab, fasted, sd	plasma	verification	Sanwald Ducray 2014
25	po, tab, fasted, sd	plasma	verification	Stockis 2013
25	po, tab, fasted, sd	plasma	verification	Toon 1987
25	po, tab, fasted, sd	plasma	verification	Toon 1990
25	po, tab, fed, sd	plasma	verification	Mallikaarjun 1999
25	po, tab, fed, sd	plasma	verification	Sidharta 2014
25	po, tab, fed, sd	plasma	verification	Soon 2006
25	po, tab, fed, sd	plasma	verification	Yin 2011
30	po, tab, fasted with meal administration 4h post-dose, sd	plasma	verification	Liao 1996

iv = intravenous; po = oral administration; tab = tablet administration; sd = single dose; qd = once daily

2.2.2.3 Model Verification R- Warfarin

The following studies were used for model verification:

Dose [mg]	Dosing	PK data	Dataset	Reference
-----------	--------	---------	---------	-----------

<<<<<<< HEAD | 25| po, tab, fasted, sd
 |plasma|verification|[Almeida 2008](#)| | 25| po, tab,
 fasted, sd |plasma|verification|[Dingemante 2013](#)| |
 25| po, tab, fasted, sd |plasma|verification|[He
 2007](#)| | 25| po, tab, fasted, sd
 |plasma|verification|[Malhotra 2011](#)| | 25| po, tab,
 fasted, sd |plasma|verification|[Rahimy 2002](#)| | 25|
 po, tab, fasted, sd |plasma|verification|[Stockis
 2013](#)| | 25| po, tab, fasted, sd
 |plasma|verification|[Toon 2087](#)| | 25| po, tab,
 fasted, sd |plasma|verification|[Toon 1990](#)| | 25| po,
 tab, fed, sd |plasma|verification|[Mallikaarjun 1999](#)|
 | 25| po, tab, fed, sd |plasma|verification|[Sidharta
 2014](#)| | 25| po, tab, fed, sd
 |plasma|verification|[Soon 2006](#)| | 25| po, tab, fed,
 sd |plasma|verification|[Yin 2011](#)|

| 25| po, tab, fasted, sd |plasma|verification|[Almeida 2008](#)| | 25| po, tab, fasted, sd |plasma|verification|[Dingemante 2013](#)| |
 25| po, tab, fasted, sd |plasma|verification|[He 2007](#)| | 25| po, tab, fasted, sd |plasma|verification|[Malhotra 2011](#)| | 25| po,
 tab, fasted, sd |plasma|verification|[Rahimy 2002](#)| | 25| po, tab, fasted, sd |plasma|verification|[Stockis 2013](#)| | 25| po, tab,
 fasted, sd |plasma|verification|[Toon 1987](#)| | 25| po, tab, fasted, sd |plasma|verification|[Toon 1990](#)| | 25| po, tab, fed, sd
 |plasma|verification|[Mallikaarjun 1999](#)| | 25| po, tab, fed, sd |plasma|verification|[Sidharta 2014](#)| | 25| po, tab, fed, sd
 |plasma|verification|[Soon 2006](#)| | 25| po, tab, fed, sd |plasma|verification|[Yin 2011](#)|

||| | c4bac1efd21d652db42ed9b81ab341fed3579aca

iv = intravenous; po = oral administration; tab = tablet administration; sd = single dose; qd = once daily

2.3 Model Parameters and Assumptions

2.3.1 Absorption

The measured solubility of Warfarin was taken from Product information from [Drugbank](#) (see [Section 2.2.1](#)).

Tablet dissolution was modeled using an empirical Weibull dissolution approach. Both enantiomer share the same absorption processes.

2.3.2 Distribution

After testing the available organ-plasma partition coefficient and cell permeability calculation methods built in PK-Sim®, observed clinical data was best described by choosing the partition coefficient calculation by `Rodgers and Rowland` and cellular permeability calculation by `Charged dependent Schmitt`.

2.3.3 Metabolism and Elimination

Clearance of warfarin was described through enzymatic and reductive pathways, with only a negligible contribution from renal excretion of unchanged parent drug. The stereospecific nature of warfarin metabolism was explicitly represented by modeling the two enantiomers separately, with different sets of enzymes responsible for their elimination.

For **S-warfarin**, metabolism is dominated by CYP2C9, which is the principal enzyme responsible for oxidative clearance. A smaller contribution from CYP3A4 was also included, together accounting for the observed in-vivo clearance. In addition, a reductive pathway leading to alcohol metabolites was implemented, using literature-based plasma clearance values and a blood-to-plasma ratio of 0.55 [Wittkowsky 2003](#), leading to a hepatic `Specific Clearance` of 0.02 1/min. The renal elimination of unchanged S-warfarin was parameterized but remains negligible, consistent with clinical evidence of ~ 1% fraction excreted in urine [FDA 2010](#). Kinetic constants for CYP2C9 and CYP3A4 were taken from in-vitro studies and the associated `Kcat` were optimized within the current model to achieve agreement with observed plasma concentrations. These optimized values reflect the dominant role of CYP2C9 and the minor but non-negligible contribution of CYP3A4 to the overall clearance of the S-enantiomer.

The clearance of **R-warfarin** was represented by three oxidative enzymes—CYP1A2, CYP2C19, and CYP3A4—together with the same reductive alcohol pathway and minor renal clearance. Experimental information on kinetic constants for R-warfarin is less abundant, and therefore the K_m and V_{max} values were adopted from a previously developed PBPK model in PK-Sim. The values for the Michaelis–Menten constants are relatively high (K_m values of 648 μM for CYP1A2, 391 μM for CYP2C19, and 586 μM for CYP3A4), which indicates a low affinity of these enzymes for R-warfarin. The reduction to alcohols and renal clearance contribute only marginally to the overall disposition of the R-enantiomer.

2.3.4 Automated Parameter Identification

This is the result of the final parameter identification.

Model Parameter	Optimized Value	Unit
<code>Intestinal permeability (transcellular)</code>	0.0015	cm/min
<code>S-Warfarin CYP2C9 kcat</code>	0.2	1/min
<code>S-warfarin CYP3A4 kcat</code>	0.09	1/min

3 Results and Discussion

The PBPK model for Warfarin, was developed and verified with clinical pharmacokinetic data.

The model was evaluated covering data from studies including in particular

- Intravenous Bolus
- Oral administration over fasted and fed states.

The next sections show:

1. the final model parameters for the building blocks: [Section 3.1](#).
2. the overall goodness of fit: [Section 3.2](#).
3. simulated vs. observed concentration-time profiles for the clinical studies used for model building and for model verification: [Section 3.3](#).

3.1 Final input parameters

The compound parameter values of the final PBPK model are illustrated below.

Compound: S-Warfarin

Parameters

Name	Value	Value Origin	Alternative	Default
Solubility at reference pH	17 mg/l	Internet-DrugBank (Tomlin 1997)	Measurement	True
Reference pH	7	Internet-DrugBank (Tomlin 1997)	Measurement	True
Lipophilicity	3.46 Log Units	Publication-logP_mem_neutral (Avdeef 1998)	Measurement	True
Fraction unbound (plasma, reference value)	0.009	Publication-Takahashi 2003	Measurement	True
Specific intestinal permeability (transcellular)	0.0015 cm/min		fitted	True
Is small molecule	Yes			
Molecular weight	308.33 g/mol			
Plasma protein binding partner	Albumin			

Calculation methods

Name	Value
Partition coefficients	Rodgers and Rowland
Cellular permeabilities	Charge dependent Schmitt

Processes

Systemic Process: Renal Clearances-Urine excretion data (Heimark, Toon, O'Reilly)

Species: Human

Parameters

Name	Value	Value Origin
Fraction unbound (experiment)	0.009	
Plasma clearance	0.07 ml/h/kg	
Specific clearance	0.02 1/min	Unknown

Metabolizing Enzyme: CYP2C9-HLM Mean

Molecule: CYP2C9

Parameters

Name	Value	Value Origin
In vitro Vmax for liver microsomes	6.9 pmol/min/mg mic. protein	
Content of CYP proteins in liver microsomes	96 pmol/mg mic. protein	Unknown
Km	4.4 µmol/l	
kcat	0.2022708099 1/min	Parameter Identification-Parameter Identification-Value updated from 'Parameter Identification' on 2025-08-01 11:48

Systemic Process: Total Hepatic Clearance-Reduction to alcohol

Species: Human

Parameters

Name	Value	Value Origin
Fraction unbound (experiment)	0.009	
Lipophilicity (experiment)	3.46 Log Units	
Blood/Plasma concentration ratio	0.55	Unknown
Plasma clearance	0.14 ml/min/kg	
Specific clearance	0.02 1/min	Unknown

Metabolizing Enzyme: CYP3A4-HLM mean

Molecule: CYP3A4

Parameters

Name	Value	Value Origin
Enzyme concentration	1 µmol/l	
Vmax	0 µmol/l/min	
Km	14.9 µmol/l	
kcat	0.0872168808 1/min	Parameter Identification-Parameter Identification-Value updated from 'Parameter Identification' on 2025-08-01 11:48

Compound: R-Warfarin

Parameters

Name	Value	Value Origin	Alternative	Default
Solubility at reference pH	17 mg/l	Internet-DrugBank (Tomlin 1997)	Measurement	True
Reference pH	7	Internet-DrugBank (Tomlin 1997)	Measurement	True
Lipophilicity	3.46 Log Units	Publication-logP_mem_neutral (Avdeef 1998)	Measurement	True
Fraction unbound (plasma, reference value)	0.009	Publication-Takahashi 2003	Measurement	True
Specific intestinal permeability (transcellular)	0.0015 cm/min		fitted	False
Is small molecule	Yes			
Molecular weight	308.33 g/mol			
Plasma protein binding partner	Albumin			

Calculation methods

Name	Value
Partition coefficients	Rodgers and Rowland
Cellular permeabilities	Charge dependent Schmitt

Processes

Metabolizing Enzyme: CYP1A2-HLM Mean

Molecule: CYP1A2

Parameters

Name	Value	Value Origin
In vitro Vmax for liver microsomes	775 pmol/min/mg mic. protein	
Content of CYP proteins in liver microsomes	45 pmol/mg mic. protein	Unknown
Km	648 µmol/l	

Metabolizing Enzyme: CYP2C19-HLM Mean

Molecule: CYP2C19

Parameters

Name	Value	Value Origin
In vitro Vmax for liver microsomes	178 pmol/min/mg mic. protein	
Content of CYP proteins in liver microsomes	19 pmol/mg mic. protein	Unknown
Km	391 µmol/l	

Metabolizing Enzyme: CYP3A4-HLM Mean

Molecule: CYP3A4

Parameters

Name	Value	Value Origin
In vitro Vmax for liver microsomes	886 pmol/min/mg mic. protein	
Km	586 µmol/l	

Systemic Process: Total Hepatic Clearance-Reduction to alcohols

Species: Human

Parameters

Name	Value	Value Origin
Fraction unbound (experiment)	0.009	
Lipophilicity (experiment)	3.46 Log Units	
Blood/Plasma concentration ratio	0.55	Unknown
Plasma clearance	0.27 ml/h/kg	

Systemic Process: Renal Clearances-Urine excretion data (Heimark, Toon, O'Reilly)

Species: Human

Parameters

Name	Value	Value Origin
Fraction unbound (experiment)	0.009	
Plasma clearance	0.07 ml/h/kg	

3.2 Diagnostics Plots S-warfarin

Below you find the goodness-of-fit visual diagnostic plots for the PBPK model performance of all data used presented in [Section 2.2.2](#).

The first plot shows observed versus simulated plasma concentration, the second weighted residuals versus time.

Table 3-1: GMFE for S-Warfarin Goodness of fit plot for concentration in plasma

Group	GMFE
S-Warfarin IV Administration (model verification)	1.14
S-Warfarin Oral Administration (model building)	1.23
S-Warfarin Oral Administration fasted (model verification)	1.40
S-Warfarin Oral Administration fed (model verification)	1.16
All	1.33

- ▲ S-Warfarin IV Administration (model verification)
- S-Warfarin Oral Administration (model building)
- S-Warfarin Oral Administration fasted (model verification)
- ◆ S-Warfarin Oral Administration fed (model verification)




Figure 3-1: S-Warfarin Goodness of fit plot for concentration in plasma




Figure 3-2: S-Warfarin Goodness of fit plot for concentration in plasma

3.3 Diagnostics Plots R-warfarin

Below you find the goodness-of-fit visual diagnostic plots for the PBPK model performance of all data used presented in [Section 2.2.2](#).

The first plot shows observed versus simulated plasma concentration, the second weighted residuals versus time.

Table 3-2: GMFE for R-Warfarin Goodness of fit plot for concentration in plasma, R-warfarin

Group	GMFE
R-Warfarin Oral Administration (model verification)	1.32




Figure 3-3: R-Warfarin Goodness of fit plot for concentration in plasma, R-warfarin




Figure 3-4: R-Warfarin Goodness of fit plot for concentration in plasma, R-warfarin

3.4 Concentration-Time Profiles

Simulated versus observed concentration-time profiles of all data listed in [Section 2.2.2](#) are presented below.




Figure 3-5: Warfarin Na 7.5mg (S-Warfarin 3.75mg) fasted + meal post-dose




Figure 3-6: Warfarin 10mg (S-Warfarin 5mg) PO fasted




Figure 3-7: Warfarin 26mg (S-Warfarin 13mg) PO fed




Figure 3-8: Warfarin Na 30mg (S-Warfarin 15mg) fasted




Figure 3-9: Warfarin Na 20mg (S-Warfarin 10mg) PO fasted




Figure 3-10: Warfarin 25mg (S-Warfarin 12.5mg) PO fasted




Figure 3-11: Warfarin Na 25mg (S-Warfarin 12.5mg) PO fed




Figure 3-12: Warfarin 7.5mg (S-Warfarin 15mg) PO fasted + meal post-dose




Figure 3-13: S-Warfarin 0.375mg/kg IV fasted




Figure 3-14: Warfarin Na 25mg (R-Warfarin 12.5mg) PO fasted

4 Conclusion

The herein presented PBPK model adequately describes the pharmacokinetics of Warfarin in adults after iv and oral administration. Since S-warfarin is the more potent enantiomer, its model development and verification were more extensive. If modeling of R-warfarin becomes necessary in the future, additional work will be required.

5 References

- Almeida 2008** Almeida L, Falcão A, Vaz-da-Silva M, Nunes T, Santos AT, Rocha JF, Neta C, Macedo T, Fontes-Ribeiro C, Soares-da-Silva P. Effect of nebicapone on the pharmacokinetics and pharmacodynamics of warfarin in healthy subjects. *Eur J Clin Pharmacol*. 2008 Oct;64(10):961-6. doi: 10.1007/s00228-008-0534-2. Epub 2008 Aug 6.
- Avdeef 1998** Avdeef A, Box KJ, Comer JE, Hibbert C, Tam KY. pH-metric logP 10. Determination of liposomal membrane-water partition coefficients of ionizable drugs. *Pharm Res*. 1998 Feb;15(2):209-15. doi: 10.1023/a:1011954332221.
- Crewe 2011** Crewe HK, Barter ZE, Yeo KR, Rostami-Hodjegan A. Are there differences in the catalytic activity per unit enzyme of recombinantly expressed and human liver microsomal cytochrome P450 2C9? A systematic investigation into inter-system extrapolation factors. *Biopharm Drug Dispos*. 2011 Sep;32(6):303-18. doi: 10.1002/bdd.760.
- Dingemans 2013** Dingemans J, Hoever P. Absence of pharmacokinetic and pharmacodynamic interactions between almorexant and warfarin in healthy subjects. *Drugs R D*. 2013 Jun;13(2):145-51. doi: 10.1007/s40268-013-0017-5.
- Drugbank** DrugBank DB00682 <https://go.drugbank.com/drugs/DB00682>, accessed 26-08-2025.
- FDA 2010** COUMADIN® TABLETS, FDA drug label. https://www.accessdata.fda.gov/drugsatfda_docs/label/2010/009218s108lbl.pdf
- Frymoyer 2010** Frymoyer A, Shugarts S, Browne M, Wu AH, Frassetto L, Benet LZ. Effect of single-dose rifampin on the pharmacokinetics of warfarin in healthy volunteers. *Clin Pharmacol Ther*. 2010 Oct;88(4):540-7. doi: 10.1038/clpt.2010.142. Epub 2010 Aug 11.
- Jones 2010** Jones DR, Kim SY, Boysen G, Yun CH, Miller GP. Contribution of three CYP3A isoforms to metabolism of R- and S-warfarin. *Drug Metab Lett*. 2010 Dec;4(4):213-9. doi: 10.2174/187231210792928242.
- Lilja 2007** Lilja JJ, Backman JT, Neuvonen PJ. Effects of daily ingestion of cranberry juice on the pharmacokinetics of warfarin, tizanidine, and midazolam--probes of CYP2C9, CYP1A2, and CYP3A4. *Clin Pharmacol Ther*. 2007 Jun;81(6):833-9. doi: 10.1038/sj.clpt.6100149. Epub 2007 Mar 28.
- Geng 2024** Geng K, Shen C, Wang X, Wang X, Shao W, Wang W, Chen T, Sun H, Xie H. A physiologically-based pharmacokinetic/pharmacodynamic modeling approach for drug-drug-gene interaction evaluation of S-warfarin with fluconazole. *CPT Pharmacometrics Syst Pharmacol*. 2024 May;13(5):853-869. doi: 10.1002/psp4.13123.
- He 2007** He YL, Sabo R, Riviere GJ, Sunkara G, Leon S, Ligueros-Saylan M, Rosenberg M, Dole WP, Howard D. Effect of the novel oral dipeptidyl peptidase IV inhibitor vildagliptin on the pharmacokinetics and pharmacodynamics of warfarin in healthy subjects. *Curr Med Res Opin*. 2007 May;23(5):1131-8. doi: 10.1185/030079907x188008.
- Kim 2001** Kim JS, Nafziger AN, Gaedigk A, Dickmann LJ, Rettie AE, Bertino JS Jr. Effects of oral vitamin K on S- and R-warfarin pharmacokinetics and pharmacodynamics: enhanced safety of warfarin as a CYP2C9 probe. *J Clin Pharmacol*. 2001 Jul;41(7):715-22. doi: 10.1177/00912700122010618.
- Kim 2019** Kim YJ, Jeong JW, Song Y, Koo TS. Effects of formulation types on pharmacodynamics of warfarin in patients with cerebral infarction and dysphagia. *Clin Pharmacol*. 2019 Mar 13;11:51-56. doi: 10.2147/CPAA.S184232.
- Liao 1996** Liao S, Palmer M, Fowler C, Nayak RK. Absence of an effect of levofloxacin on warfarin pharmacokinetics and anticoagulation in male volunteers. *J Clin Pharmacol*. 1996 Nov;36(11):1072-7. doi: 10.1177/009127009603601111.
- Macha 2013** Macha S, Rose P, Mattheus M, Pinnetti S, Woerle HJ. Lack of drug-drug interaction between empagliflozin, a sodium glucose cotransporter 2 inhibitor, and warfarin in healthy volunteers. *Diabetes Obes Metab*. 2013 Apr;15(4):316-23. doi: 10.1111/dom.12028. Epub 2012 Nov 22.
- Maddison 2013** Maddison J, Somogyi AA, Jensen BP, James HM, Gentgall M, Rolan PE. The pharmacokinetics and pharmacodynamics of single dose (R)- and (S)-warfarin administered separately and together: relationship to VKORC1

genotype. *Br J Clin Pharmacol.* 2013 Jan;75(1):208-16. doi: 10.1111/j.1365-2125.2012.04335.x.

Malhotra 2011 Malhotra B, Alvey C, Gong J, Li X, Duczynski G, Gandelman K. Effects of fesoterodine on the pharmacokinetics and pharmacodynamics of warfarin in healthy volunteers. *Br J Clin Pharmacol.* 2011 Aug;72(2):257-62. doi: 10.1111/j.1365-2125.2011.03989.x.

Mallikaarjun 1999 Mallikaarjun S, Bramer SL. Effect of cilostazol on the pharmacokinetics and pharmacodynamics of warfarin. *Clin Pharmacokinet.* 1999;37 Suppl 2:79-86. doi: 10.2165/00003088-199937002-00009.

Ngo 2010 Ngo N, Brantley SJ, Carrizosa DR, Kashuba AD, Dees EC, Kroll DJ, Oberlies NH, Paine MF. The warfarin-cranberry juice interaction revisited: A systematic in vitro-in vivo evaluation. *J Exp Pharmacol.* 2010 Jul;2010(2):83-91. doi: 10.2147/JEP.S11719.

O'Sullivan 1993 O'Sullivan TA, Wang JP, Unadkat JD, al-Habet SM, Trager WF, Smith AL, McNamara S, Aitken ML. Disposition of drugs in cystic fibrosis. V. In vivo CYP2C9 activity as probed by (S)-warfarin is not enhanced in cystic fibrosis. *Clin Pharmacol Ther.* 1993 Sep;54(3):323-8. doi: 10.1038/clpt.1993.154.

Ouellet 2006 Ouellet D, Bramson C, Carvajal-Gonzalez S, Roman D, Randinitis E, Remmers A, Gardner MJ. Effects of lasofoxifene on the pharmacokinetics and pharmacodynamics of single-dose warfarin. *Br J Clin Pharmacol.* 2006 Jun;61(6):741-5. doi: 10.1111/j.1365-2125.2006.02589.x.

Rahimy 2002 Rahimy M, Hallén B, Narang P. Effect of tolterodine on the anticoagulant actions and pharmacokinetics of single-dose warfarin in healthy volunteers. *Arzneimittelforschung.* 2002;52(12):890-5. doi: 10.1055/s-0031-1299986.

Rettie 1992 Rettie AE, Korzekwa KR, Kunze KL, Lawrence RF, Eddy AC, Aoyama T, Gelboin HV, Gonzalez FJ, Trager WF. Hydroxylation of warfarin by human cDNA-expressed cytochrome P-450: a role for P-4502C9 in the etiology of (S)-warfarin-drug interactions. *Chem Res Toxicol.* 1992 Jan-Feb;5(1):54-9. doi: 10.1021/tx00025a009.

Sanwald Ducray 2014 Sanwald-Ducray P, Jamois C, Banken L. The effect of aleglitazar on the pharmacokinetics and pharmacodynamics of S- and R-warfarin in healthy male subjects. *J Cardiovasc Pharmacol.* 2014 Feb;63(2):152-7. doi: 10.1097/FJC.0000000000000033.

Shaik 2016 Shaik AN, Grater R, Lulla M, Williams DA, Gan LL, Bohnert T, LeDuc BW. Comparison of enzyme kinetics of warfarin analyzed by LC-MS/MS QTrap and differential mobility spectrometry. *J Chromatogr B Analyt Technol Biomed Life Sci.* 2016 Jan 1;1008:164-173. doi: 10.1016/j.jchromb.2015.11.036. Epub 2015 Nov 23.

Schwartz 2008 Schwartz JI, Dunbar S, Yuan J, Li S, Gipson A, Rosko K, Johnson-Levonas AO, Lasseter KC, Addy C, Stoch AS, Wagner JA. Influence of taranabant, a cannabinoid-1 receptor inverse agonist, on pharmacokinetics and pharmacodynamics of warfarin. *Adv Ther.* 2008 Nov;25(11):1175-90. doi: 10.1007/s12325-008-0116-9.

Sidharta 2014 Sidharta PN, Dietrich H, Dingemanse J. Investigation of the effect of macitentan on the pharmacokinetics and pharmacodynamics of warfarin in healthy male subjects. *Clin Drug Investig.* 2014 Aug;34(8):545-52. doi: 10.1007/s40261-014-0207-0.

Soon 2006 Soon D, Kothare PA, Linnebjerg H, Park S, Yuen E, Mace KF, Wise SD. Effect of exenatide on the pharmacokinetics and pharmacodynamics of warfarin in healthy Asian men. *J Clin Pharmacol.* 2006 Oct;46(10):1179-87. doi: 10.1177/0091270006291622.

Stockis 2013 Stockis A, van Lier JJ, Cawello W, Kumke T, Eckhardt K. Lack of effect of lacosamide on the pharmacokinetic and pharmacodynamic profiles of warfarin. *Epilepsia.* 2013 Jul;54(7):1161-6. doi: 10.1111/epi.12192.

Takahashi 2003 Takahashi H, Wilkinson GR, Caraco Y, Muszkat M, Kim RB, Kashima T, Kimura S, Echizen H. Population differences in S-warfarin metabolism between CYP2C9 genotype-matched Caucasian and Japanese patients. *Clin Pharmacol Ther.* 2003 Mar;73(3):253-63. doi: 10.1067/mcp.2003.26a.

Toon 1987 Toon S, Hopkins KJ, Garstang FM, Rowland M. Comparative effects of ranitidine and cimetidine on the pharmacokinetics and pharmacodynamics of warfarin in man. *Eur J Clin Pharmacol.* 1987;32(2):165-72. doi:

Toon 1990 Toon S, Holt BL, Mullins FG, Bullingham R, Aarons L, Rowland M. Investigations into the potential effects of multiple dose ketorolac on the pharmacokinetics and pharmacodynamics of racemic warfarin. *Br J Clin Pharmacol.* 1990 Nov;30(5):743-50. doi: 10.1111/j.1365-2125.1990.tb03845.x.

Turpeinen 2013 Turpeinen M, Uusitalo J, Lehtinen T, Kailajärvi M, Pelkonen O, Vuorinen J, Tapanainen P, Stjernschantz C, Lammintausta R, Scheinin M. Effects of ospemifene on drug metabolism mediated by cytochrome P450 enzymes in humans in vitro and in vivo. *Int J Mol Sci.* 2013 Jul 5;14(7):14064-75. doi: 10.3390/ijms140714064.

Uygungül 2014 Uygungül E, Ayrik C, Narci H, Erdoğan S, Toker I, Demir F, Karaaslan U. Determining risk factors of bleeding in patients on warfarin treatment. *Adv Hematol.* 2014;2014:369084. doi: 10.1155/2014/369084.

Weber 1999 Weber C, Banken L, Birnboeck H, Schulz R. Effect of the endothelin-receptor antagonist bosentan on the pharmacokinetics and pharmacodynamics of warfarin. *J Clin Pharmacol.* 1999 Aug;39(8):847-54. doi: 10.1177/00912709922008380.

Wittkowsky 2003 Wittkowsky AK. Warfarin and other coumarin derivatives: pharmacokinetics, pharmacodynamics, and drug interactions. *Semin Vasc Med.* 2003 Aug;3(3):221-30. doi: 10.1055/s-2003-44457.

Yin 2011 Yin OQ, Gallagher N, Fischer D, Zhao L, Zhou W, Leroy E, Golor G, Schran H. Effects of nilotinib on single-dose warfarin pharmacokinetics and pharmacodynamics: a randomized, single-blind, two-period crossover study in healthy subjects. *Clin Drug Investig.* 2011;31(3):169-79. doi: 10.2165/11538700-000000000-00000.

Yong 2009 Yong WP, Kim TW, Undevia SD, Innocenti F, Ratain MJ. R(+)-XK469 inhibits hydroxylation of S-warfarin by CYP2C9. *Eur J Cancer.* 2009 Jul;45(11):1904-8. doi: 0.1016/j.ejca.2009.04.032.