

A second year mathematics degree

Yu Coughlin

Contents

L1	1	Euclidean spaces	2
	1.1	Euclidean norm	2
	1.2	Convergence in \mathbb{R}^n	2
	2	Continuity and limits of functions	3
	2.1	Open sets	3
	2.2	Continuity	3
	3	Derivative of maps of Euclidean spaces	3
	3.1	Total derivatives	3
	3.2	Directional and partial derivatives	3
	3.3	Higher order derivatives	4
	4	Inverse and implicit function theorems	5
	4.1	Inverse function theorem	5
	4.2	Implicit function theorem	5
	5	Metric spaces	5
	5.1	Introduction	5
	5.2	Normed vector spaces	6
	5.3	Open and closed sets	6
	5.4	Separable space	7
	6	Continuous maps in metric spaces	7
	6.1	Convergence	7
	6.2	Continuity of maps	8
	6.3	Metric homeomorphisms	8
	7	Topological spaces	8
	7.1	Topologies and their spaces	8
	7.2	Bases	9
	7.3	Closed sets	9
	7.4	Convergence and Hausdorff property	9
	7.5	Continuous maps	10
	7.6	Subspaces	10
	8	Connectedness	10
	8.1	Definition	10
	8.2	Continuous maps	11
	8.3	Path connected sets	11
	9	Compactness	11
	9.1	Covers	11
	9.2	Sequential compactness	11
	9.3	Continuous maps	11
	9.4	Arzelá-Ascoli theorem	11
	10	Completeness	11
	10.1	Banach spaces	11
	10.2	Fixed point theorem	11
	1	Groups and Rings	12
	1	Quotient groups	13
	1.1	Group homomorphisms	13
	1.2	Normal subgroups	13
	1.3	Quotient groups	13
	1.4	Isomorphism theorems	13

	1.5	Centres	15
	1.6	Commutators	15
	1.7	Torsion and p -primary subgroups	15
	1.8	Generators	15
	1.9	Classification of finitely generated Abelian groups	16
2	Group actions		16
	2.1	Actions	16
	2.2	Orbit-stabiliser theorem	16
	2.3	Jordan's theorem	17
3	Rings		17
	3.1	Rings	17
	3.2	Ring homomorphisms	17
	3.3	Ideals	18
4	Integral domains		18
	4.1	Integral domains	18
	4.2	Charateristic	18
	4.3	Polynomial rings	19
5	PIDs and UFDs		19
	5.1	Euclidian domains	19
	5.2	Principal ideal domains	19
	5.3	Unique factorisation domains	19
6	Fields		19
	6.1	Vector spaces	19
	6.2	Field extensions	20
	6.3	Constructing fields	20
	6.4	Existence of finite fields	20
2	Lebesgue Measure and Integration		21
L1	1	Motivation	24
	2	Measures	24
	2.1	Algebras and σ -algebras	24
	2.2	Measures	24
	2.3	Complete measure spaces	24
	3	Constructing measures	24
	3.1	Pre-measure	24
	3.2	Outer measure	24
	3.3	Restriction	24
	3.4	Lebesgue measure	24
	4	Measurable functions	24
	4.1	Defintion	24
	4.2	Properties	24
	4.3	Continuity	24
	5	Lebesgue integral	24
	5.1	Construction	24
	5.2	Properties	24
	6	Convergence	24
	6.1	Monotone convergence	24
	6.2	Fatou's lemma	24
	6.3	Lebesgue dominated convergence	24
	6.4	Vitali's theorem	24
	7	L^p spaces	24
	7.1	Norms	24
	7.2	L^p spaces	24
	7.3	Normed vector spaces	24
	7.4	Completeness	24
	8	Product measures	24
	8.1	Products of sets	24
	8.2	σ -algebras on product sets	24
	8.3	Product measures	24

9	Fubini's theorem	24
9.1	Motivations	24
9.2	Setup	24
9.3	Fubini's theorem	24
10	Differentiation	24
10.1	Hardy-Littlewood maximal function	24
10.2	Compact support spaces	24
10.3	Lebesgue's differentiation theorem	24
11	Decomposition	24
11.1	Signed measures	24
11.2	Hahn decomposition theorem	24
11.3	Mutually singular measures	24
11.4	Jordan decomposition theorem	24
11.5	Lebesgue decomposition theorem	24
11.6	Radon-Nikodym theorem	24
3	Categories	25
1	Basic definitions	26
1.1	Categories	26
1.2	Functors	26
1.3	Natural transformations	27
1.4	Equivalence of categories	27
1.5	Representable functors	27
1.6	Yoneda lemma	27

Real Analysis and Topology

Lectured by Someone
Typed by Yu Coughlin
Season Year

Introduction

The following are complementary reading for the course.

- G. Grimmett and D. J. A. Welsh, Probability: An Introduction, 1986
- J. K. Blitzstein and J. Hwang, Introduction to Probability, 2019
- D. F. Anderson et al, Introduction to Probability, 2018
- S. M. Ross, Introduction to Probability Models, 2014
- G. Grimmett and D. Stirzaker, Probability and Random Processes, 2001
- G. Grimmett and D. Stirzaker, One Thousand Exercises in Probability, 2009

Notation. Unbracketed superscripts are used to label the components of vectors, with unbracketed subscripts labelling different vectors.

1 Euclidean spaces

Definition 1.0.1 (\mathbb{R}^n). The set $\mathbb{R}^n = \{(x^1, x^2, \dots, x^n) : x^i \in \mathbb{R}, \forall i \in [1, n]\}$ will be considered with the operations to make it a real vector space.

1.1 Euclidean norm

Definition 1.1.1 (Inner product). We will have the **inner product** on \mathbb{R}^n by $\langle \cdot, \cdot \rangle : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ satisfying:

$$\langle x, y \rangle := \sum_{i=1}^n x^i y^i,$$

with the **Euclidean norm** given by,

$$\|\cdot\| : \mathbb{R}^n \rightarrow [0, \infty) \text{ with } \|x\| = \sqrt{\langle x, x \rangle}.$$

Proposition 1.1.2 (Properties of the Euclidean norm). The Euclidean norm satisfies the following properties:

(N1) for all $x \in \mathbb{R}^n$, $\|x\| \geq 0$ achieving equality iff $x = 0$,

(N2) for all $x \in \mathbb{R}^n$ and $\lambda \in \mathbb{R}$, $\|\lambda x\| = |\lambda| \cdot \|x\|$,

(N3) for all $x, y \in \mathbb{R}^n$: $\|x + y\| \leq \|x\| + \|y\|$,

Theorem 1.1.3 (Cauchy-Swartz inequality). For all $x, y \in \mathbb{R}^n$, $|\langle x, y \rangle| \leq \|x\| \cdot \|y\|$.

Theorem 1.1.4 (Reverse triangle inequality). For all $x, y \in \mathbb{R}^n$, $|\|x\| - \|y\|| \leq \|x - y\|$.

Proposition 1.1.5. For $x = (x^1, x^2, \dots, x^n) \in \mathbb{R}^n$,

$$\max_{k \in [1, n]} |x^k| \leq \|x\| \leq \sqrt{n} \max_{k \in [1, n]} |x^k|.$$

Proof. Exercise □

1.2 Convergence in \mathbb{R}^n

Definition 1.2.1 (Open ball). In \mathbb{R}^n we define the **open ball** around $x \in \mathbb{R}^n$ of size $r \in \mathbb{R}$ as

$$B_r(x) := \{y \in \mathbb{R}^n : \|x - y\| < r\}.$$

This will be analogous to the notion of open intervals used throughout analysis 1.

Definition 1.2.2 (Sequence in \mathbb{R}^n). A **sequence** in \mathbb{R}^n is an ordered list $x_0, x_1, \dots, x_i \dots$ with $x_i \in \mathbb{R}^n$ for all $i \in \mathbb{N}$, written $(x_i)_{i=0}^\infty$

Definition 1.2.3 (Convergence in \mathbb{R}^n). We say a sequence in \mathbb{R}^n , $(x_i)_{i=0}^\infty$ **converges to** $x \in \mathbb{R}^n$ iff

$$\forall \epsilon > 0 \exists N \in \mathbb{N} \text{ such that, } \forall n \geq N, \|x_i - x\| < \epsilon$$

and we write $x_i \rightarrow x$ as $i \rightarrow \infty$ or $\lim_{i \rightarrow \infty} x_i = x$.

Lemma 1.2.4. The sequence of vectors in \mathbb{R}^n , $(x_i)_{i=0}^\infty$, converges to some $x = (x^1, x^2, \dots, x^n) \in \mathbb{R}^n$ iff each component of x_i converges to the corresponding component in x :

$$\forall k \in [1, n] \lim_{i \rightarrow \infty} x_i^k = x^k.$$

Proof. (\implies) Given $\lim_{i \rightarrow \infty} x_i^k = x^k$ for all $k \in [1, n]$ we have that for all $\epsilon > 0$, $|x_i^k - x^k| < \frac{\epsilon}{\sqrt{n}}$ for all $i \geq N_k$ for each $k \in [1, n]$ respectively. We take $N = \max_{k \in [1, n]} N_k$ and now have:

$$\max_{k \in [1, n]} |x_i^k - x^k| < \frac{\epsilon}{\sqrt{n}} \implies \|x_i - x\| \leq \sqrt{n} \max_{k \in [1, n]} |x_i^k - x^k| < \epsilon.$$

(\impliedby) Similarly, given $\lim_{i \rightarrow \infty} x_i = x \implies \|x_i - x\| < \epsilon$ for all $\epsilon > 0$:

$$|x_i^k - x^k| \leq \max_{k \in [1, n]} |x_i^k - x^k| \leq \|x_i - x\| < \epsilon,$$

therefore $\lim_{i \rightarrow \infty} x_i^k = x^k$ for all $k \in [1, n]$. □

2 Continuity and limits of functions

2.1 Open sets

Definition 2.1.1 (Open set in \mathbb{R}^n). A subset $U \subseteq \mathbb{R}^n$ is **open** in \mathbb{R}^n iff:

$$\forall x \in U, \exists r > 0 \text{ such that } B_r(x) \subseteq U.$$

2.2 Continuity

Definition 2.2.1 (Continuity). Let $A \subseteq \mathbb{R}^n$ then we have $f : A \rightarrow \mathbb{R}^m$ **continuous at** some $p \in A$ iff

$$\forall \epsilon > 0, \exists \delta > 0 \text{ such that } \forall x \in A \text{ with } \|x - p\| < \delta, \|f(x) - f(p)\| < \epsilon.$$

If f is continuous at all $p \in A$ we say f is **continuous on** A .

Theorem 2.2.2. Let $A \subseteq \mathbb{R}^n$ and $B \subseteq \mathbb{R}^m$ with $f : A \rightarrow B$ continuous at $p \in A$. Suppose $g : B \rightarrow \mathbb{R}^l$ is continuous at $f(p)$, then $g \circ f : A \rightarrow \mathbb{R}^l$ is continuous at p .

Proof. Given any $\epsilon > 0$ have $\|x - p\| < \delta_f \circ \delta_g(\epsilon) \implies \|f(x) - f(p)\| < \delta_g(\epsilon) \implies \|g \circ f(x) - g \circ f(p)\| < \epsilon$. \square

3 Derivative of maps of Euclidean spaces

3.1 Total derivatives

Definition 3.1.1 (Total derivative). Given open $\Omega \subset \mathbb{R}^n$, the function $f : \Omega \rightarrow \mathbb{R}^m$ is **differentiable at** $p \in \Omega$ iff there is a linear map $\Lambda : \mathbb{R}^n \rightarrow \mathbb{R}^m$ satisfying:

$$\lim_{x \rightarrow p} \frac{\|f(x) - f(p) - \Lambda(x - p)\|}{\|x - p\|} = 0.$$

Have $Df(p) := \Lambda$ be the **total derivative** of f at p .

Remark 3.1.2. Given $f : (a, b) \rightarrow \mathbb{R}$ differentiable at $p \in (a, b)$, we have

$$\begin{aligned} \lim_{x \rightarrow p} \frac{\|f(x) - f(p) - \Lambda(x - p)\|}{\|x - p\|} &= \lim_{x \rightarrow p} \frac{|f(x) - f(p) - \lambda \cdot (x - p)|}{|x - p|} = \lim_{x \rightarrow p} \left| \frac{f(x) - f(p)}{x - p} - \lambda \right| = 0 \\ \implies \lim_{x \rightarrow p} \left| \frac{f(x) - f(p)}{x - p} \right| &= \lambda, \text{ which satisfies the normal definition for a derivative.} \end{aligned}$$

Theorem 3.1.3 (Uniqueness of total derivative). If the total derivative of a function $f : \Omega \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$ exists, then it is unique.

Proof. \square

Theorem 3.1.4 (Chain rule). Let $\Omega \subset \mathbb{R}^n$, $\Omega' \subset \mathbb{R}^m$ be open and have $g : \Omega \rightarrow \Omega'$, $f : \Omega' \rightarrow \mathbb{R}^l$ differentiable at $p, g(p)$ respectively and let $h := f \circ g$, $Dh(p) = Df(g(p)) \circ Dg(p)$.

Proof. \square

3.2 Directional and partial derivatives

Definition 3.2.1 (Directional derivative). Suppose $\Omega \subseteq \mathbb{R}^n$ is open with $f : \Omega \rightarrow \mathbb{R}^m$ differentiable at $p \in \Omega$. For all $v \in \mathbb{R}^n$ the **directional derivative** of f at p in the direction of v is:

$$\frac{\partial f}{\partial v}(p) := \lim_{t \rightarrow 0} \frac{f(p + tv) - f(p)}{t} = Df(p)[v].$$

With the partial derivatives of f given by:

$$D_i f(p) := \frac{\partial f}{\partial e_i}(p), \text{ for all } i \in [1, n].$$

Remark 3.2.2. If the total derivative of a function exists, then so do all of its directional derivatives.

Theorem 3.2.3. If $\Omega \subset \mathbb{R}^n$ is open with $f : \Omega \rightarrow \mathbb{R}$ with all partial derivatives existing for all $x \in \Omega$. If the map $x \mapsto D_i f(x)$ is continuous at $p \in \Omega$ for all partial derivatives, then f is differentiable at p .

Proof. \square

3.3 Higher order derivatives

Definition 3.3.1 (Second order partial derivatives). Let $\Omega \subset \mathbb{R}^n$ be open with differentiable $f : \Omega \rightarrow \mathbb{R}$ written as $(f^1, f^2, \dots, f^n)^T$, the ik th second partial derivative at p is

$$D_k D_i f^j(p) := \lim_{t \rightarrow 0} \frac{D_i f^j(p + te_k) - D_i f^j(p)}{t}.$$

This can naturally be extended to n th order partial derivatives.

Theorem 3.3.2. Given open $\Omega \subseteq \mathbb{R}^n$ and $f : \Omega \rightarrow \mathbb{R}^m$ differentiable on Ω , consider the map:

$$\begin{aligned} Df &: \Omega \longrightarrow \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m) \cong M_{n \times m}(\mathbb{R}) \cong \mathbb{R}^{m \times n} \\ p &\longmapsto Df(p) \end{aligned},$$

which we can now show to be continuous or differentiable at $p \in \Omega$, when differentiable we can take $DDf(p) \in \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$. The components of the corresponding matrix are give by:

$$[DDf(p)[h]]_{ij} = \sum_{k=1}^n D_k D_i f^j(p) h^k.$$

Proof. □

Remark 3.3.3. The condition of a function being k times differentiable at a point p can is often difficult to establish, instead the continuous existence of all $k - th$ partial derivatives in a neighbourhood of p is a preferable question which implies the former statement.

Theorem 3.3.4 (Schwartz's theorem). Suppose $\Omega \subseteq \mathbb{R}^n$ is open and $f : \Omega \rightarrow \mathbb{R}^m$ is differentiable on Ω with $D_i D_j f(p), D_j D_i f(p)$ both exist continuous only Ω ; then we have

$$D_i D_j f(p) = D_j D_i f(p) \text{ for all } p \in \Omega.$$

Proof. □

Notation 3.3.5. We need the following necessary notation around an n -vector of non-negative integers, $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n) \in (\mathbb{Z}_{>0})^n$ for some $n \in \mathbb{Z}_{>0}$, to easily express Taylor's theorem in multiple dimensions:

1. $|\alpha| = \alpha_1 + \alpha_2 + \dots + \alpha_n$,
2. $D^\alpha f = (D_1)^{\alpha_1} (D_2)^{\alpha_2} \dots (D_n)^{\alpha_n}$,
3. for some vector $h = (h^1, h^2, \dots, h^n) \in \mathbb{R}^n$, $h^\alpha = ((h^1)^{\alpha_1}, (h^2)^{\alpha_2}, \dots, (h^n)^{\alpha_n})$,
4. $\alpha! = \alpha_1! \alpha_2! \dots \alpha_n!$.

Theorem 3.3.6 (Taylor's theorem). Given $p \in \mathbb{R}^n$ with $f : B_r(p) \rightarrow \mathbb{R}$, for some $r > 0$, k -times continuous differentiable on $B_r(p)$ and some $\|h\| < r$; we have:

$$f(p + h) = \sum_{|\alpha| \leq k-1} \frac{h^\alpha}{\alpha!} D^\alpha f(p) + R_k(p, h).$$

Where the remainder term, $R_k(p, h)$ is given by:

$$R_k(p, h) = \sum_{|\alpha|=k} \frac{h^\alpha}{\alpha!} D^\alpha f(x).$$

Proof. □

4 Inverse and implicit function theorems

4.1 Inverse function theorem

Theorem 4.1.1 (Inverse function theorem). Have $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ continuous differentiable on $\Omega \subseteq \mathbb{R}^n$ and $Df(p)$ be invertible for a $p \in \Omega$. There exists open sets $U \in \Omega$ and $V \in \mathbb{R}^n$ such that $f : U \rightarrow V$ is a bijection. Furthermore, $f^{-1} : V \rightarrow U$ is continuous differentiable on V with:

$$Df^{-1}(y) = [Df(f^{-1}(y))]^{-1}.$$

Lemma 4.1.2. Have $B_r(p) \subset \mathbb{R}^n$ with $f : B_r(p) \rightarrow \mathbb{R}^n$ continuously differentiable. If there exists some $M \in \mathbb{R}_{>0}$ with $|D_j f^i(x)| < M$ for all $x \in B_r(p)$ then

$$\|f(x) - f(y)\| \leq nM\|x - y\|, \text{ for all } x, y \in B_r(p).$$

Proof.

□

Lemma 4.1.3.

Lemma 4.1.4.

Lemma 4.1.5.

Proof of Theorem 4.1.1 (Inverse function theorem).

□

4.2 Implicit function theorem

Theorem 4.2.1 (Implicit function theorem). Given $\Omega \subseteq \mathbb{R}^n$ and $\Omega' \subseteq \mathbb{R}^m$ both open with $f : \Omega \times \Omega' \rightarrow \mathbb{R}^m$ continuous differentiable on $\Omega \times \Omega'$. If there is some $p \in \Omega \times \Omega'$ with $f(p) = 0$ and $D_{n+j} f^i(p)$ invertible for $1 \leq i, j \leq m$. Then, there are open sets $A \in \Omega$ and $B \in \Omega'$ containing a and b respectively such that for all $x \in A$ there is a unique and differentiable $g(x) \in B$ with $f(x, g(x)) = 0$.

Proof.

□

5 Metric spaces

5.1 Introduction

Definition 5.1.1 (Metric). A **metric** on some arbitrary set X is a function:

$$d : X \times X \rightarrow \mathbb{R}$$

that satisfies the following properties for all $x, y, z \in X$:

(M1) $d(x, y) \geq 0$ with $d(x, y) = 0$ iff $x = y$ (positivity),

(M2) $d(x, y) = d(y, x)$ (symmetry),

(M3) $d(x, y) \leq d(x, z) + d(z, y)$ (triangle inequality).

Definition 5.1.2 (Metric space). A **metric space** is a pair consisting of a set and a metric on said set, often denoted $M = (X, d)$. The elements of X are called **points** and for any two points of M , x, y , their **distance (with respect to d)** is $d(x, y)$.

Examples 5.1.3. The following are common examples of metric spaces:

1. have $X = \mathbb{R}$ and $d_1 : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ by $d_1(x, y) := |x - y|$,

2. have $X = \mathbb{R}^n$ and have $d(x, y) := \sqrt{\sum_{i=1}^n (x^i - y^i)^2}$,

3. for an arbitrary non-empty set X we have $d_{\text{disc}} : X \times X \rightarrow \mathbb{R}$ by $d_{\text{disc}}(x, y) := 0$ iff $x = y$ and 1 otherwise (discrete metric),

4. have X be the set of bounded real sequences, then we can have $d_\infty : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ given by $d_\infty(x, y) := \sup_{k \geq 1} |x^k - y^k|$,
5. let X be the set of continuous real functions on $[a, b]$ with $d(f, g) := \int_{t=a}^b |f(t) - g(t)| dt$.

Definition 5.1.4 (Induced metric). Given the metric space (X, d) and some $Y \subset X$, we have $d_Y : Y \times Y \rightarrow \mathbb{R}$ with $d_Y(x, y) = d(x, y)$ for all $x, y \in Y$ as the **induced metric** on Y . (Y, d_Y) is a **metric subspace** of (X, d) .

5.2 Normed vector spaces

Definition 5.2.1 (Normed vector spaces). Given a real-vector space V , a function $\|\cdot\| : V \rightarrow \mathbb{R}$ is a **norm** on V iff the following hold for all $u, v \in V$:

- (N1) $\|v\| \geq 0$ with $\|v\| = 0$ iff $v = 0_V$,
- (N2) for all $\lambda \in \mathbb{R}$, $\|\lambda v\| = |\lambda| \cdot \|v\|$,
- (N3) $\|u + v\| \leq \|u\| + \|v\|$.

A vector space together with a norm is a **normed vector space**.

Lemma 5.2.2. If $(V, \|\cdot\|)$ is a normed vector space, $d_{\|\cdot\|} : V \times V \rightarrow \mathbb{R}$ with $d_{\|\cdot\|}(u, v) = \|u - v\|$ is a metric on V .

Proof.

□

5.3 Open and closed sets

Definition 5.3.1 (ϵ -ball). Given a point x in the metric space (X, d) and a real $\epsilon > 0$, the **ball** of radius ϵ centred at x is the set,

$$B_\epsilon(x) := \{y \in X : d(x, y) < \epsilon\},$$

which is sometimes referred to as a neighbourhood of x .

Definition 5.3.2 (Open sets). Given metric space (X, d) a set $U \subseteq X$ is **open** in (X, d) iff, for all $u \in U$ there exists some $\delta > 0$ such that $B_\delta(u) \subseteq U$.

Proposition 5.3.3. Have $\mathcal{X} = (X, d)$ a metric space, the follow hold true:

1. \emptyset and \mathcal{X} are open in \mathcal{X} ,
2. for all $x \in \mathcal{X}$ and $\epsilon > 0$, $B_\epsilon(x)$ is open in \mathcal{X} ,
3. the union of (up to uncountably many) open sets in \mathcal{X} are open in \mathcal{X} ,
4. the intersection of finitely many open sets in \mathcal{X} is open in \mathcal{X} .

Proof.

□

Definition 5.3.4 (Topological equivalence). Two metrics d, d' on X are **topologically equivalent** iff $U \subseteq X$ is open in (X, d) iff it is also open in (X, d') .

Definition 5.3.5 (Closed sets). Given the metric space (X, d) with $U \subseteq X$, U is **closed** iff $X \setminus U$ is open.

Proposition 5.3.6. A set $U \subseteq X$ with (X, d) a metric space is closed iff, every convergent sequence in V has a limit in V .

Proof.

□

Proposition 5.3.7. The intersection of (up to countable many) closed sets in a metric space is closed; the union of finitely many sets in a metric space is closed.

Proof.

□

5.4 Separable space

Definition 5.4.1 (Interior, isolated, limits and boundary points). We will have (X, d) be a metric space with $V \subseteq X$ and $x \in X$:

- x is an **interior point** of V if there is some $\delta > 0$ with $B_\delta(x) \subseteq V$,
- x is an **isolated point** of V if there is some $\delta > 0$ such that $V \cap B_\delta(x) = \{x\}$,
- x is a **limit point** of V if for all $\delta > 0$, we have $(B_\delta(x) \cap V) \setminus \{x\} \neq \emptyset$,
- x is a **boundary point** of V if it is a limit point, under the previous definition, and $B_\delta(x) \setminus V \neq \emptyset$.

Remark 5.4.2. Interior and isolated points are necessarily in V , but limit points and boundary points need not be elements of V .

Definition 5.4.3 (Interior, closure and boundary). Once again, we will have (X, d) a metric space with $V \subseteq X$:

- the **interior** of V is the set of all $v \in V$ with v an interior point of V , denoted V° ,
- the **closure** of V is the union of V with the set of limit points of V , denoted \bar{V} ,
- the **boundary** of V is the set of boundary points of V , denoted ∂V .

Proposition 5.4.4. $\partial V = \bar{V} \setminus V^\circ$.

Proof.

□

Definition 5.4.5 (Dense set). Have (X, d) a metric space, $V \subseteq X$ is **dense** in (X, d) iff $\bar{V} = X$.

Definition 5.4.6 (Separable space). We say the metric space (X, d) is **separable** if there is a countable, dense set in X .

6 Continuous maps in metric spaces

6.1 Convergence

Definition 6.1.1 (Convergence in metric spaces). Let $(x_n)_{n \geq 1}$ be a sequence in the metric space (X, d) . We say $(x_n)_{n \geq 1}$ **converges** in (X, d) iff:

$$\exists x \in X \text{ such that, } \forall \epsilon > 0, \exists N \in \mathbb{Z}_{>0} \text{ with } d(x_n, x) < \epsilon \text{ for all } n \geq N.$$

And we say $(x_n)_{n \geq 1}$ converges to x in (X, d) , or any other equivalent phrasing from analysis.

Definition 6.1.2 (Cauchy sequences). A sequence $(x_n)_{n \geq 1}$ is **Cauchy** in (X, d) iff

$$\forall \epsilon > 0, \exists N \in \mathbb{Z}_{>0} \text{ such that } \forall n, m \geq N, d(x_n, x_m) < \epsilon.$$

Lemma 6.1.3 (Uniqueness of limits). If the sequence $(x_n)_{n \geq 1}$ converges to some x in the metric space (X, d) then this limit is unique.

Proof.

□

Theorem 6.1.4. Given two topologically equivalent metrics d, d' on X , the sequence $(x_n)_{n \geq 1}$ converges in (X, d) iff it also converges in (X, d') .

Proof.

□

6.2 Continuity of maps

Definition 6.2.1 (Continuous map). Given the metric spaces $(X, d_X), (Y, d_Y)$ and $f : X \rightarrow Y$:

1. f is **continuous at** $x \in X$ iff for all $x' \in X$:

$$\forall \epsilon > 0, \exists \delta > 0 \text{ such that } d_X(x, x') < \delta \implies d_Y(f(x), f(x')) < \epsilon,$$

2. f is **continuous on** $U \subseteq X$ if f is continuous at every $u \in U$,
3. f is **uniformly continuous** on $U \subseteq X$ if f is continuous on U and $\delta = \delta(\epsilon)$ does not depend on x .

Theorem 6.2.2. Let $(X, d_X), (Y, d_Y)$ be metric spaces, a function $f : X \rightarrow Y$ is continuous iff the pre-image of any open $U \subseteq Y$ is open in X .

Proof. □

Proposition 6.2.3. If, similarly, $(X, d_X), (Y, d_Y)$ are metric spaces with $f : X \rightarrow Y$, the following are equivalent:

1. f is continuous at $x \in X$,
2. if a sequence $(x_n)_{n \geq 1}$ converges to $x \in X$ then $(f(x_n))_{n \geq 1}$ converges to $f(x) \in Y$.

Proof. □

6.3 Metric homeomorphisms

Definition 6.3.1 (Homeomorphism). Have $(X, d_X), (Y, d_Y)$ be metric spaces, a mapping $f : X \rightarrow Y$ is a **homeomorphism** if it is a bijection with f, f^{-1} both continuous. Metric spaces with homeomorphisms between them are **homeomorphic**.

Definition 6.3.2 (Lipschitz). Given metric spaces $(X, d_X), (Y, d_Y)$ and $f : X \rightarrow Y$ we say:

1. f is **Lipschitz** if there is some $M > 0$ with:

$$d_Y(f(x_1), f(x_2)) \leq M \cdot d_X(x_1, x_2) \text{ for all } x_1, x_2 \in X,$$

2. f is **bi-Lipschitz** if there is some $M_1, M_2 > 0$ with:

$$M_1 \cdot d_X(x_1, x_2) \leq d_Y(f(x_1), f(x_2)) \leq M_2 \cdot d_X(x_1, x_2) \text{ for all } x_1, x_2 \in X,$$

3. f is **isometric** if,

$$d_Y(f(x_1), f(x_2)) = d_X(x_1, x_2) \text{ for all } x_1, x_2 \in X.$$

Remark 6.3.3. An isometry between metric spaces is a bi-Lipschitz map with two unit constants.

7 Topological spaces

7.1 Topologies and their spaces

Definition 7.1.1 (Topology). Given a non-empty set X , we say τ , a collection of subsets of X , is a **topology** on X if it satisfies the following conditions:

(T1) $\emptyset, X \subseteq \tau$,

(T2) if $X_i \in \tau$ for all i in a indexing set \mathcal{I} , $\bigcup_{i \in \mathcal{I}} X_i \in \tau$,

(T3) if $X_1, X_2, \dots, X_n \in \tau$, $\bigcap_{i=1}^n X_i \in \tau$.

The pair (X, τ) is called a **topological space** with elements of X called **points** and elements of τ called open sets. If $x \in X$ and $x \in U \in \tau$, U is a neighbourhood of x .

Examples 7.1.2. These are some common examples of topological spaces:

1. for any set X have $\tau = \{\emptyset, X\}$, the trivial topology on X ,

2. instead have τ be the collection of subsets of X , the discrete topology on X ,
3. if (X, d) is a metric space, $\tau := \{U \subseteq X : U \text{ is open in } (X, d)\}$ the metric topology on X ,
4. for a non-empty set X , $\tau = \{\emptyset, V, X\}$ for some non-empty $V \subset X$,
5. if $X = \{a, b\}$ and $\tau = \{\emptyset, \{a, b\}, \{b\}\}$ is the smallest topological space that is neither trivial nor discrete (called the Sierpinski topology).

Definition 7.1.3 (Metrisability). A topological space (X, τ) is **metrisable** iff it is the topology induced by some metric.

Definition 7.1.4 (Coarser and finer topologies). Given two topologies τ_1, τ_2 both on X , we say τ_1 is **coarser** than τ_2 , and equivalently τ_2 is **finer** than τ_1 , iff $\tau_2 \subseteq \tau_1$.

7.2 Bases

Definition 7.2.1 (Basis). Given a topological space (X, τ) we call a subfamily $B \subseteq \tau$ a **basis** for τ iff every open set in τ is the union of open sets in B .

7.3 Closed sets

Definition 7.3.1 (Closed sets). Given a topological space (X, τ) , we say $V \subseteq X$ is **closed** iff $X \setminus V$ is open.

Proposition 7.3.2. Closed sets in any given topological space (X, τ) satisfy the following:

- (C1) X, \emptyset are closed,
- (C2) if C_1, C_2 are closed, $C_1 \cup C_2$ is closed,
- (C3) the (up to uncountable) intersection of closed sets is closed.

Proof. □

Definition 7.3.3 (Closure). Given an open set U in the topological space (X, τ) the **closure** of U in (X, τ) is given by:

$$\bar{U} := \bigcap_{\substack{V \subseteq X \\ V \text{ closed}, A \subseteq V}} V.$$

Definition 7.3.4 (Point of closure). Given the topological space \mathcal{X} with $A \subseteq \mathcal{X}$, $x \in \mathcal{X}$ is a **point of closure** of A iff every open set U with $x \in U$ has $U \cap A \neq \emptyset$.

Proposition 7.3.5. $\bar{A} = \{x \in X : x \text{ is a point of closure for } A\}$.

Proof. □

7.4 Convergence and Hausdorff property

Definition 7.4.1 (Convergence). For a sequences $(x_n)_{n \geq 1}$ in a topological space (X, τ) we say $(x_n)_{n \geq 1}$ **converges** (in (X, τ)) to $x \in X$ iff

$$\forall T \in \tau \text{ with } x \in T, \exists N \in \mathbb{Z}_{>0} \text{ such that } \forall n \geq N, x_n \in T.$$

Definition 7.4.2 (Hausdorff). A topological space (X, τ) is **Hausdorff** iff for all $x, y \in X$ with $x \neq y$ there are open sets U, V containing x, y respectively with $U \cap V = \emptyset$. With U and V **separating** x and y .

Theorem 7.4.3. Limits of convergent sequences in Hausdorff spaces are unique.

Proof. □

Definition 7.4.4 (Regular spaces). A topological space (X, τ) is **regular** iff for every closed subset $C \subseteq X$ with point $p \notin C$ there are open sets $U, V \in \tau$ such that $p \in U$, $C \subseteq V$ and $U \cap V = \emptyset$.

7.5 Continuous maps

Definition 7.5.1 (Continuous map). Given two topological spaces $(X, \tau_X), (Y, \tau_Y)$ the map $f : X \rightarrow Y$ is **continuous** iff $f^{-1}(U) \in \tau_X$ for all $U \in \tau_Y$.

Definition 7.5.2 (Continuity at points). The map $f : X \rightarrow Y$, with $(X, \tau_X), (Y, \tau_Y)$ topological spaces, is **continuous at** $x \in X$ iff $f^{-1}(U) \in \tau_X$ for all $U \in \tau_Y$ with $f(x) \in U$.

Definition 7.5.3 (Homeomorphism). A **homeomorphism** between topological spaces is a bijection map, f , where both f and f^{-1} are continuous. Spaces with homeomorphisms between them are **topologically equivalent**.

7.6 Subspaces

Definition 7.6.1 (Subspace). If (X, τ) is a topological space and $A \subseteq X$, the **subspace topology** on A is $\tau_A = \{A \cap U : U \in \tau\}$, (A, τ_A) is a topological space called the **subspace** of (X, τ) .

Proof of topological space. □

Proposition 7.6.2 (Universal property). Given topological spaces $(X, \tau_X), (Y, \tau_Y)$ with $A \subseteq X$ with its subspace topology and $g : Y \rightarrow A$, g is continuous iff $i \circ g$ is continuous, where i is the inclusion map,

$$\begin{array}{ccc} & X & \\ i \circ g \nearrow & \uparrow i & \\ Y & \xrightarrow{g} & A \end{array}.$$

Proof. □

Theorem 7.6.3. Given the topological space (X, τ) and $A \subseteq X$, the subspace topology is the only topology such that for all (Y, τ_Y) , $g : Y \rightarrow A$ is continuous iff $(i \circ g)$ is continuous.

Proof. □

Lemma 7.6.4. If B is a basis for the topological space (X, τ) and $A \subseteq X$, $B_A := \{U \cap A : U \in B\}$ is a basis for τ_A .

Proof. □

Proposition 7.6.5. For a metric space (X, d) with $A \subseteq X$, the two canonical topologies on A , τ_{d_A} and T_A are equal.

Proof. □

8 Connectedness

8.1 Definition

Definition 8.1.1 (Disconnected sets). Let

8.2 Continuous maps**8.3 Path connected sets****9 Compactness****9.1 Covers****9.2 Sequential compactness****9.3 Continuous maps****9.4 Arzelá-Ascoli theorem****10 Completeness****10.1 Banach spaces****10.2 Fixed point theorem**

Chapter 1

Groups and Rings

Lectured by Someone
Typed by Yu Coughlin
Autumn 2024

Introduction

The following are complementary reading for the course.

- G. Grimmett and D. J. A. Welsh, Probability: An Introduction, 1986
- J. K. Blitzstein and J. Hwang, Introduction to Probability, 2019
- D. F. Anderson et al, Introduction to Probability, 2018
- S. M. Ross, Introduction to Probability Models, 2014
- G. Grimmett and D. Stirzaker, Probability and Random Processes, 2001
- G. Grimmett and D. Stirzaker, One Thousand Exercises in Probability, 2009

1 Quotient groups

1.1 Group homomorphisms

Definition 1.1.1 (Group isomorphism). Given groups G, H , a function $f : G \rightarrow H$ is a **group isomorphism** if it is a bijective group homomorphism. If there exists an isomorphism between groups, G is **isomorphic** to H written $G \cong H$.

Definition 1.1.2 (Group automorphism). Given G a group, an isomorphism $f : G \xrightarrow{\sim} G$ is a **group automorphism**.

Theorem 1.1.3. $\text{Aut } G$ (the set of automorphisms of a group G) is a group under function composition.

Proof. By examining the definition of $\text{Aut } G$, taking $e = \text{id}$ and showing association elementwise. \square

Theorem 1.1.4. Given groups G, H , if $f : G \xrightarrow{\sim} H$ then $f^{-1} : H \xrightarrow{\sim} G$.

Proof. $f^{-1}(f(g_1))f^{-1}(f(g_2)) = g_1g_2 = f^{-1}(f(g_1g_2)) = f^{-1}(f(g_1)g(g_2))$ is sufficient as f is surjective. \square

1.2 Normal subgroups

Definition 1.2.1 (Normal subgroup). A subgroup N of G is **normal**, written $N \trianglelefteq G$, if it satisfies any of these equal properties:

- (N1) N is the kernel of some group homomorphism ϕ ,
- (N2) N is stable under conjugations ($\forall n \in N$ and $g \in G$, $gng^{-1} \in N$),
- (N3) for all $g \in G$ $gN = Ng$.

Proof of equivalence. (N1 \implies N2): $\phi(gng^{-1}) = \phi(g)\phi(n)\phi(g^{-1}) = \phi(g)\phi(g)^{-1} = e_H$.

(N2 \implies N3): $gng^{-1} \in N \implies gn \in Ng$ by g^{-1} so $gN \subseteq Ng$, similarly for $Ng \subseteq gN$ with g^{-1} replacing g .

(N3 \implies N2): The set of left and right cosets of G by N are isomorphic with N as the kernel. \square

1.3 Quotient groups

Definition 1.3.1 (Quotient groups). Let $N \trianglelefteq G$, the **quotient group** of G modulo N , written G/N , is the group with elements as left cosets of N in G with $(g_1N) \cdot (g_2N) = (g_1g_2N)$.

Proof. One can easily check this satisfies all of the group axioms. \square

Remark 1.3.2. By Lagrange's theorem $|G/N| = |G|/|N|$.

Definition 1.3.3 (Simple group). A group G is **simple** if it has no normal subgroups except $\{e_G\}$ and G .

1.4 Isomorphism theorems

Theorem 1.4.1 (First isomorphism theorem). If $f : G \rightarrow H$ is a group homomorphism, $G/\ker f \cong \text{im } f$.

Proof. Have $\phi : G/\ker f \rightarrow \text{im } f$ with $\phi : g\ker f \mapsto f(g)$.

well defined: if $g\ker f = h\ker f$, $gh^{-1}\ker f = \ker f \implies f(g) = f(gh^{-1}h) = f(gh^{-1})f(h) = f(h)$.

homomorphism: $\phi((g\ker f)(h\ker f)) = \phi(gh\ker f) = f(gh) = f(g)f(h) = \phi(g\ker f)\phi(h\ker f)$.

surjective: any $h = f(g) \in \text{im } f$ is clearly $\phi(g\ker f)$ for any $g \in G$.

injective: if $\phi(g\ker f) = e_H$, $f(g) = e_H \implies g \in \ker f$ so $\ker f = \{\ker \phi\} = \{e_{G/\ker \phi}\}$. By a lemma from *Linear algebra and groups*, we now have ϕ injective. \square

Theorem 1.4.2 (Universal property of quotients). Let $N \trianglelefteq G$ and $f : G \rightarrow H$ be a group homomorphism such that $N \subseteq \ker f$. There exists a *unique* homomorphism $\tilde{f} : G/N \rightarrow H$ such that the diagram

$$\begin{array}{ccc} G & & \\ \pi \downarrow & \searrow f & \\ G/N & \xrightarrow{\tilde{f}} & H \end{array}$$

commutes, (here $\pi : G \rightarrow G/N$ is the projection map with $\pi : g \rightarrow gN$).

Proof. The proof is essentially that of Theorem 1.4.1 with $H = \text{im } f$. □

Lemma 1.4.3. If $N \trianglelefteq G$ and $N \leq H \leq G$ then $N \trianglelefteq H$.

Proof. $gN = Ng$ for all $g \in G$ so also for all $g \in H$. □

Theorem 1.4.4 (Second isomorphism theorem). Let $K, L \trianglelefteq G$ with $K \leq L$, $G/L \cong (G/K)/(L/K)$

Proof. Have $f : G/K \rightarrow G/L$, via same arguments in Theorem 1.4.1, f is a surjective group homomorphism, $gK \in \ker f \implies f(gK) = gL = L$ so $g \in L$ and $\ker f = L/K$. By Theorem 1.4.1, $(G/K)/(\ker f) = (G/K)/(L/K) \cong (G/L)$. □

Definition 1.4.5 (Frobenius product). Given $A, B \subseteq G$ a group, the **(Frobenius) product** of A and B is


$$AB := \{ab \in G : a \in A, b \in B\}.$$

Lemma 1.4.6. Given $H, N \leq G$ a group, N is normal $\implies HN \leq G$ and N, H normal $\implies HN \trianglelefteq G$.

Proof. 1. HN is nonempty with $(h_1n_1)(h_2n_2) = (n_1n_3)(h_1h_2) \in NH$ for some $n_3 \in N$ and $(hn)^{-1} = n^{-1}h^{-1} \in Nh^{-1} = h^{-1}N \subseteq HN$. □

2. $gHNg^{-1} = gHg^{-1} \cdot gNg^{-1} = HN$. □

Theorem 1.4.7 (Third isomorphism theorem). If $H \leq G$ and $N \trianglelefteq G$, $H/(H \cap N) \cong (HN)/N$. This is ometimes called the *diamond theorem* due to the shape of the subgroup lattice it produces:



where arrows point to subgroups.

Proof. Have $\phi : H \rightarrow G/N$ be the canonical map, $\ker \phi = H \cap N$ as $hN = N$ iff $h \in N$, $\text{im } \phi = \{hN : h \in H\} = HN/N$, Theorem 1.4.1 on ϕ gives the result. □

Note 1.4.8. The naming of the group isomorphism theorems throughout literatue is very inconsistent.

1.5 Centres

Definition 1.5.1 (Inner automorphisms). Given the group G the conjugations by elements of G form the group $\text{Inn } G \trianglelefteq \text{Aut } G$.

Proof. Have $\phi : G \rightarrow \text{Aut}(G)$ assigning to each element in $g \in G$ the conjugation map by G , $\text{Inn}(G) = \text{im } \phi \subset \text{Aut}(G)$. \square

Definition 1.5.2 (Centre of group). Given the group G the elements of G that commute with all other elements form the **centre** of G , $Z(G) \trianglelefteq G$.

Proof of normality. Have $\phi : G \rightarrow \text{Aut } G$ with $\phi : g \mapsto \text{conjugation by } g$, $\ker \phi = Z(G)$. \square

Proposition 1.5.3. If $G/Z(G)$ is cyclic, G is Abelian.

Proof. $G/Z(G) = \langle aZ(G) \rangle$ for some $a \in G$, for all $g \in G$ $gZ(G) = [aZ(G)]^m = a^m Z(G)$ for some $m \in \mathbb{N}$ therefore $a^{-m}g = z \in Z(G)$ so $g = a^m z$ and for all $g, h \in G$ we have $gh = a^n z_g a^m z_h = a^{n+m} z_g z_h = a^m z_h a^n z_g = hg$. \square

1.6 Commutators

Definition 1.6.1 (Commutator). For $a, b \in G$ a group, we have $[a, b] := aba^{-1}b^{-1}$ the **commutator** of a and b . $[G, G]$ is the smallest subgroup of G containing all commutators of elements of G , called the **commutator** of G .

Remark 1.6.2. A group G is Abelian iff $[G, G] = e_G$.

Theorem 1.6.3. Given G a group, $[G, G] \trianglelefteq G$ with its quotient in G Abelian.

Theorem 1.6.4. Let $N \trianglelefteq G$, G/N is Abelian iff $[G, G] \subseteq N$.

Theorem 1.6.5. Given a group G with $A, B \trianglelefteq G$, $A \cap B = \{e_G\}$ and $AB = G$; $A \times B \cong G$.

1.7 Torsion and p -primary subgroups

Definition 1.7.1 (Torsion subgroup). Given an abelian group G , the set of elements of G with finite order form the **torsion subgroup** of G , denoted G_{tors} . When $G = G_{\text{tors}}$, we call G a **torsion Abelian group**.

Definition 1.7.2 (p -primary subgroups). Given an abelian group G , the set of elements of G with order p (a prime) is the **p -primary subgroup** of G , written $G\{p\}$. When $G = G\{p\}$, we call G a **p -primary torsion Abelian group**.

Theorem 1.7.3. Let the prime factorisation of $n \in \mathbb{N}$ be $p_1^{a_1} p_2^{a_2} \dots p_m^{a_m}$ with C_n the cyclic group of order n .

$$C_n \cong C_{p_1^{a_1}} \times C_{p_2^{a_2}} \times \dots \times C_{p_m^{a_m}}.$$

Proof. \square

1.8 Generators

Lemma 1.8.1. Given an indexing set \mathcal{I} , and a sequence of subgroups $(H_i)_{i \in \mathcal{I}} \leq H$, $\bigcap_{i \in \mathcal{I}} H_i \leq G$.

Definition 1.8.2 (Subgroup generated by a set). Given $S \subseteq G$ a group,

$$\langle S \rangle := \left(\bigcap_{S \subseteq H \leq G} H \right) \leq G$$

is the **subgroup of G generated by S** . If $\langle S \rangle = G$ then we say S **generates G** and G is **finitely generated** if S is finite.

1.9 Classification of finitely generated Abelian groups

Definition 1.9.1 (Free Abelian group of rank n). The **Free Abelian group of rank n** is the group \mathbb{Z}^n under addition. The free abelian group of rank 0 is the trivial group.

Lemma 1.9.2. If $\mathbb{Z}^m \cong \mathbb{Z}^n$ then $n = m$, so the rank of a free abelian group is well defined.

Lemma 1.9.3. Any subgroup of \mathbb{Z}^n is isomorphic to some \mathbb{Z}^m for some $m \leq n$.

Theorem 1.9.4. Every finitely generated Abelian group is isomorphic to a product of finitely many cyclic groups.

Theorem 1.9.5. Every finitely generated Abelian group is isomorphic to a product of finitely many infinite cyclic groups and finitely many cyclic groups of prime order. The number of infinite cyclic factors and the number of cyclic factors of order p^r , where p is prime and $r \in \mathbb{N}$ is determined solely by the group.

Theorem 1.9.6. A finitely generated Abelian group, G , is not cyclic iff there exists a prime p such that $G \cong C_p \times C_p$.

2 Group actions

2.1 Actions

Definition 2.1.1 (Actions). Given a group G and a set X , a **group action** is: a binary operation

$$\begin{aligned} \cdot : G \times X &\longrightarrow X \\ (g, x) &\longmapsto g \cdot x \end{aligned}$$

with $e_G \cdot x = x$ for all $x \in X$ and $(g_1 g_2) \cdot x = g_1 \cdot (g_2 x)$ for all $g_1, g_2 \in G$ and $x \in X$; or, equivalently, a homomorphism $\rho : G \rightarrow \text{Sym}(X)$.

Definition 2.1.2 (Faithful set). An action of a group G on a set X is **faithful** if the map $\rho : G \rightarrow \text{Sym}(X)$ is injective.

2.2 Orbit-stabiliser theorem

Definition 2.2.1 (Orbit). Given a group G acting on a set X , the **G -orbit** of $x \in X$ is

$$G(x) := \{g \cdot x : g \in G\} \subseteq X.$$

Orbits partition X into X/G .

Definition 2.2.2 (Stabiliser). Given a group G acting on a set X , the **stabiliser** of $x \in X$ is

$$\text{Stab}_G(x) := \{g \in G : g \cdot x = x\} \subseteq G.$$

Stabilisers also partition G .

Remark 2.2.3 (Conjugacy classes). When G acts on itself by conjugations, orbits of G are the **conjugacy classes**, x^G of G and the stabilisers of G are the centralisers of G .

Lemma 2.2.4. Given a group G acting on a set X , $\text{Stab}_G(g \cdot x) = g \text{Stab}_G(x) g^{-1}$

Theorem 2.2.5 (Orbit-stabiliser theorem). Given a group G acting on a set X . For all $x \in X$, we have $\phi_x : G/\text{Stab}(x) \xrightarrow{\sim} G(x)$ by $\phi_x : g \text{Stab}(x) \mapsto g \cdot x$, giving $|G(x)| = [G : \text{Stab}(x)] = |G|/|\text{Stab}(x)|$.

Proof. asdfs □

Corollary 2.2.6. $|X| = \sum_{i=1}^n |G(x_i)| = \sum_{i=1}^n [G : \text{Stab}(x_i)]$.

Corollary 2.2.7 (Cayley's theorem). Let G be a finite group of order n . Then $S_n \cong \text{Sym}(G)$ contains a finite subgroup isomorphic to G .

Corollary 2.2.8 (Cauchy's theorem). Let G be a finite group of order n and let p be a prime factor of n . Then G contains an element of order p .

Definition 2.2.9 (p -group). A finite group G is a **p -group** if the order of G is a power of prime p .

Theorem 2.2.10. Let G be a p -group, $Z(G) \neq \{e_G\}$.

Proof. □

2.3 Jordan's theorem

Definition 2.3.1 (Transitive action). Given a group G acting on a set X , if X is a G -orbit then we say G acts **transitively** on X .

Definition 2.3.2 (Fixed points). Given a group G acting on a set X , an element $x \in X$ is a fixed point of $g \in G$ iff $g \cdot x = x$. We have $\text{Fix}(g) \subseteq X$ the set of fixed points of $g \in G$ satisfying:

$$\text{Stab}(x) \xleftarrow{\pi_G} \{(x, g) \in X \times G; g \cdot x = x\} \xrightarrow{\pi_X} \text{Fix}(g) .$$

Theorem 2.3.3 (Jordan's theorem). Let G act transitively on a finite set X , we have

$$\sum_{g \in G} |\text{Fix}(g)| = |G|,$$

with there being some element $g \in G$ such that $\text{Fix}(g) = \emptyset$.

Corollary 2.3.4 (Burnside's lemma). Given a group G acting on a finite set X :

$$|X/G| = \frac{1}{|G|} \sum_{g \in G} |\text{Fix}(g)|.$$

3 Rings

3.1 Rings

Definition 3.1.1 (Ring). A ring (with 1) is a set R with elements $0, 1$ and binary operations $+, \times$ such that

1. $(R, +)$ is an abelian group with identity 0 ,
2. (R, \times) is a semigroup with 1 as the identity,
3. both left and right multiplication are distributive over addition.

Examples 3.1.2. $\mathbb{Z}, \mathbb{Q}, \mathbb{R}, \mathbb{C}$ are all rings with their normal operations. $\mathbb{R}[x]$ is the set of real-valued polynomials and is also a ring.

Definition 3.1.3 (Subring). A subset of a ring which is itself a ring under the same operators with the same 1 is a **subring**.

Definition 3.1.4 (Commutative ring). A ring, R , is **commutative** iff $a + b = b + a$ for all $a, b \in R$.

Definition 3.1.5 (Invertible). An element x of a ring R is invertible if there exists $y, z \in R$ with $yx = zx = 1$.

Definition 3.1.6 (Division ring). A ring R is called a **division ring** if $R \setminus \{0\}$ is a group under multiplication with identity 1 .

Remark 3.1.7. A commutative division ring is a field.

Definition 3.1.8 (Integral domain). A commutative ring R is an integral domain iff $0 \neq 1$ and for all $a, b \in R$ $ab = 0 \implies a = 0$ or $b = 0$.

3.2 Ring homomorphisms

Definition 3.2.1 (Ring homomorphism). Let R, S be rings, a function $f : R \rightarrow S$ is a **ring homomorphism** iff it satisfies

1. $f : (R, +) \rightarrow (S, +)$ is a group homomorphism,
2. $f(xy) = f(x)f(y)$ for all $x, y \in R$,
3. $f(1_R) = 1_S$.

Lemma 3.2.2. Given the ring homomorphism $f : R \rightarrow S$ the kernel of f is a subgroup of $(R, +)$ which satisfies $xr, rx \in \ker f$ for all $x \in \ker f$ and $r \in R$.

3.3 Ideals

Definition 3.3.1 (Ideal). For a ring R , a subset $I \subseteq R$ is a **left ideal**, denoted $I \trianglelefteq R$ iff

1. $(I, +)$ is a subgroups of $(R, +)$,
2. if $r \in R$ and $i \in I$, $ri \in I$.

Similarly, for **right ideals**. A subset I is a bi-ideal if it is both a left and right ideal.

Definition 3.3.2 (Quotient ring). Given ring R with proper ideal $I \subset R$, The quotient abelian group R/I , with natural multiplication, forms the **quotient ring** of R by I .

Definition 3.3.3 (Principal ideal). Given a commutative ring R and some $a \in R$, $aR := \{ax : x \in R\}$ is an ideal called a **principal ideal** with **generator** a .

Definition 3.3.4. A bijective ring homomorphism is a **ring isomorphism**, a ring homomorphism $f : R \rightarrow R$ is a **ring endomorphism**, an isomorphic ring endomorphism is **ring automorphism**.

Proposition 3.3.5. Given the ring homomorphism $f : R \rightarrow S$, $f(R) = \text{im } R$ is a subring of S which is isomorphic to $R/\ker f$.

Proposition 3.3.6. A commutative ring is a field iff its only proper ideal is the trivial / zero ideal.

Proposition 3.3.7. Given $f : R \rightarrow S$ a ring homomorphism with J a left (or right or bi) ideal of S , $f^{-1}(J)$ is a left (respectively) ideal of R .

Definition 3.3.8 (Prime ideal). Let R be a commutative ring, a proper ideal $I \subset R$ is a **prime ideal** iff $ab \in I$ for $a, b \in R \implies a \in I$ or $b \in I$.

Theorem 3.3.9. If $I \subset R$ is a prime ideal, R/I is an integral domain

Definition 3.3.10 (Maximal ideal). A proper ideal I in a commutative ring R is **maximal** iff there are no other proper ideals J with $I \subset J$.

Theorem 3.3.11. I is a maximal ideal of R iff R/I is a field.

4 Integral domains

Throughout this section we will always have R be an integral domain.

4.1 Integral domains

Theorem 4.1.1. $ab = ac \implies b = c$ for all $a, b, c \in R$. (the cancellation law holds for all integral domains)

Proposition 4.1.2. For $a, b \in R$, $aR = bR$ iff $a = br$ for some $r \neq 0 \in R$.

Proof.

□

Theorem 4.1.3. All fields are integral domains and all finite integral domains are fields.

Remark 4.1.4. The ring $\mathbb{Z}/n\mathbb{Z}$ is an integral domain iff it is a field $\iff n$ is prime.

Definition 4.1.5 (Unit). $r \in R$ is a **unit** if there exists some $y \in R$ with $x \times y = 1_R$. We write R^\times for the group of units in R under multiplication.

Definition 4.1.6 (Irreducible). $r \in R \setminus R^\times$ is **irreducible** if it cannot be written as the product of two elements of $R \setminus R^\times$.

4.2 Characteristic

Lemma 4.2.1. For any ring S there is a unique ring homomorphism $f : \mathbb{Z} \rightarrow S$.

Proof. Have $f(0_R) = 0$, $f(1) \rightarrow 1_S$ and inductively have $f(n)$ be the sum of 1_S n times.

□

Lemma 4.2.2. The kernel of the unique homomorphism $\mathbb{Z} \rightarrow R$ is either $\{0\}$ or $p\mathbb{Z}$ for some prime p .

Definition 4.2.3 (Characteristic). The **characteristic** of R is the unique non-negative generator of the kernel of $\mathbb{Z} \rightarrow R$, denoted $\text{char } R$.

4.3 Polynomial rings

Definition 4.3.1 (Polynomial ring). $R[t]$ is, formally, the set of infinite sequences of elements of R with finitely many non-zero terms, but more helpfully: the set of polynomials in t with coefficients in R .

Definition 4.3.2 (Polynomial degree). The **degree** of a polynomial, $r_0 + r_1t + r_2t^2 + \dots + r_it^i + \dots \in R[t]$, is the unique maximum $i \in \mathbb{N}$ with $r_i \neq 0$ and 0 otherwise.

Lemma 4.3.3. Given $p(t), q(t) \in R$, $\deg(p(t)q(t)) = \deg(p(t)) + \deg(q(t))$, $R[t]$ is an integral domain and $R[t]^* = R^*$.

Theorem 4.3.4. If k is a field with $a(t), b(t) \in k[t]$ with $b(t) \neq 0$, there exists $q(t), r(t) \in k[t]$ such that $a(t) = q(t)b(t) + r(t)$ with $\deg(r(t)) < \deg(b(t))$ and $q(t), r(t)$ unique.

5 PIDs and UFDs

5.1 Euclidian domains

Definition 5.1.1 (Euclidian domain). An integral domain R is a Euclidian domain if there exists some $\phi : R^* \rightarrow \mathbb{N}_0$ satisfying:

1. $\phi(ab) \leq \phi(a)$ for all $a, b \neq 0$,
2. for all $a, b \in R$ there exists $q, r \in R$ with $a = qb + r$ with $r = 0$ or $\phi(r) < \phi(b)$.

5.2 Principal ideal domains

Definition 5.2.1 (Principal integral domain). An integral domain R is a **principal integral domain** iff every ideal of R is principal.

Theorem 5.2.2. R is a Euclidian domain $\implies R$ is a principal integral domain.

Proof.

□

Corollary 5.2.3. F is a field $\implies F[t]$ is a PID.

5.3 Unique factorisation domains

Definition 5.3.1 (Unique factorisation domain). An integral domain R is a **unique factorisation domain** iff every element of $R \setminus R^\times$ can be written as the product of a single unit and finitely many irreducibles in R which is unique up to rearrangement.

Definition 5.3.2 (Division). Given a, b in the integral domain R , we say a **divides** b , written $a|b$ iff $b = ra$ for some $r \in R$ and **properly divides** if $r \notin R^\times$.

Lemma 5.3.3. Given $p, a, b \in R$ a UFD, if p is irreducible then $p|ab \implies p|a$ or $p|b$.

Lemma 5.3.4. There is no infinite sequence of non-zero $r_1, r_2, \dots \in R$ a UFD such that r_{n+1} properly divides r_n for all $n \geq 1$.

Theorem 5.3.5. The integral domain R is a UFD iff the properties in Lemma 5.3.3 and Lemma 5.3.4 hold.

Theorem 5.3.6. Every principal ideal domain is a unique factorisation domain.

6 Fields

6.1 Vector spaces

Throughout this section let k be a field.

Definition 6.1.1 (Vector space). A k -vector space V is an abelian group with an action of k on the elements of V satisfying

1. $1_kv = v$ for all $v \in V$,
2. $(x + y)V = xV + yV$ for all $x, y \in k$ and $v \in V$,

3. $x(v + w) = xv + xw$ for all $x \in k$ and $v, w \in V$.

Proposition 6.1.2. If $\text{ch } k = 0$ then k contains a unique subfield isomorphic to \mathbb{Q} . Otherwise, if $\text{ch } k = p$ then k contains a unique subfield isomorphic to \mathbb{F}_p .

Theorem 6.1.3. Every finite field has p^n elements for some prime p and $n \in \mathbb{N}$.

6.2 Field extensions

Definition 6.2.1 (Field extension). A **field extension** F of k is a k -vector space.

Proposition 6.2.2. All homomorphisms between fields and rings are injective.

Proof. The only possible maps between fields are field extensions, the only proper ideal of a field is the zero ideal. \square

Definition 6.2.3 (Finite field extension). An extension of the fields $k \subset K$ is **finite** iff K is a finite dimensional vector space over k with $\dim K$ the **degree** of the extension

Theorem 6.2.4. If $k \subset F \subset K$ are field extensions, K is a finite extension of k iff K is a finite extension of F and F is a finite extension of k . We then have $[K : k] = [K : F][F : k]$.

Remark 6.2.5. Degree 2 and 3 field extensions are called quadratics and cubics respectively.

6.3 Constructing fields

Lemma 6.3.1. Given R a PID with $a \neq 0 \in R$, aR is maximal iff a is irreducible.

Proof. \square

Corollary 6.3.2. Given R a PID with reducible $a \in R$, R/aR is a field.

Theorem 6.3.3. A polynomial $f(t) \in k[t]$ of degree 2 or 3 is irreducible iff it has no root in k .

Definition 6.3.4 (Non-Square). $a \in k$ is non-square if there is no element $b \in k$ with $b^2 = a$.

Lemma 6.3.5. Let p be an odd prime. The field \mathbb{F}_p contains $(p-1)/2$ non-squares. For all non-square $a \in \mathbb{F}_p$, $t^2 - a$ is irreducible in $\mathbb{F}_p[t]$.

Theorem 6.3.6. For all $p(t) \in k[t]$, there exists a finite field extension $k \subset K$ such that:

$$p(t) = c \prod_{i=1}^n (t - a_i),$$

for some $c \in k^\times$ and $a_i \in K$ for all $i \in [1, n]$.

6.4 Existence of finite fields

Theorem 6.4.1. Let k have characteristic $p \neq 0$, for all $x, y \in k$ and $m \in \mathbb{Z}^{\geq 0}$,

$$(x + y)^{p^m} = x^{p^m} + y^{p^m}.$$

Definition 6.4.2 (Derivative). Let $p(t) = a_0 + a_1 t + \dots + a_n t^n \in k[t]$, the **derivative** of $p(t)$ is

$$p'(t) := a_1 + 2a_2 t + \dots + na_n t^{n-1}.$$

Lemma 6.4.3. Let $p(t) = (x - a_1)(x - a_2) \dots (x - a_n) \in k[t]$, $a_i \neq a_j$ for all $i \neq j$ iff $p(t)$ and $p'(t)$ have no common roots.

Theorem 6.4.4. For all prime p and natural n , there exists a field with p^n elements.

Chapter 2

Lebesgue Measure and Integration

Lectured by Someone
Typed by Yu Coughlin
Season Year

Introduction

The following are complementary reading for the course.

- G. Grimmett and D. J. A. Welsh, Probability: An Introduction, 1986
- J. K. Blitzstein and J. Hwang, Introduction to Probability, 2019
- D. F. Anderson et al, Introduction to Probability, 2018
- S. M. Ross, Introduction to Probability Models, 2014
- G. Grimmett and D. Stirzaker, Probability and Random Processes, 2001
- G. Grimmett and D. Stirzaker, One Thousand Exercises in Probability, 2009

Lecture 1
Monday
30/10/2023

1 Motivation

2 Measures

2.1 Algebras and σ -algebras

2.2 Measures

2.3 Complete measure spaces

3 Constructing measures

3.1 Pre-measure

3.2 Outer measure

3.3 Restriction

3.4 Lebesgue measure

4 Measurable functions

4.1 Definition

4.2 Properties

4.3 Continuity

5 Lebesgue integral

5.1 Construction

5.2 Properties

6 Convergence

6.1 Monotone convergence

6.2 Fatou's lemma

6.3 Lebesgue dominated convergence

6.4 Vitali's theorem

7 L^p spaces

7.1 Norms

7.2 L^p spaces

7.3 Normed vector spaces

7.4 Completeness

8 Product measures

8.1 Products of sets

8.2 σ -algebras on product sets

8.3 Product measures

9 Fubini's theorem

9.1 Motivations

9.2 Setup

9.3 Fubini's theorem

10 Differentiation

Chapter 3

Categories

Lectured by noone
Typed by Yu Coughlin
Season Year

Introduction

The following are complementary reading for the course.

- G. Grimmett and D. J. A. Welsh, Probability: An Introduction, 1986
- J. K. Blitzstein and J. Hwang, Introduction to Probability, 2019
- D. F. Anderson et al, Introduction to Probability, 2018
- S. M. Ross, Introduction to Probability Models, 2014
- G. Grimmett and D. Stirzaker, Probability and Random Processes, 2001
- G. Grimmett and D. Stirzaker, One Thousand Exercises in Probability, 2009

1 Basic definitions

1.1 Categories

Definition 1.1.1 (Category). A category \mathcal{C} contains the following data:

1. a *collection* of objects, $\text{Ob}(\mathcal{C})$,
2. for every $x, y \in \text{Ob}(\mathcal{C})$ a collection of morphisms $\text{Hom}_{\mathcal{C}}(x, y)$ from x to y ,
3. an identity morphism $\text{id}_x \in \text{Hom}_{\mathcal{C}}(x, x)$ for all $x \in \text{Ob}(\mathcal{C})$,
4. a composition map of morphisms, $\circ : \text{Hom}_{\mathcal{C}}(y, z) \times \text{Hom}_{\mathcal{C}}(x, y) \rightarrow \text{Hom}_{\mathcal{C}}(x, z)$ for all $x, y, z \in \text{Ob}(\mathcal{C})$.

Which satisfy the two axioms:

1. for all $f \in \text{Hom}_{\mathcal{C}}(x, y)$ with $x, y \in \text{Ob}(\mathcal{C})$ we have $f \circ \text{id}_x = f = \text{id}_y \circ f$,
2. for compatible morphisms f, g, h we have $f \circ (g \circ h) = (f \circ g) \circ h$.

We will use the shorthand $x \in \mathcal{C}$ for $x \in \text{Ob} \mathcal{C}$, $\text{Hom}(x, y)$ for $\text{Hom}_{\mathcal{C}}(x, y)$ when \mathcal{C} is obvious and $\text{End}(x)$ for $\text{Hom}(x, x)$.

Note 1.1.2. Note that in our definition the term *collection* is used instead of set, this is commonplace and necessary to prevent paradoxes when constructing the category of sets.

Examples 1.1.3. The following are all categories:

1. **Set** with sets as objects and functions as their morphisms,
2. **Grp** with groups as objects and their homomorphisms as morphisms,
3. **Ab**, **Grp** restricted to abelian groups,
4. for a field k , **Vect_k** with k -vector spaces as objects and linear transformations as morphisms,
5. **Cat** with categories as objects and soon to be defined **functors** as morphisms,
6. **Top**, **Rng**, **Meas**, **Poset**, **Man** with their objects and morphisms all defined similarly
7. Given a category \mathcal{C} , \mathcal{C}^{op} which has the same objects as \mathcal{C} but $\text{Hom}_{\mathcal{C}^{op}}(x, y) = \text{Hom}_{\mathcal{C}}(y, x)$ for all $x, y \in \mathcal{C}$,
8. Any set X with objects as elements in X and no morphisms except the identities
9. (\mathbb{R}, \leq) with objects as \mathbb{R} and a morphisms from x to y iff $x \leq y$ for all $x, y \in \mathbb{R}$.

Definition 1.1.4 (Isomorphism). A morphism $f \in \text{Hom}(x, y)$ is an **isomorphism** iff there is a morphism $f^{-1} \in \text{Hom}(y, x)$ with $f \circ f^{-1} = \text{id}_y$ and $f^{-1} \circ f = \text{id}_x$.

1.2 Functors

Definition 1.2.1 ((Covariant) Functor). Given categories \mathcal{C}, \mathcal{D} a **(covariant) functor** $F : \mathcal{C} \rightarrow \mathcal{D}$ is the following data:

1. a map $\text{Ob}(\mathcal{C}) \rightarrow \text{Ob}(\mathcal{D})$ (also denoted F),
2. for any two objects $x, y \in \mathcal{C}$ a map $\text{Hom}_{\mathcal{C}}(x, y) \rightarrow \text{Hom}_{\mathcal{D}}(F(x), F(y))$ (also also denoted F)

satisfying the properties:

1. for all $x \in \mathcal{C}$, $F(\text{id}_x) = \text{id}_{F(x)}$,
2. for all x, y, z with f, g in $\text{Hom}_{\mathcal{C}}(y, z), \text{Hom}_{\mathcal{C}}(x, y)$, $F(f \circ g) = F(f) \circ F(g)$.

Definition 1.2.2 (Contravariant functor). A **contravariant functor** from \mathcal{C} to \mathcal{D} is a covariant functor from \mathcal{C}^{op} to \mathcal{D} .

Definition 1.2.3 (Hom-functor). The **hom-functor** for a given category \mathcal{C} is $\text{Hom}_{\mathcal{C}} : \mathcal{C}^{op} \times \mathcal{C} \rightarrow \text{Set}$ sending a pair of elements $c, d \in \mathcal{C}$ to $\text{Hom}_{\mathcal{C}}(c, d)$.

1.3 Natural transformations

Definition 1.3.1 (Natural transformation). Given categories \mathcal{C}, \mathcal{D} with functors $F, G : \mathcal{C} \rightarrow \mathcal{D}$, a **natural transformation** $\eta : F \rightarrow G$ consists of morphisms η_x for all $x \in \mathcal{C}$ such that the diagram,

$$\begin{array}{ccc} F(x) & \xrightarrow{F(f)} & F(y) \\ \downarrow \eta_x & & \downarrow \eta_y \\ G(x) & \xrightarrow{G(f)} & G(y) \end{array}$$

commutes for all $x, y \in \mathcal{C}$ and $f \in \text{Hom}_{\mathcal{C}}(x, y)$.

Remark 1.3.2. By constructing the category of functors from \mathcal{C} to \mathcal{D} , denoted $\text{Fun}(\mathcal{C}, \mathcal{D})$, morphisms are natural transformations. **Natural isomorphisms** are defined as isomorphisms in this category.

1.4 Equivalence of categories

Definition 1.4.1 (Equivalence). Given categories \mathcal{C}, \mathcal{D} an **equivalence of categories** is a pair of functors $F : \mathcal{C} \rightarrow \mathcal{D}$ and $G : \mathcal{D} \rightarrow \mathcal{C}$ with natural isomorphisms $FG \xrightarrow{\sim} \text{id}_{\mathcal{D}}$ and $\text{id}_{\mathcal{C}} \xrightarrow{\sim} GF$.

Definition 1.4.2 (Adjunction). An **adjunction** between categories \mathcal{C}, \mathcal{D} is a pair of functors $F : \mathcal{C} \rightarrow \mathcal{D}$ and $G : \mathcal{D} \rightarrow \mathcal{C}$ such that for all $x \in \mathcal{C}$ and $y \in \mathcal{D}$, there exists an $\eta_{x,y} : \text{Hom}_{\mathcal{C}}(x, G(y)) \xrightarrow{\sim} \text{Hom}_{\mathcal{D}}(F(x), y)$ such that the diagram

$$\begin{array}{ccccc} \text{Hom}_{\mathcal{D}}(F(x'), y) & \xrightarrow{\circ F(f)} & \text{Hom}_{\mathcal{D}}(F(x), y) & \xrightarrow{g \circ} & \text{Hom}_{\mathcal{D}}(F(x), y') \\ \uparrow \eta_{x',y} & & \uparrow \eta_{x,y} & & \uparrow \eta_{x,y'} \\ \text{Hom}_{\mathcal{C}}(x', G(y)) & \xrightarrow{\circ f} & \text{Hom}_{\mathcal{C}}(x, G(y)) & \xrightarrow{G(g) \circ} & \text{Hom}_{\mathcal{C}}(x, G(y')) \end{array}$$

commutes for all $x, x' \in \mathcal{C}$; $y, y' \in \mathcal{D}$; $f : x \rightarrow x'$ and $g : y \rightarrow y'$.

Theorem 1.4.3. If F, G form an equivalence of the categories \mathcal{C}, \mathcal{D} then F, G are an adjunction.

Examples 1.4.4 (Adjunctions in group theory). Consider the **forgetful functor** $F : \text{Ab} \rightarrow \text{Grp}$ which simply forgets the Abelian property of a group. We also have the **abelianisation functor** $(-)^{\text{ab}} : \text{Grp} \rightarrow \text{Ab}$ which maps $G \mapsto G^{\text{ab}} := G/[G, G]$. F and $(-)^{\text{ab}}$ form an adjunction between Grp and Ab .

1.5 Representable functors

Definition 1.5.1 (Yoneda functor). Given some x in a category \mathcal{C} , there is a functor $\text{Hom}_{\mathcal{C}}(-, x) : \mathcal{C}^{\text{op}} \rightarrow \text{Set}$ which satisfies the required properties to have the **Yoneda functor**:

$$Y : \mathcal{C} \rightarrow \text{Fun}(\mathcal{C}^{\text{op}}, \text{Set}).$$

Which sends an element $y \in \mathcal{C}$ to the functor from objects in \mathcal{C}^{op} to the set of morphisms from these objects to y .

Lemma 1.5.2. The Yoneda functor and the hom-functor form an adjunction in Cat .

Definition 1.5.3 (Representable). A functor $F \in \text{Fun}(\mathcal{C}^{\text{op}}, \text{Set})$ is **representable** if $F \cong Y(c)$ for some $c \in \mathcal{C}$.

Example 1.5.4. Consider the functor $F : \text{Set}^{(\text{op})} \rightarrow \text{Set}$ sending a set to its powerset. F is clearly isomorphic to the functor $\text{Hom}(-, \{0, 1\})$ from subsets to indicator functions on X . This is the image of the Yoneda functor so F is representable.

1.6 Yoneda lemma

Theorem 1.6.1 (Yoneda lemma). Given some $x \in \mathcal{C}$ and $F \in \text{Fun}(\mathcal{C}^{\text{op}}, \text{Set})$ we have

$$\text{Hom}_{\text{Fun}(\mathcal{C}^{\text{op}}, \text{Set})}(Y(x), F) \cong F(x).$$

Remark 1.6.2. This is a generalisation of Cayley's theorem which shows that we can study a group by instead studying the permutations of its underlying set.