## Exposures negatively associated with ovarian cancer risk
## meta−analysis of dietary−intake studies

| Exposure | # of studies | Sample size | Cases | Pooled RR (95% CI) |
|---|---|---|---|---|
| *CAROTENOIDS* | | | | |
| Lutein | 4 | 529,715 | 3,624 | *0.66 (0.45−0.98)* |
| beta−Carotene | 8 | 611,872 | 4,692 | *0.72 (0.57−0.91)* |
| *VITAMINS A, C, D, E, K* | | | | |
| Vitamin D | 2 | 4,611 | 1,993 | *0.52 (0.38−0.71)* |
| Vitamin E | 6 | 591,029 | 4,497 | *0.66 (0.48−0.92)* |
| Vitamin C | 3 | 583,089 | 2,528 | *0.71 (0.38−1.29)* |
| Vitamin A | 6 | 1,588,466 | 9,351 | *0.97 (0.86−1.09)* |
| *B VITAMINS* | | | | |
| Folic Acid | 8 | 723,200 | 3,495 | *0.76 (0.56−1.02)* |
| *ANTIOXIDANTS* | | | | |
| Antioxidants | 4 | 523,871 | 2,791 | *0.59 (0.36−0.99)* |
| *MINERALS & TRACE ELEMENTS* | | | | |
| Selenium | 2 | 1,633 | 611 | *0.58 (0.38−0.9)* |
| Calcium | 3 | 36,513 | 1,667 | *0.66 (0.54−0.81)* |
| *POLYPHENOLS & FLAVONOIDS* | | | | |
| Tea | 8 | 307,213 | 4,002 | *0.77 (0.55−1.08)* |
| *FRUITS & VEGETABLES* | | | | |
| Legumes | 2 | 63,479 | 506 | *0.58 (0.22−1.49)* |
| *FATTY ACIDS & LIPIDS* | | | | |
| Omega−6 FA | 2 | 6,630 | 3,238 | *0.78 (0.67−0.9)* |
| *PHYTOESTROGENS* | | | | |
| Soy | 2 | 1,906 | 754 | *0.37 (0.22−0.64)* |
| *OTHER* | | | | |
| Alcohol | 18 | 819,111 | 17,098 | *0.95 (0.82−1.11)* |
| Caffeine | 11 | 336,054 | 38,479 | *0.95 (0.79−1.15)* |




**Exposure group**

Antioxidants | B Vitamins | Carotenoids | Fatty Acids & Lipids | Fruits & Vegetables
Minerals & Trace Elements | Other | Phytoestrogens | Polyphenols & Flavonoids | Vitamins A, C, D, E, K

*RR < 1 = inversely associated with ovarian cancer risk. | [*] pooled RR (size proportional to # of studies) | [ ] red outline = statistically significant (95% CI excludes 1.0) | [*] faded = non−significant | dashed line = 95% prediction interval*