


Exposures negatively associated with breast cancer risk				
Exposure	# of studies	Sample size	Cases	Pooled RR (95% CI)
类胡萝卜素				
叶黄素	13	1,072,840	51,083	0.62 (0.52–0.74)
番茄红素	13	141,249	13,606	0.68 (0.57–0.81)
–胡萝卜素	39	1,817,569	128,023	0.73 (0.66–0.8)
维生素 A、C、D、E、K				
维生素 D	75	1,315,839	99,816	0.82 (0.79–0.86)
维生素 E	22	328,582	33,305	0.82 (0.68–0.97)
维生素 A	14	322,062	20,383	0.9 (0.74–1.09)
维生素 C	16	512,782	35,415	0.94 (0.82–1.06)
B族维生素				
维生素 B6	7	92,079	8,361	0.56 (0.42–0.76)
维生素 B12	8	88,394	11,275	0.68 (0.48–0.95)
维生素 B1	2	38,639	853	0.75 (0.61–0.93)
叶酸	37	1,078,947	52,167	0.8 (0.73–0.89)
抗氧化剂				
抗氧化剂	14	397,778	21,055	0.65 (0.53–0.8)
矿物质与微量元素				
钼	3	4,820	2,330	0.42 (0.24–0.76)
钾	2	1,370	587	0.55 (0.3–1.03)
钙	27	894,018	76,477	0.91 (0.86–0.96)
硒	19	346,098	21,359	0.95 (0.88–1.03)
多酚与黄酮类				
绿茶	14	274,406	18,211	0.89 (0.77–1.02)
水果与蔬菜				
蘑菇	5	4,428	1,957	0.59 (0.24–1.43)
大蒜	4	243,855	1,388	0.71 (0.55–0.94)
豆类	15	273,560	15,479	0.74 (0.65–0.85)
橄榄	11	249,237	17,161	0.75 (0.64–0.87)
发酵食品与益生菌				
味噌	3	87,065	751	0.8 (0.56–1.13)
酸奶	5	20,962	40,377	0.82 (0.73–0.94)
发酵食品	13	218,102	46,171	0.86 (0.69–1.07)
脂肪酸与脂质				
Omega–3 脂肪酸	21	267,959	13,978	0.67 (0.58–0.79)
鱼油	7	511,508	63,626	0.76 (0.59–0.97)
鳕鱼肝油	2	4,001	1,731	0.78 (0.68–0.9)
植物雌激素				
大豆	40	961,262	34,226	0.75 (0.65–0.86)
草本与植物制品				
黑升麻	3	31,455	5,515	0.74 (0.6–0.92)
其他				
燕麦	2	28,604	1,075	0.7 (0.3–1.65)
地中海饮食	24	639,823	40,922	0.76 (0.7–0.83)
咖啡因	11	872,194	38,461	0.97 (0.92–1.01)
乳制品	46	2,055,428	92,012	0.97 (0.92–1.03)
代谢物与氨基酸				
胆碱	4	80,262	6,805	0.76 (0.52–1.11)
甜菜碱	3	77,198	5,297	0.77 (0.56–1.07)
支链氨基酸 (BCAAs)	3	223,922	13,868	0.98 (0.87–1.1)
激素与内源性物质				
褪黑素	10	62,422	3,175	0.85 (0.73–1)



Exposure group

- 抗氧化剂

■ B族维生素

■ 类胡萝卜素

■ 脂肪酸与脂质

■ 发酵食品与益生菌
- 水果与蔬菜

■ 草本与植物制品

■ 激素与内源性物质

■ 代谢物与氨基酸

■ 矿物质与微量元素
- 其他

■ 植物雌激素

■ 多酚与黄酮类

■ 维生素 A、C、D、E、K