


Exposures inversely associated with ovarian cancer risk

Exposure	N studies	N	Cases	Pooled RR (95% CI)
CAROTENOÏDEN				
Luteïne	4	529,715	3,624	0.69 (0.48-0.98)
Bètacaroteen	9	678,322	30,201	0.87 (0.78-0.98)
VITAMINEN A, C, D, E EN K				
Vitamine E	7	657,479	30,006	0.65 (0.48-0.88)
Vitamine C	3	583,089	2,528	0.72 (0.41-1.29)
Vitamine D	13	426,489	19,570	0.81 (0.73-0.91)
Vitamine A	6	1,588,466	9,351	0.97 (0.86-1.09)
B-VITAMINEN				
Foliumzuur	9	712,037	10,452	0.87 (0.68-1.1)
MINERALEN EN SPORENELEMENTEN				
Seleen	3	1,947	768	0.23 (0.06-0.93)
Calcium	7	42,324	4,508	0.69 (0.53-0.9)
POLYFENOLEN EN FLAVONOÏDEN				
Thee	6	238,872	3,158	0.77 (0.53-1.13)
FRUIT EN GROENTEN				
Peulvruchten	2	63,479	506	0.6 (0.25-1.43)
VETZUREN EN LIPIDEN				
Omega 6-vetzuren	2	6,630	3,238	0.8 (0.69-0.91)
FYTO-OESTROGENEN				
Soja	2	1,906	754	0.41 (0.25-0.67)
OVERIG				
Alcohol	18	2,268,288	41,778	0.96 (0.84-1.09)
Cafeïne	12	380,116	39,052	0.97 (0.86-1.09)



Exposure group

- B-vitaminen
- Carotenoïden
- Vetzuren en lipiden
- Fruit en groenten
- Mineralen en sporenelementen
- Overig
- Fyto-oestrogenen
- Polyfenolen en flavonoïden
- Vitaminen A, C, D, E en K

RR < 1 = inversely associated with ovarian cancer risk. | [*] pooled RR (size proportional to k studies) | [] red outline = statistically significant (95% CI excludes 1.0) | [*] faded = non-significant | dashed line = 95% prediction interval