


Exposures positively associated with breast cancer risk
meta-analysis of dietary-intake studies

Exposure	N studies	N	Cases	Pooled RR (95% CI)
비타민 A, C, D, E, K				
비타민 K	2	5,048	2,592	1.01 (0.68-1.5)
무기질 및 미량 원소				
셀레늄	6	175,826	12,809	1.02 (0.87-1.2)
철	8	385,425	22,415	1.22 (1.04-1.43)
폴리페놀 및 플라보노이드				
차	26	1,631,244	201,438	1.03 (0.96-1.12)
과일 및 채소				
자몽	2	164,504	5,404	1.1 (0.8-1.52)
지방산 및 지질				
공액리놀레산	4	380,443	17,329	1.06 (0.92-1.22)
오메가-6 지방산	3	5,439	2,664	1.45 (1.08-1.94)
기타				
달걀	9	29,487	5,481	1.15 (0.95-1.39)
붉은 고기	34	1,178,141	60,075	1.33 (1.24-1.42)
알코올	191	16,169,285	1,285,264	1.37 (1.31-1.42)



Exposure group

-  지방산 및 지질
-  과일 및 채소
-  무기질 및 미량 원소
-  기타
-  폴리페놀 및 플라보노이드
-  비타민 A, C, D, E, K

RR > 1 = positively associated with breast cancer risk. | [*] pooled RR (size proportional to k studies) | [] red outline = statistically significant (95% CI excludes 1.0) | [*] faded = non-significant | dashed line = 95% prediction interval